

**ANTI-DIARRHEAL- loperamide hcl tablet**  
**Chain Drug Consortium**

-----  
**Premier Value 44-375-Delisted**

***Active ingredient (in each caplet)***

Loperamide HCl 2 mg

***Purpose***

Anti-diarrheal

***Use***

controls symptoms of diarrhea, including Travelers' Diarrhea

***Warnings***

**Allergy alert:** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.

**Heart alert:** Taking more than directed can cause serious heart problems or death.

**Do not use**

if you have bloody or black stool.

**Ask a doctor before use if you have**

- a fever
- mucus in the stool
- a history of liver disease
- a history of abnormal heart rhythm

**Ask a doctor or pharmacist before use if you are**

taking a prescription drug. Loperamide may interact with certain prescription drugs.

**When using this product,**

tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.

**Stop use and ask a doctor if**

- symptoms get worse
- diarrhea lasts for more than 2 days
- you get abdominal swelling or bulging. These may be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

- **drink plenty of clear fluids to help prevent dehydration caused by diarrhea**
- find right dose on chart. If possible, use weight to dose; otherwise, use age.

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

### **Other information**

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store between 20°-25°C (68°-77°F)
- see end flap for expiration date and lot number

### **Inactive ingredients**

corn starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

### **Questions or comments?**

**Call 1-800-426-9391** 8:30 AM-4:00 PM ET, Monday-Friday

### **Principal Display Panel**

\*COMPARE TO THE ACTIVE INGREDIENT IN IMODIUM® A-D

**Premier  
Value®**

## Anti-diarrheal

Loperamide HCl Tablets, 2 mg

### ANTI-DIARRHEAL

Controls the symptoms  
of diarrhea

6 Caplets

actual size

**TAMPER EVIDENT: DO NOT USE IF PACKAGE  
IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS  
ANY SIGNS OF TAMPERING**

\*This product is not manufactured or distributed by Johnson &  
Johnson Corporation, owner of the registered trademark Imodium® A-D.

50844 ORG061937545

**Distributed by:**  
**Pharmacy Value Alliance, LLC**  
**407 East Lancaster Avenue, Wayne, PA 19087**

INDEPENDENTLY TESTED

PV

SATISFACTION GUARANTEED

**If for any reason you are not satisfied with  
this product, please return it to the store  
where purchased for a full refund.**

**Drug Facts**

**KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

**Active ingredient (in each caplet)**  
Loperamide HCl 2 mg..... Anti-diarrheal

**Purpose**  
Use controls symptoms of diarrhea, including Travelers' Diarrhea

**no print / no varnish area**  
lot no. & exp. date

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS  
OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR  
SHOWS ANY SIGNS OF TAMPERING**

**Premier Value®**

**\*COMPARE TO THE ACTIVE INGREDIENT  
IN IMODIUM® A-D**

**Anti-diarrheal**  
Loperamide HCl Tablets, 2 mg  
**ANTI-DIARRHEAL**

actual size

Controls the symptoms  
of diarrhea

1. To open, tear  
at perforations and  
remove a section.

2. Use % to cut  
through plastic  
for caplet.

Manufactured or distributed by Johnson &  
Johnson, owner of the registered trademark Imodium®  
50844 ORG061937545

Alliance, LLC  
407 East Lancaster Avenue, Wayne, PA 19087

Reason you are not satisfied with  
this product, please return it to the store  
where purchased for a full refund.



6 Caplets



\*This product is not  
 Johnson Corporation  
 A-D, 50844 0  
 Distributed By:  
 Pharmacy Value  
 407 East Lancoas  
 If for any n  
 this product  
 where part

B-1590-375-45  
 OR6061937545

**Drug Facts (continued)**  
**Other information** ■ store between 20°-25°C (68°-77°F) ■ see end flap for expiration date and lot number  
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN  
**Inactive ingredients** com starch, D&C yellow #10 aluminum lake, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, magnesium stearate, microcrystalline cellulose, silicon dioxide  
**Questions or comments?** Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

**Drug Facts (continued)**  
**Warnings** Do not use if you have ever had a rash or other allergic reaction to loperamide HCl.  
 Heart alert: Taking more than directed can cause serious heart problems or death.  
 Do not use if you have bloody or black stool.  
 Ask a doctor before use if you have ■ a fever ■ mucus in the stool ■ a history of liver disease  
 ■ a history of abnormal heart rhythm  
 Ask a doctor or pharmacist before use if you are taking a prescription drug. Loperamide may interact with certain prescription drugs.  
 When using this product, tiredness, drowsiness, or dizziness may occur. Be careful when driving or operating machinery.  
 Stop use and ask a doctor if ■ symptoms get worse ■ diarrhea lasts for more than 2 days  
 ■ you get abdominal swelling or bulging. These may be signs of a serious condition.  
 If pregnant or breast-feeding, ask a health professional before use.  
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.  
**Directions** ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea  
 ■ find right dose on chart. If possible, use weight to dose; otherwise, use age.  

adults and children 12 years and over	2 caplets after the first loose stool; 1 caplet after each subsequent loose stool; but no more than 4 caplets in 24 hours
children 9-11 years (60-95 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 3 caplets in 24 hours
children 6-8 years (48-59 lbs)	1 caplet after the first loose stool; 1/2 caplet after each subsequent loose stool; but no more than 2 caplets in 24 hours
children 2-5 years (34-47 lbs)	ask a doctor
children under 2 years (up to 33 lbs)	do not use

**Premier Value 44-375**

**ANTI-DIARRHEAL**

loperamide hcl tablet

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:68016-301
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>LOPERAMIDE HYDROCHLORIDE</b> (UNII: 77TI35393C) (LOPERAMIDE - UNII:6X9OC3H4II)	LOPERAMIDE HYDROCHLORIDE	2 mg

**Inactive Ingredients**

Ingredient Name	Strength
-----------------	----------

<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>DIBASIC CALCIUM PHOSPHATE DIHYDRATE</b> (UNII: O7TSZ97GEP)	
<b>FD&amp;C BLUE NO. 1 ALUMINUM LAKE</b> (UNII: J9EQA3S2JM)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	

### Product Characteristics

<b>Color</b>	green	<b>Score</b>	2 pieces
<b>Shape</b>	OVAL	<b>Size</b>	10mm
<b>Flavor</b>		<b>Imprint Code</b>	44;375
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:68016-301-18	3 in 1 CARTON	12/06/2019	11/30/2026
1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:68016-301-24	4 in 1 CARTON	12/06/2019	11/30/2026
2		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:68016-301-06	1 in 1 CARTON	12/06/2019	11/30/2022
3		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076497	12/06/2019	11/30/2026

### Labeler - Chain Drug Consortium (101668460)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(68016-301) , pack(68016-301)

### Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(68016-301)

