

**ARIELLA HEMORRHOID GEL- lidocaine, phenylephrine hcl ointment**  
**YITONGBADA (SHENZHEN) INTERNATIONAL TRADE CO., LTD**

-----

**Ariella Hemorrhoid Gel**

Drug Facts

## Drug Facts

### Active ingredients

	Purpose
Lidocaine 6% .....	Protectant
Phenylephrine HCL 0.25%.....	Vasoconstrictor

### Uses

-helps relieve itching and discomfort associated with hemorrhoids - temporarily shrinks swelling and relieves burning and pain -soothes and protects irritated tissue

### Warnings

**For external and/or intrarectal use only**

#### Ask a doctor before use if you have

-heart disease -high blood pressure - thyroid disease - diabetes -difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are** presently taking a prescription drug for high blood pressure or depression.

**When using this product** do not exceed the recommended daily dosage unless directed by a doctor

**Stop and ask a doctor if** - bleeding occurs -condition worsens or does not improve within 7 days -introduction of the applicator into the rectum causes additional pain -an allergic reaction occurs - symptom does not improve or if redness, irritation, swelling, pain or other symptoms develop or increase

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of accidental ingestion, seek medical advice immediately or contact a Poison Control Center right away. (1-800-222-1222)

### Directions

-when practical, cleanse the affected area by patting or blotting with an appropriate cleansing wipe. Gently dry by patting or blotting with a tissue or a soft cloth before application.

-apply half the gel in the applicator to the external affected area (*only applies if you have external symptoms*)

-find a comfortable position and gently insert about ½ inch into the rectum.

-push the end of the applicator to apply the remaining gel inside the rectum (*Both steps are necessary for both internal and external symptoms as external hemorrhoid often comes with internal hemorrhoid*)

-dispose the applicator after use.

-avoid movement for 1-3 hours

-Children under 12 years of age: ask a doctor

**Other information** store at 15-25°C (59-77°F)

**Inactive ingredients** Borneol, Carbomer Homopolymer, Chlorhexidine Acetate, Glycerin, Thymol

**Questions or comments?** Contact us at [www.tryariella.com](http://www.tryariella.com) or email [cs@tryariella.com](mailto:cs@tryariella.com)

Active ingredients Purpose

Lidocaine 6% .....Protectant Phenylephrine HCL 0.25%  
.....Vasoconstrictor

Uses

-helps relieve itching and discomfort associated with hemorrhoids - temporarily shrinks swelling and relieves burning and pain -soothes and protects irritated tissue

For external and/or intrarectal use only

Ask a doctor before use if you have

-heart disease -high blood pressure - thyroid disease - diabetes -difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are presently taking a prescription drug for high blood pressure or depression.

When using this product do not exceed the recommended daily dosage unless directed by a doctor

Stop and ask a doctor if - bleeding occurs -condition worsens or does not improve within 7 days -introduction of the applicator into the rectum causes additional pain -an allergic reaction occurs - symptom does not improve or if redness, irritation, swelling, pain or other symptoms develop or increase

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental ingestion, seek medical advice immediately or contact a Poison Control Center right away. (1-800-222-1222)

## Warnings

## Directions

-when practical, cleanse the affected area by patting or blotting with an appropriate cleansing wipe. Gently dry by patting or blotting with a tissue or a soft cloth before application.

-apply half the gel in the applicator to the external affected area (only applies if you have external symptoms)

-find a comfortable position and gently insert about ½ inch into the rectum.

-push the end of the applicator to apply the remaining gel inside the rectum (Both steps are necessary for both internal and external symptoms as external hemorrhoid often comes with internal hemorrhoid)

-dispose the applicator after use.

-avoid movement for 1-3 hours

-Children under 12 years of age: ask a doctor

Other information store at 15-25°C (59-77°F)

## Inactive ingredients

Borneol, Carbomer Homopolymer, Chlorhexidine Acetate, Glycerin, Thymol

## **ARIELLA HEMORRHOID GEL**

lidocaine, phenylephrine hcl ointment

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83364-007(NDC:84010-001)
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	0.25 g in 100 g
<b>LIDOCAINE</b> (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	4 g in 100 g

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>BORNEOL</b> (UNII: M89NIB437X)	
<b>CARBOMER HOMOPOLYMER, UNSPECIFIED TYPE</b> (UNII: 0A5MM307FC)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>CHLORHEXIDINE ACETATE</b> (UNII: 5908Z UF22Y)	
<b>THYMOL</b> (UNII: 3J50XA376E)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:83364-007-01	1 in 1 BOX	06/25/2024	
1		6 in 1 APPLICATOR		
1		3 g in 1 APPLICATOR; Type 0: Not a Combination Product		

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M005	06/25/2024	

**Labeler** - YITONGBADA (SHENZHEN) INTERNATIONAL TRADE CO., LTD (725220463)

Revised: 6/2024

YITONGBADA (SHENZHEN) INTERNATIONAL TRADE CO., LTD