

**PULMOTIL 90- tilmicosin phosphate granule**  
**Elanco US Inc.**

-----  
**Elanco™**

AF0472

***Pulmotil™ 90***

**Net Weight:**

**10 kg (22.0 lb)**

***Tilmicosin***

**Type A Medicated Article**

**CAUTION:** Federal law restricts medicated feed containing this veterinary feed directive (VFD) drug to use by or on the order of a licensed veterinarian.

**For Use in Swine and Cattle Feeds Only.**

**Do not feed undiluted.**

**Active Drug Ingredient:** Tilmicosin (as tilmicosin phosphate) 90.7 g per lb (200 g per kg)

**Inert Ingredients:** Ground corncobs.

**Description:**

Pulmotil is a formulation of the antibiotic tilmicosin. Tilmicosin is produced semi-synthetically and is in the macrolide class of antibiotics. Each kilogram of Type A Medicated Article contains 200 grams (0.44 lbs) of tilmicosin adsorbed onto ground corncobs.

**Indications:**

**Swine:** For the control of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae* and *Pasteurella multocida*.

**Cattle:** For the control of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida* and *Histophilus somni* in groups of beef and non-lactating dairy cattle, where active BRD has been diagnosed in at least 10% of the animals in the group.

**Feeding Directions:**

**Swine:** Tilmicosin is to be fed continuously at 181 grams to 363 grams per ton (200 ppm to 400 ppm) of Type C medicated feed as the sole ration for a 21-day period, beginning approximately 7 days before an anticipated disease outbreak.

**Cattle:** Tilmicosin is to be fed continuously for a single, 14 day period at 568 grams to 757 grams (626 ppm to 834 ppm) per ton on a 100% dry matter basis of Type C medicated feed as the sole ration to provide 12.5 mg/kg of body weight/day.

**IMPORTANT: Must be thoroughly mixed in swine or cattle feeds before use.**

**Mixing Directions:**

**For Incorporation into Swine Feeds:** Thoroughly mix Pulmotil Type A medicated article with feed to provide a Type B medicated feed containing up to 36,300 grams tilmicosin per ton or to provide a complete Type C medicated feed containing 181 to 363 g tilmicosin per ton. Do not use in any feeds containing bentonite. Bentonite in feeds may affect the efficacy of tilmicosin.

Starting concentration of Pulmotil 90 Type A Medicated Article <sup>a</sup>	Amount of Type A Medicated Article to add per ton	Resulting concentration in Type B Medicated Feed	
		grams per ton	grams per pound
90.7	400	36,300	18.1
	300	27,200	13.6
	200	18,100	9.1

  

Starting concentration of Pulmotil 90 Type A Medicated Article <sup>a</sup>	Amount of Type A Medicated Article to add per ton	Resulting concentration in Type C Medicated Feed
		grams per ton
90.7	4	363
	3	272
	2	181

<sup>a</sup>Pulmotil 90 contains 90.7 g tilmicosin phosphate per pound

**For Incorporation into Cattle Feeds:** Thoroughly mix Pulmotil Type A medicated article with feed to provide a Type B medicated feed containing up to 36,300 grams tilmicosin per ton on a 100% dry matter basis or to provide a complete Type C medicated feed containing 568 to 757 g tilmicosin per ton on a 100% dry matter basis. Complete Type C medicated feeds should not be pelleted. Do not use in any feeds containing bentonite, cottonseed meal, or cottonseed hulls. Bentonite, cottonseed meal, or cottonseed hulls in feeds may affect the efficacy of tilmicosin.

Starting concentration of Pulmotil 90 Type A Medicated Article <sup>a</sup>	Amount of Type A Medicated Article to add per ton	Resulting concentration in Type B Medicated Feed <sup>b</sup>	
		grams per ton	grams per pound
90.7	400	36,300	18.1
	200	18,100	9.1

	100	9,070	4.5
Starting concentration of Pulmotil 90 Type A Medicated Article <sup>a</sup>	Amount of Type A Medicated Article to add per ton	Resulting concentration in Type C Medicated Feed <sup>b</sup>	
grams per pound	pounds	grams per ton	
90.7	8.35	757	
	6.26	568	

<sup>a</sup>Pulmotil 90 contains 90.7 g tilmicosin phosphate per pound

<sup>b</sup>100% dry matter basis

### CAUTION:

Do not allow horses or other equines access to feeds containing tilmicosin. The safety of tilmicosin has not been established in cattle or male swine intended for breeding purposes.

To assure both food safety and responsible use in cattle, the treatment of cattle with this medicated feed is required to be initiated within the first 45 days of the production period. The treatment should not occur concurrent with or following administration of an injectable macrolide, or within 3 days following administration of a non-macrolide injectable BRD therapy.

**Swine:** Feed containing tilmicosin shall not be fed to pigs for more than 21 days during each phase of production without ceasing administration for reevaluation of antimicrobial use by a licensed veterinarian before re-initiating a further course of therapy with an appropriate antimicrobial. Veterinary Feed Directive (VFD) expiration date for swine must not exceed 90 days from the time of issuance. VFDs for tilmicosin phosphate shall not be refilled.

**Cattle:** Use only in cattle fed in confinement for slaughter. Tilmicosin medicated feed treatment has not been evaluated in cattle with severe clinical disease. Cattle with severe clinical illness should be evaluated for individual treatment with an alternative non-macrolide therapy. The expiration date for a tilmicosin Veterinary Feed Directive (VFD) for cattle must not exceed 45 days from the time of issuance. VFDs for tilmicosin phosphate shall not be refilled.

### WARNINGS:

**RESIDUE WARNING: Swine:** Swine intended for human consumption must not be slaughtered within 7 days of the last treatment of this drug product.

**RESIDUE WARNING: Cattle:** Cattle intended for human consumption must not be slaughtered within 28 days of the last treatment with this drug product.

This drug product is not approved for use in female dairy cattle 20 months of age or older. Use in these cattle may cause drug residues in milk. This drug product is not approved for use in calves intended to be processed for veal. A withdrawal period has not been established in pre-ruminating calves.

**User Safety Warnings:** Avoid inhalation, oral exposure and direct contact with skin or

eyes. Operators mixing and handling Pulmotil 90 should use protective clothing, impervious gloves, goggles and a NIOSH-approved dust mask. Wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water. If irritation persists, seek medical attention. Not for human consumption. Keep out of reach of children. The Safety Data Sheet contains more detailed occupational safety information. To report adverse effects in users, to obtain more information, or to obtain a Safety Data Sheet, call 1-800-428-4441.

**Clinical Pharmacology:** Oral dosing of tilmicosin phosphate to swine at 181 to 363 g/ton of feed results in serum tilmicosin levels, which do not correlate with efficacy. Lung concentrations of tilmicosin are significantly higher than serum. Following 7 consecutive days of administering tilmicosin-medicated feeds to swine, the concentration of tilmicosin in respiratory tissues, phagocytic cells, and nasal secretions was significantly higher than that of plasma or serum. Lung levels are achieved within 2 days after beginning feeding and plateau by 4 days. Using *in-vitro* incubation techniques, the ratio of intracellular to extracellular concentrations of tilmicosin for neutrophils, monocyte-macrophages and alveolar macrophages were 69, 19 and 17, respectively, after four hours of incubation. Although lower levels of accumulation were observed *in-vivo*, swine alveolar macrophages have been shown *in-vitro* and *in-vivo* to concentrate large amounts of tilmicosin; these cells may be important for *in-vivo* distribution of the drug and may serve as an important reservoir for tilmicosin in lung tissue.

Oral dosing of tilmicosin phosphate to cattle to target a dose of 12.5 mg/kg body weight resulted in serum tilmicosin concentrations above the analytical limit of quantification (0.5 ng/mL) within 12 hours following treatment administration. The relationship of serum tilmicosin concentration to lung tilmicosin concentration has not been determined following oral administration of tilmicosin.

**Toxicology:** The cardiovascular system is the target of toxicity in laboratory and domestic animals given tilmicosin by oral or parenteral routes. Primary cardiac effects are increased heart rate (tachycardia) and decreased contractility (negative inotropy). Given orally, the median lethal dose is 800 mg/kg in fasted rats and 2250 mg/kg in non-fasted rats. No compound-related lesions were found at necropsy. Results of genetic toxicology studies were all negative. Results of teratology and reproduction studies in rats were all negative. The no effect level in dogs after daily oral doses for up to one year is 4 mg/kg of body weight. Tilmicosin was included in the diet of 18 adult horses for a period of 14 days at dose levels of 400, 1200 and 2000 ppm. Some horses at both the low and high dose levels demonstrated gastrointestinal disturbance with more severe colic evident at the higher levels. One horse died after consuming the 2000 ppm diet. A study was conducted in cattle administered oral tilmicosin at 12.5, 25.0 or 37.5 mg/kg for 42 days or administered 12.5 mg/kg of oral tilmicosin for 14 days followed by 20 mg/kg injection of tilmicosin or saline (volume equivalent). Cardiac lesions observed (one animal in the 12.5 mg/kg for 42 days treatment group; one animal in the 12.5 mg/kg for 14 days followed by tilmicosin injection treatment group) were not considered clinically significant as no other abnormalities were seen and the affected animals were clinically normal.

To report adverse effects, access medical information or obtain additional product information, call 1-800-428-4441.

**Storage Information:** Store at less than or equal to 25°C (77°F). Excursions to 40°C (104°F) are acceptable. Avoid excessive moisture.

Not to be used after the date printed on the bag.

Restricted Drug (California) - Use Only as Directed

Approved by FDA under NADA # 141-064

Manufactured For: Elanco US Inc, Greenfield, IN 46140, USA

Pulmotil, Elanco and the diagonal bar logo are trademarks of Elanco or its affiliates.

**TAKE TIME TO**



**OBSERVE LABEL DIRECTIONS**

**BG280428X**

**Principal Display Panel - 22 lb Bag Label**

***Elanco***<sup>™</sup> AF0472

***Pulmotil***<sup>™</sup> 90

**Net Weight:**  
**10 kg (22.0 lb)**

***Tilmicosin***



Elanco™

AF0472

# Pulmotil™ 90

Net Weight:  
10 kg (22.0 lb)

## Tilmosicin

### Type A Medicated Article

Caution: Federal law prohibits dispensing this veterinary feed directive (VFD) drug to use beyond the order of a licensed veterinarian.

#### For Use in Swine and Cattle Feeds Only

##### Directions

Swine: 10 mg/kg (0.225 mg/lb) of live weight (0.225 g per 100 g of feed)

##### Directions

For the control of acute respiratory disease (ARD) associated with *Actinobaculum suis* and *Streptococcus suis* in swine, the recommended dose is 10 mg/kg of live weight (0.225 mg/lb) of live weight (0.225 g per 100 g of feed) in the feed.

**Caution:** For the control of acute respiratory disease (ARD) associated with *Actinobaculum suis* and *Streptococcus suis* in swine, the recommended dose is 10 mg/kg of live weight (0.225 mg/lb) of live weight (0.225 g per 100 g of feed) in the feed.

**Caution:** For the control of acute respiratory disease (ARD) associated with *Actinobaculum suis* and *Streptococcus suis* in swine, the recommended dose is 10 mg/kg of live weight (0.225 mg/lb) of live weight (0.225 g per 100 g of feed) in the feed.

##### Directions

For the control of acute respiratory disease (ARD) associated with *Actinobaculum suis* and *Streptococcus suis* in swine, the recommended dose is 10 mg/kg of live weight (0.225 mg/lb) of live weight (0.225 g per 100 g of feed) in the feed.

Lot No.	Net Weight (kg)	Net Weight (lb)
001	10.0	22.0
002	10.0	22.0
003	10.0	22.0
004	10.0	22.0
005	10.0	22.0
006	10.0	22.0
007	10.0	22.0
008	10.0	22.0
009	10.0	22.0
010	10.0	22.0

Lot No.	Net Weight (kg)	Net Weight (lb)
011	10.0	22.0
012	10.0	22.0
013	10.0	22.0
014	10.0	22.0
015	10.0	22.0
016	10.0	22.0
017	10.0	22.0
018	10.0	22.0
019	10.0	22.0
020	10.0	22.0

Lot No.	Net Weight (kg)	Net Weight (lb)
021	10.0	22.0
022	10.0	22.0
023	10.0	22.0
024	10.0	22.0
025	10.0	22.0
026	10.0	22.0
027	10.0	22.0
028	10.0	22.0
029	10.0	22.0
030	10.0	22.0

Lot No.	Net Weight (kg)	Net Weight (lb)
031	10.0	22.0
032	10.0	22.0
033	10.0	22.0
034	10.0	22.0
035	10.0	22.0
036	10.0	22.0
037	10.0	22.0
038	10.0	22.0
039	10.0	22.0
040	10.0	22.0

Lot No.	Net Weight (kg)	Net Weight (lb)
041	10.0	22.0
042	10.0	22.0
043	10.0	22.0
044	10.0	22.0
045	10.0	22.0
046	10.0	22.0
047	10.0	22.0
048	10.0	22.0
049	10.0	22.0
050	10.0	22.0

Caution: Federal law prohibits dispensing this veterinary feed directive (VFD) drug to use beyond the order of a licensed veterinarian.



LIT No:  
D.O.M.  
EXP:

# PULMOTIL 90

tilmosicin phosphate granule

Product Information			
Product Type	VFD TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:58198-0472
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
	Ingredient Name	Basis of Strength	Strength
	tilmosicin phosphate (UNII: SMH7U1S683) (tilmosicin - UNII:XL4103X2E3)	tilmosicin phosphate	200 g in 1 kg

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58198-0472-9	10 kg in 1 BAG		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA141064	10/18/2012	

**Labeler** - Elanco US Inc. (966985624)

Revised: 12/2024

Elanco US Inc.