

**MEIJER DAYTIME COLD AND FLU SOFTGEL- acetaminophen,
dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid
filled
Meijer, Inc.**

meijer Daytime Cold & Flu SOFTGEL

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion • cough due to minor throat & bronchial irritation • sore throat
- headache • minor aches & pains • fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs

for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease • heart disease • diabetes
- high blood pressure • thyroid disease
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

When using this product, do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 8 softgels per 24 hrs

adults & children 12 yrs & over	2 softgels with water every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- store at room temperature between 20°-25°C (68°-77°F)
- protect from light, heat, and moisture

Inactive ingredients

edible printing ink, FD&C red no. 40, FD&C yellow no. 6, gelatin (bovine), glycerin, polyethylene glycol 400, povidone K-17, propylene glycol, purified water, sorbitol sorbitan solution

Questions or comments?

1-888-577-8033 Monday - Friday 8am - 4pm EST

COMPARE TO VICKS® DAYQUIL™ COLD & FLU LIQUICAPS™ ACTIVE INGREDIENTS*

ANTIHISTAMINE FREE

*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the trademarks Vicks®, DayQuil™ and LiquiCaps™.

**DIST. BY MEIJER DISTRIBUTION, INC.
GRAND RAPIDS, MI 49544**


WWW.MEIJER.COM

PRODUCT OF UNITED ARAB EMIRATES

READ AND KEEP OUTER CARTON FOR COMPLETE PRODUCT WARNINGS AND INFORMATION

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

Packaging



*This product is not manufactured or distributed by The Procter & Gamble Company, owner of the trademarks Vicks®, DayQuil™ and LiquiCaps™.
 OUR QUALITY GUARANTEE
 DIST. BY MEIJER DISTRIBUTION, INC. GRAND RAPIDS, MI 49544
 WWW.MEIJER.COM
 PRODUCT OF UNITED ARAB EMIRATES
 1-888-577-8033 Monday - Friday 8am - 4pm EST
 Net wt 20.25g (68.77g)
 edible printing ink, FD&C red no. 40, FD&C yellow, polyethylene glycol 400, povidone K-17, propylene glycol, purified water, sorbitol sorbitan solution

NDC 79481-0807-4

meijer.

COMPARE TO VICKS® DAYQUIL™ COLD & FLU LIQUICAPS™ ACTIVE INGREDIENTS*

Daytime Cold & Flu

PAIN RELIEVER/FEVER REDUCER,
COUGH SUPPRESSANT,
NASAL DECONGESTANT

Acetaminophen | 325mg
Dextromethorphan HBr | 10mg
Phenylephrine HCl | 5mg

24 SOFTGELS**
(**Liquid-filled Capsules)

ime Cold & Flu

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg.....Pain reliever/fever reducer
 Dextromethorphan HBr 10 mg.....Cough suppressant
 Phenylephrine HCl 5 mg.....Nasal decongestant

Purpose

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

Uses

temporarily relieves common cold/flu symptoms:

- more than 8 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 softgels in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor

Drug Facts (continued)

- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

When using this product, do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy, or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- do not exceed 8 softgels per 24 hrs



DRUG FACTS LABEL

Drug Facts Active ingredients (in each softgel) Acetaminophen 325 mg.....Pain reliever/fever reducer Dextromethorphan HBr 10 mg.....Cough suppressant Phenylephrine HCl 5 mg.....Nasal decongestant		Drug Facts (continued) ■ trouble urinating due to enlarged prostate gland ■ cough that occurs with too much phlegm (mucus) ■ persistent or chronic cough as occurs with smoking, asthma, or emphysema Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin		Drug Facts (continued) Other information ■ store at room temperature between 20°-25°C (68°-77°F) ■ protect from light, heat, and moisture Inactive ingredients edible printing ink, FD&C red no. 40, FD&C yellow no. 6, gelatin (bovine), glycerin, polyethylene glycol 400, povidone K-17, propylene glycol, purified water, sorbitol solution							
Purpose temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ cough due to minor throat & bronchial irritation ■ sore throat ■ headache ■ minor aches & pains ■ fever		When using this product, do not use more than directed Stop use and ask a doctor if ■ you get nervous, dizzy, or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.		Questions or comments? 1-888-577-8033 Monday - Friday 8am - 4pm EST							
Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 8 softgels in 24 hours, which is the maximum daily amount for this product ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away. Sore throat warning: If sore throat is severe, lasts for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.		If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.									
Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.		Directions ■ take only as directed ■ do not exceed 8 softgels per 24 hrs									
Ask a doctor before use if you have ■ liver disease ■ heart disease ■ diabetes ■ high blood pressure ■ thyroid disease		<table border="1"> <tr> <td>adults & children 12 yrs & over</td> <td>2 softgels with water every 4 hrs</td> </tr> <tr> <td>children 4 to under 12 yrs</td> <td>ask a doctor</td> </tr> <tr> <td>children under 4 yrs</td> <td>do not use</td> </tr> </table>		adults & children 12 yrs & over	2 softgels with water every 4 hrs	children 4 to under 12 yrs	ask a doctor	children under 4 yrs	do not use		
adults & children 12 yrs & over	2 softgels with water every 4 hrs										
children 4 to under 12 yrs	ask a doctor										
children under 4 yrs	do not use										

MEIJER DAYTIME COLD AND FLU SOFTGEL

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79481-0807
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN TYPE B BOVINE (230 BLOOM) (UNII: WL1404U79)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
POVIDONE K17 (UNII: C67P1734QJ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
SORBITAN (UNII: 6O92ICV9RU)	

Product Characteristics

Color	red	Score	no score
Shape	OVAL	Size	21mm
Flavor		Imprint Code	807
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79481-0807-4	2 in 1 CARTON	06/27/2024	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/27/2024	

Labeler - Meijer, Inc. (006959555)