

VIVIER SHEER SPF 30 MINERAL- titanium dioxide, zinc oxide lotion
Vivier Pharma, Inc.

Vivier Sheer SPF 30 Mineral

Drug Facts

Active ingredients (w/w)

Octinoxate 7.5% Avobenzone 2.5%

Ensulizole 2%

Purpose

Sun Protectant

Uses

- Helps prevent sunburn • Sun Protection Factor 30 • If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Do not use

on broken skin.

When using this product

avoid contact with eyes. If contact occurs, rinse thoroughly with water.

Stop use and ask a doctor if

rash occurs.

Keep out of reach of children.

If swallowed, call a poison control centre or get medical help right away.

Directions

Adults and children over 6 months of age: • Apply liberally and evenly 15 minutes before sun exposure • Reapply at least every 2 hours • Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15

or higher and other sun protection measures including: • Limit time in the sun, especially from 10 a.m.– 2 p.m.; and • wear long-sleeved shirts, pants, hats, and sunglasses.

Other information

Store at room temperature (15-30°C / 59-86°F).

Inactive ingredients

Butylated Hydroxytoluene, Butylparaben, Carbomer, Cetyl Alcohol, Dimethicone, Disodium EDTA, D-Panthenol, Ethylparaben, Hydrogenated Palm Glycerides, Methylparaben, Octyldodecanol, Octyldodecyl Neopentanoate, Phenoxyethanol, Potassium Cetyl Phosphate, Propylene Glycol, Propylparaben, Sodium Hydroxide, Stearic Acid, Tocopheryl Acetate, Triethanolamine, Water.

Questions?

1 877 484 8437

Package Labeling:

FRONT PANEL

BACK PANEL

Lot XXXXX UNPRINTABLE AREA (12mm)	EXP YYYY-MM
<p>NPN 80002049 NDC 67226-xxxx-x</p> <p> VIVIER®</p> <p>SHEER SPF FPS 30 MINERAL</p> <p>BROAD SPECTRUM UVA/UVB SUNSCREEN WITH ANTIOXIDANTS</p> <p>—</p> <p>MINERAL ÉCRAN SOLAIRE À LARGE SPECTRE UVA / UVB AVEC ANTIOXYDANTS</p> <p>60 ml 2 fl oz</p>	<p>Directions Adults and children over 6 months of age: Apply daily to face, neck and chest. Apply liberally and evenly before 15 minutes before sun exposure. Reapply at least every 2 hours. Use a water resistant sunscreen if swimming or sweating. Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including: limit time in the sun, especially from 10 a.m.-2 p.m.; and wear long-sleeved shirts, pants, hats and sunglasses. Store at room temperature (15-30°C / 59-86°F).</p> <p>Warnings For external use only. Do not use on broken skin. When using this product avoid contact with eyes. If contact occurs, rinse thoroughly with water. Stop use and ask a doctor if rash occurs. Keep out of reach of children. If swallowed, call a poison control centre or get medical help right away.</p> <p>This unit is packaged in an outer container. Please see outer container for complete product facts information.</p> <p>Medicinal ingredients / Ingrédients médicinaux Titanium Dioxide 10% / Dioxyde de Titane 10 % Zinc Oxide 8% / Oxyde de Zinc 8 %</p> <p>Mode d'emploi Adultes et enfants de plus de 6 mois : Appliquer quotidiennement au visage, cou et décolleté. Appliquer abondamment et uniformément 15 minutes avant l'exposition au soleil. Appliquer de nouveau au moins toutes les 2 heures. Utiliser un écran solaire résistant à l'eau lors d'une activité aquatique ou s'il y a transpiration. Mesures de protection contre le soleil: Passer du temps au soleil augmente le risque de cancer de la peau et du vieillissement prématûre de la peau. Afin de diminuer ce risque, il convient d'utiliser régulièrement un écran solaire avec un F.P.S. à large spectre d'au moins 15 et d'appliquer d'autres mesures de protection contre le soleil, telles que: limiter le temps passé au soleil, notamment entre 10h-14h; et porter des chemises à manches longues, des pantalons, un chapeau et des lunettes de soleil. Conserver à la température ambiante (15-30 °C / 59-86 °F).</p> <p>Mises en garde Pour usage externe seulement. Ne pas utiliser sur une peau éraflée. Lorsque vous utilisez ce produit évitez tout contact avec les yeux. Le cas échéant, bien rincer avec de l'eau. Cessez d'utiliser et consultez un médecin si une éruption cutanée se développe. Garder hors de la portée des enfants. En cas d'ingestion, appeler immédiatement un centre antipoison ou obtenir une assistance médicale. Cette unité est emballée dans un contenant extérieur. Veuillez vous y référer pour consulter toute l'information sur le médicament.</p> <p>Questions? 1 877 484 8437</p> <p> VIVIER® Vivier Canada Inc. Mfd. for Vivier Pharma Vaudreuil-Dorion QC Canada J7V 5V5 Champlain NY USA 12919 vivierskin.com</p> <p>Made in Canada 01-8065-01D07</p>

**Product Facts (continued) /
Informations sur le produit (suite)**

**Non-medicinal ingredients /
Ingrédients non-médicinaux**

Alumina, Ammonium Acryloyldimethyltaurate/VP Copolymer, Bis PEG/PPG-16/16 PEG/PPG-16/16 Dimethicone, Butylparaben, C12-15 Alkyl Benzoate, Camellia Sinensis (Green Tea) Extract, Caprylic/Capric Triglyceride, Ceteareth 20, Cetearyl Alcohol, Cetyl Dimethicone, Citrus Aurantium Dulcis (Orange) Peel Extract, Citrus Limon (Lemon) Peel Extract, Cyclohexasiloxane, Cyclopentasiloxane, Dimethicone, Dipropylene Glycol Dibenzoate, Ethylparaben, Glycerin, Glyceryl Stearate, Isopropyl Titanium Triisostearate, Methylparaben, 2-Methylpropan-2-ol, Phenoxethanol, Polyhydroxystearic Acid, Polysorbate 20, PPG-15 Stearyl Ether Benzozate, Propylene Glycol, Propylparaben, Pyrus Malus (Apple) Fruit Extract, Saccharum Officinarum (Sugar Cane) Extract, Sodium Ascorbyl Phosphate, Tocopherol Acetate, Triethoxycaprylylsilane, VP/Eicosene Copolymer, Water/Eau, Xanthan Gum.

Questions? 1 877 484 8437

**THE BEAUTY OF RESULTS®
LA BEAUTÉ DES RÉSULTATS®**



Vivier Canada Inc. | Mfd. for Vivier Pharma
Vaudreuil-Dorion QC Canada J7V 5V5
Champlain NY USA 12919
vivierskin.com

Made in Canada

© Vivier Canada Inc. - used under license



Product Facts

Medicinal ingredients	Purpose
Titanium Dioxide 10% Zinc Oxide 8%	Sunscreen

Uses • Helps prevent sunburn and provides a sun protection factor of 30. • If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun.

Warnings

For external use only.

Do not use on broken skin.

When using this product avoid contact with eyes. If contact occurs, rinse thoroughly with water.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children.

If swallowed, call a poison control centre or get medical help right away.

Directions Adults and children over 6 months of age: • Apply daily to face, neck and chest. • Apply liberally and evenly 15 minutes before sun exposure.

• Reapply at least every 2 hours. • Use a water resistant sunscreen if swimming or sweating. • Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:

- limit time in the sun, especially from 10 a.m. - 2 p.m.; and
- wear long-sleeved shirts, pants, hats, and sunglasses. • Ideal for daily use.

Other information

Store at room temperature (15-30°C / 59-86°F). ►

NPN 80002049 | NDC 67226-xxxx-x



**SHEER
SPF | FPS**

30

MINERAL

BROAD SPECTRUM UVA/UVB
SUNSCREEN WITH ANTIOXIDANTS

MINÉRAL

ÉCRAN SOLAIRE À LARGE
SPECTRE UVA/UVB AVEC
ANTIOXYDANTS

60 ml | 2 fl oz

Informations sur le produit

Ingrédients médicinaux	Utilité
Dioxyde de Titane 10 % Oxyde de Zinc 8 %	Écran solaire

Usages • Aide à prévenir les coups de soleil et offre un facteur de protection solaire de 30. • Si le produit est utilisé selon les instructions et avec d'autres mesures de protection contre le soleil (voir le Mode d'emploi), le risque de cancer de la peau et de vieillissement prématuré de la peau causé par le soleil est réduit.

Mises en garde

Pour usage externe seulement.

Ne pas utiliser sur une peau éraflée.

Lorsque vous utilisez ce produit évitez tout contact avec les yeux. Le cas échéant, bien rincer avec de l'eau.

Cessez d'utiliser et consultez un médecin si une éruption cutanée se développe.

Garder hors de la portée des enfants.

En cas d'ingestion, appeler immédiatement un centre antipoison ou obtenir une assistance médicale.

Mode d'emploi Adultes et enfants de plus de 6 mois: • Appliquer quotidiennement au visage, cou et décolleté. • Appliquer abondamment et uniformément 15 minutes avant l'exposition au soleil. • Appliquer de nouveau au moins toutes les 2 heures. • Utiliser un écran solaire résistant à l'eau lors d'une activité aquatique ou s'il y a transpiration. • Mesures de protection contre le soleil: Passer du temps au soleil augmente le risque de cancer de la peau et du vieillissement prématuré de la peau. Afin de diminuer ce risque, il convient d'utiliser régulièrement un écran solaire avec un F.P.S. à large spectre d'au moins 15 et d'appliquer d'autres mesures de protection contre le soleil, telles que: • limiter le temps passé au soleil, notamment entre 10h - 14h; et • porter des chemises à manches longues, des pantalons, un chapeau et des lunettes de soleil. • Idéal pour utilisation quotidienne.

Autre renseignement Conserver à la température ambiante (15-30 °C / 59-86 °F). ►

VIVIER SHEER SPF 30 MINERAL

titanium dioxide, zinc oxide lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67226-2850
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ZINC OXIDE (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	80 mg in 1 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	100 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
DIPROPYLENE GLYCOL DIBENZOATE (UNII: 60A5ZDY41O)	
ETHYLPARABEN (UNII: 14255EXE39)	
GLYCERIN (UNII: PDC6A3C00X)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

PHENOXYETHANOL (UNII: HIE492ZZ3T)

ETHYL BENZOATE (UNII: J115BRJ15H)

POLYSORBATE 20 (UNII: 7T1F30V5YH)

PROPYLPARABEN (UNII: Z8IX2SC1OH)

APPLE (UNII: B423VGH5S9)

SUGARCANE (UNII: 81H2R5AOH3)

SODIUM ASCORBYL PHOSPHATE (UNII: 836SJG51DR)

TRIETHOXYCAPRYLYLSILANE (UNII: LDC331P08E)

XANTHAN GUM (UNII: TTV12P4NEE)

.ALPHA.-TOCOPHEROL ACETATE (UNII: 9E8X80D2L0)

WATER (UNII: 059QF0KO0R)

ALUMINUM OXIDE (UNII: LMI26O6933)

ISOPROPYL TITANIUM TRIISOSTEARATE (UNII: 949E3KB1I)

METHYLPARABEN (UNII: A2I8C7HI9T)

GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)

AMMONIUM ACRYLOYLDIMETHYLTAURATE/VP COPOLYMER (UNII: W59H9296ZG)

BIS-PEG/PPG-16/16 PEG/PPG-16/16 DIMETHICONE (UNII: 55A74AJ3KB)

BUTYLPARABEN (UNII: 3QPI1U3FV8)

ALKYL (C12-15) BENZOATE (UNII: A9EJ3J61HQ)

GREEN TEA LEAF (UNII: W2ZU1RY8B0)

MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)

POLYOXYL 20 CETOSTEARYL ETHER (UNII: YRC528SWUY)

CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)

ORANGE PEEL (UNII: TI9T76XD44)

LEMON PEEL (UNII: 72O054U628)

CYCLOMETHICONE 6 (UNII: XHK3U310BA)

CYCLOMETHICONE 5 (UNII: 0THT5PCI0R)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67226-2850-0	1 in 1 CARTON	01/05/2023	
1		60 mL in 1 TUBE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	01/05/2023	

Labeler - Vivier Pharma, Inc. (250996550)

Revised: 10/2023

Vivier Pharma, Inc.