

**MUCUS RELIEF CONGESTION AND COUGH MAXIMUM STRENGTH-  
dextromethorphan hbr, guaifenesin, phenylephrine hci liquid  
TARGET Corporation**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredients (in each 20 mL)**

Dextromethorphan HBr 20 mg

Guaifenesin 400 mg

Phenylephrine HCl 10 mg

**Purposes**

Cough suppressant

Expectorant

Nasal decongestant

**Uses**

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritant
  - the intensity of coughing
  - the impulse to cough to help you get to sleep
  - nasal congestion due to a cold

**Warnings**

**Do not use**

- if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

**Ask a doctor before use if you have**

- heart disease
- high blood pressure

- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, emphysema
- cough that occurs with too much phlegm (mucus)

### **When using this product**

**do not use more than directed.**

### **Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back, or occurs with fever, rash, or headache that lasts.

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- mL= milliliter
- shake well before using
- adult and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- Children under 12 years of age do not use

### **Other information**

- each 20 mL contains: **sodium 17 mg**
- store between 20-25°C (68-77°F). Do not refrigerate.

### **Inactive ingredients**

citric acid, disodium EDTA, FD&C blue #1 FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

## Questions or comments?

Call 1-800-910-6874

## Principal Display Panel

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough\*

maximum strength

fast mucus relief severe congestion and cough

dextromethorphan HBr 20 mg (cough suppressant)

guaifenesin 400 mg (expectorant)

phenylephrine HCl 10 mg (nasal decongestant)

controls cough

relieves nasal and chest congestion

thins and loosens mucus

AGES 12 + YEARS

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough.

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.**

Dist. by Target Corp.

Minneapolis, MN 55403

## Product Label

NDC 11673-337-06

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough\*

maximum strength  
**fast mucus relief**  
**severe congestion**  
**and cough**

dextromethorphan HBr 20 mg  
(cough suppressant)  
guaifenesin 400 mg (expectorant)  
phenylephrine HCl 10 mg  
(nasal decongestant)

controls cough  
relieves nasal and chest congestion  
thins and loosens mucus



AGES  
**12+**  
YEARS

6 FL OZ (177 mL)

PLD-B283D LB004222

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND OR UNDER CAP IS BROKEN OR MISSING.

\*This product is not manufactured or distributed by Reckitt Benckiser, distributor of Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough.



094 14 0290 R01  
C-000274-01-049  
Dist. by Target Corp.  
Minneapolis, MN 55403

**PARENTS:**  
Learn about teen medicine abuse  
[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

TM & ©2021 Target Brands, Inc. PLD-B283D LB004223

### Drug Facts

Active ingredients (in each 20 mL)	Purposes
Dextromethorphan HBr 20 mg	Cough suppressant
Guaifenesin 400 mg	Expectorant
Phenylephrine HCl 10 mg	Nasal decongestant

PEEL CORNER FOR MORE DRUG FACTS

### Drug Facts (continued)

#### Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves
  - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
  - the intensity of coughing
  - the impulse to cough to help you get to sleep
- nasal congestion due to a cold

#### Warnings

##### Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- for children under 12 years of age

### Drug Facts (continued)

#### Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product, do not use more than directed.

#### Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough comes back, or occurs with fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

#### Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### Drug Facts (continued)

#### Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter
- shake well before using
- adults and children 12 years of age and older: 20 mL in dosing cup provided every 4 hours
- children under 12 years of age: do not use

#### Other information

- each 20 mL contains: sodium 17 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

#### Inactive ingredients

citric acid, disodium EDTA, FD&C blue #1, FD&C red #40, flavor, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose, xanthan gum

### Drug Facts (continued)

#### Questions or comments?

Call 1-800-910-6874

PEEL CORNER FOR MORE DRUG FACTS

**TARGET Maximum Strength Mucus Relief Severe Congestion and Cough**

# MUCUS RELIEF CONGESTION AND COUGH MAXIMUM STRENGTH

dextromethorphan hbr, guaifenesin, phenylephrine hci liquid

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11673-337
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 20 mL
<b>GUAIFENESIN</b> (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	400 mg in 20 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 20 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYL GALLATE</b> (UNII: 8D45NN7V92)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11673-337-06	177 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/31/2016	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/31/2016	

