

**SPF 50 MINERAL SUNSCREEN LIP- zinc oxide , titanium dioxide cream  
Guangzhou Tata Biotechnology Co., Ltd.**

-----  
**SPF 50 MINERAL SUNSCREEN LIP**

Zinc Oxide 4.85%  
Titanium Dioxide 8%

Sunscreen

Helps prevent sunburn.

if used as directed with other sun protection measures (see Directions)decreases the risk of skin cancer and early skin agingcaused by the sun.

For external use only.

Do not use on damaged or broken skin

When using this product keep out of eyes.

Rinse with water to remove.

Stop use and ask a doctor if rash occurs.

Keep out of reach of children.If swallowed, getmedical help or contact Posion Center right away.

Directions

·Applyliberally15 minutes before sun exposure·Reapply:·after 80 minutes of swimming or sweating·after towel drying, swimming, or·at least every 2 hours ·children under 6 months of age: ask a doctor.

.Protect this product from excessive heat and direct sun.Sun alert: Limiting sunexposure, wearing protective clothing, and using sunscreens may reduce the risk of skin aging, skin cancer,and other harmful effects ofthe sun.

Water,Dimethicone, Isohexadecane, EthylhexylPalmitate, Butylene Glycol, Glycerin, Isopropyl PalmitateSorbitan Sesquioleate, Cetyl Peg/Ppg-10/ Dimethicone.Silica, Disteardimonium Hectorite, Sodium Chloride,Trimethylsiloxysilicate, Peg-30 Dipolyhydroxystearate,

Tribehenin, Magnesium Stearate, Phenoxyethanol,Triethoxycaprylylsilane Ethylhexylglycerin, PentaerythrytyTetra-Di-T-Buty Hydroxyhydrocinnamate

Please call:1-817-398-8817 or emus at info@redi-usa.com.

48mm

# REDI

## SPF 50 MINERAL SUNSCREEN

Broad Spectrum SPF 50 with Zinc Oxide & UVA/UVB Protection, Water Resistant (80 minutes).  
1 fl. oz (29.6 ml)

**Drug Facts**

Active Ingredients	Purpose
Zinc Oxide 4.85%	Sunscreen
Titanium Dioxide 8%	Sunscreen

**Uses** - Helps prevent sunburn. If used as directed with other sun protection measures (see Directions), decreases the risk of skin cancer and early skin aging caused by the sun.

**Warnings**  
**For external use only.**  
 Do not use on damaged or broken skin.  
 When using this product keep out of eyes. Rinse with water to remove.  
 Stop use and ask a doctor if rash occurs.  
 Keep out of reach of children. If swallowed, get medical help or contact Poison Center right away.

**Directions** - Apply liberally 15 minutes before sun exposure. Reapply after 80 minutes of swimming or sweating, after towel-drying, swimming or, at least every 2 hours, children under 6 months of age, ask a doctor.

**Other Information** - Protect this product from excessive heat and direct sun. **Sun alert:** Limiting sun exposure, wearing protective clothing, and using sunscreens may reduce the risk of skin aging, skin cancer, and other harmful effects of the sun.

**Inactive Ingredients**  
 Water, Dimethicone, Behenylalcohol, Ethylhexyl Palmitate, Butylene Glycol, Chlorophyllin, Lecithin, Palmitate, Sorbitan Sesquisebacate, Cetyl Peg-9 Glycerin Ether, Dimethicone, Silica, Disteardimonium Hectorite, Sodium Chloride, Trimethylsiloxysulfate, Peg-50 Dipolyhydroxystearate, Tribohenin, Magnesium Stearate, Phenoxethanol, Triethoxycellulose, Ethylhexylglycerin, Pentaerythrityl Tetra-Di-T-Butyl Hydroxyhydroquinamate

**Questions?** Please call 1-877-398-8877 or email us at [info@redi-usa.com](mailto:info@redi-usa.com).

Fragrance Free    Oxibenzonone Free    Hypoallergenic  
 Parabens Free    Octinoxate Free    Vegan  
 Dfep: Redi SunK LLC Austin, TX 78744 USA    redi.us    @redisunscreen  
 Made in China

89mm

SPF 50 MINERAL SUNSCREEN LIP			
zinc oxide , titanium dioxide cream			
Product Information			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:84066-104
<b>Route of Administration</b>	TOPICAL		
Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	4.85 mg in 100 mL	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	8 mg in 100 mL	
Inactive Ingredients			
Ingredient Name	Strength		
<b>WATER</b> (UNII: 059QF0KO0R)			
<b>PENTAERYTHRITOL TETRAKIS(3-(3,5-DI-TERT-BUTYL-4-HYDROXYPHENYL)PROPIONATE)</b> (UNII: 255PIF62MS)			
<b>ISOPROPYL PALMITATE</b> (UNII: 8CRQ2TH63M)			

<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)
<b>ETHYLHEXYL PALMITATE</b> (UNII: 2865993309)
<b>TRIMETHYLSILOXYSILICATE (M/Q 0.8-1.0)</b> (UNII: 25LXE464L2)
<b>DISTEARDIMONIUM HECTORITE</b> (UNII: X687XDK09L)
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)
<b>TRIETHOXYCAPRYLYLSILANE</b> (UNII: LDC331P08E)
<b>ISOHEXADECANE</b> (UNII: 918X1OUF1E)
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)
<b>SORBITAN SESQUIOLEATE</b> (UNII: 0W8RRI5W5A)
<b>CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2)</b> (UNII: V2W71V8T0X)
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)
<b>PEG-30 DIPOLYHYDROXYSTEARATE</b> (UNII: 9713Q0S7FO)
<b>TRIBEHENIN</b> (UNII: 8OC9U7TQZ0)
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)
<b>PHENOXYETHANOL</b> (UNII: HIE492ZZ3T)
<b>ETHYLHEXYLGLYCERIN</b> (UNII: 147D247K3P)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84066-104-01	29.6 mL in 1 BOTTLE; Type 0: Not a Combination Product	07/03/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	07/03/2024	

**Labeler** - Guangzhou Tata Biotechnology Co., Ltd. (699605102)

**Registrant** - Guangzhou Tata Biotechnology Co., Ltd. (699605102)

### Establishment

Name	Address	ID/FEI	Business Operations
Guangzhou Tata Biotechnology Co., Ltd.		699605102	manufacture(84066-104)

Revised: 7/2024

Guangzhou Tata Biotechnology Co., Ltd.