

**FEXOFENADINE HCL AND PSEUDOEPHEDRINE HCl- fexofenadine hcl and pseudoephedrine hcl tablet, extended release  
Wal-mart Stores Inc.**

-----  
**Fexofenadine HCl 60 mg and Pseudoephedrine HCl 120 mg ER Tablets USP**

**Active ingredient(s)**

Fexofenadine HCl USP, 60 mg

Pseudoephedrine HCl USP, 120 mg

**Purpose**

Antihistamine

Nasal decongestant

**Use(s)**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies:
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages
- temporarily relieves sinus congestion and pressure
- temporarily restores freer breathing through the nose

**Warnings**

**Do not use**

- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have difficulty swallowing

**Ask a doctor before use if you have**

- heart disease
- thyroid disease
- glaucoma
- high blood pressure

- diabetes
- trouble urinating due to an enlarged prostate gland
- kidney disease. Your doctor should determine if you need a different dose.

**When using this product**

- **do not take more than directed**
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

**Stop use and ask doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

**If pregnant or breast-feeding**

ask a health professional before use.

**Keep out of reach of children**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not divide, crush, chew or dissolve the tablet; swallow tablet whole

adults and children 12 years of age and over	take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
children under 12 years of age	do not use
adults 65 years of age and older	ask a doctor
consumers with kidney disease	ask a doctor

**Other information**

- safety sealed: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° to 25°C (68° to 77°F) store between 20° to 25°C (68° to 77°F)
- this product meets the requirements of USP dissolution test 3.

**Inactive ingredients**

corn starch, croscarmellose sodium, colloidal silicon dioxide, ferric oxide, hypromellose, kollidon SR, magnesium stearate, mannitol, powder cellulose and triethyl citrate.

Questions?

Call 1-888-375-3784

Principal Display Panel

**Drug Facts (continued)**

**Ask a doctor before use if you have** ■ heart disease ■ thyroid disease ■ glaucoma ■ high blood pressure ■ diabetes ■ kidney disease. Your doctor should determine if you need a different dose.

**When using this product:**

- do not take more than directed
- do not take at the same time as aluminum or magnesium antacids
- do not take with fruit juices (see Directions)
- the tablet coating may be seen in the stool (this is normal). Continue to take as directed (see Directions).

**Stop use and ask a doctor if**

- an allergic reaction to this product occurs. Seek medical help right away.
- symptoms do not improve within 7 days or are accompanied by a fever
- you get nervous, dizzy, or sleepless

**Before use,**

- Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).

**Drug Facts (continued)**

**Directions**

- do not use if safety sealed: do not use if carton is opened or if individual blister units are torn or opened
- store between 20° and 25°C (68° and 77°F)
- this product meets the requirements of USP Dissolution Test 3

**Other information**

- children under 12 years of age do not use
- adults and children 12 years of age and older: take 1 tablet with a glass of water every 12 hours on an empty stomach; do not take more than 2 tablets in 24 hours
- ask a doctor if you have difficulty swallowing

**Questions ? 1-888-287-1915**

TAMPER EVIDENT: INDIVIDUAL BLISTER UNIT SEALED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL BLISTER UNIT IS OPEN OR TORN. IMPORTANT: Read the directions and warnings before use. Keep the carton. It contains important information.

**Drug Facts**

**Active ingredients (in each tablet)**

**Purpose**

**Warnings**

- Do not use if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or 2 weeks after stopping the MAOI drug.
- MAOI, ask a doctor or pharmacist before taking this product.

**Uses**

- temporarily relieves these symptoms due to hay fever or other upper respiratory allergies: sneezing; runny nose; itchy, watery eyes; itching of the nose or throat
- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- reduces swelling of nasal passages

**Drug Facts (continued)**

**Directions**

- temporarily relieves sinus congestion and pressure
- temporarily relieves nasal breathing through the nose



NON-DROWSY

Allergy Relief D

Fexofenadine HCl 60mg/Antihistamine

Pseudoephedrine HCl 120mg/Nasal Decongestant

Extended-Release Tablets USP

Allergy & Congestion Indoor/Outdoor Relief of:

- Nasal and sinus congestion due to colds or allergies
- Sneezing
- Runny nose
- Itchy, watery eyes
- Itchy nose or throat due to allergies



20 EXTENDED RELEASE TABLETS

Compare to Allegra-D® 12 Hour Allergy & Congestion Tablets active ingredients\*

Satisfaction guaranteed - Or we'll replace it or give you your money back. For questions or comments or to report an undesired reaction or side effect, please call 1-888-287-1915.

DISTRIBUTED BY: Wal-Mart Inc., Bentonville, AR 72716 PRODUCT OF INDIA \*This product is not manufactured or distributed by Chatham, Inc. (part of the Sanofi Group), distributor of Allegra-D® 12 Hour Allergy & Congestion Tablets. Allegra-D® is a registered trademark of Aventis Inc.



how2recycle.info

LOT EXP



## FEXOFENADINE HCL AND PSEUDOEPHEDRINE HCl

fexofenadine hcl and pseudoephedrine hci tablet, extended release

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49035-273
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>FEXOFENADINE HYDROCHLORIDE</b> (UNII: 2S068B75ZU) (FEXOFENADINE - UNII:E6582LOH6V)	FEXOFENADINE HYDROCHLORIDE	60 mg
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	120 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>MANNITOL</b> (UNII: 3OWL53L36A)	
<b>POWDERED CELLULOSE</b> (UNII: SMD1X3XO9M)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>CROSCARMELLOSE SODIUM</b> (UNII: M28OL1HH48)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>FERRIC OXIDE RED</b> (UNII: 1K09F3G675)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)	

### Product Characteristics

<b>Color</b>	WHITE (buff white to pale yellow color and other layer light red to red color)	<b>Score</b>	no score
<b>Shape</b>	CAPSULE	<b>Size</b>	16mm
<b>Flavor</b>		<b>Imprint Code</b>	R;195
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:49035-273-20	4 in 1 CARTON	12/31/2019	
1		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:49035-273-30	6 in 1 CARTON	12/31/2019	
2		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA076667	12/31/2019	

**Labeler** - Wal-mart Stores Inc. (051957769)

Revised: 9/2021

Wal-mart Stores Inc.