

**HYDROGEN PEROXIDE 3 PERCENT- hydrogen peroxide liquid
Cardinal Health (Leader) 70000**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient

Hydrogen peroxide 3%

Purpose

First aid antiseptic/oral debriding agent

Uses

- first aid to help prevent the risk of infection in minor cuts, scrapes, and burns
- aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth

Warnings

For external use only.

Do not use

- in the eyes or apply over large areas of the body
- longer than one week

Ask a doctor before use if you have

- deep puncture wounds, animal bites or serious burns

Stop use and ask a doctor if

- swelling, rash, or fever develops
- the condition persists or gets worse
- sore mouth symptoms do not improve in 7 days
- irritation, pain, or redness persists or worsens

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

For use as a first aid antiseptic:

- clean affected area
- if bandaged, let dry first
- apply small amount of product on affected area 1-3 times a day
- may be covered with a sterile bandage

For use as an oral debriding agent (oral rinse):

- **adults and children 2 years of age and over:**
 - mix with an equal amount of water
 - swish around in the mouth over affected area for at least 1 minute and then spit it out. Do not swallow.
 - use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor
- **children under 12 years** should be supervised in the use of this product
- **children under 2 years of age:** consult a doctor or dentist

Other information

- keep tightly closed and in a dark place at room temperature
- do not shake bottle. Hold away from face when opening

Inactive ingredient

purified water

Questions or comments?

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

Principal Display Panel

Hydrogen Peroxide

3% USP

First Aid Antiseptic

FL OZ (mL)

**TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS
BROKEN OR MISSING.**

DISTRIBUTED BY CARDINAL HEALTH, DUBLIN OHIO 43017

www.myleader.com

Package Label

LEADERTM

NDC 70000-0500-1

Hydrogen Peroxide

3% USP
First Aid Antiseptic
Oral Debriding Agent

100% Money Back Guarantee

32 FL OZ (1 QT) 946 mL

PLD-A220C LB006564

Drug Facts

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING.

Active ingredient Hydrogen peroxide 3%.....	Purpose First aid antiseptic/oral debriding agent
Uses <ul style="list-style-type: none"> ■ first aid to help prevent the risk of infection in minor cuts, scrapes, and burns ■ aids in the removal of phlegm, mucus, or other secretions associated with occasional sore mouth 	
Warnings For external use only. <ul style="list-style-type: none"> Do not use <ul style="list-style-type: none"> ■ in the eyes or apply over large areas of the body ■ longer than one week Ask a doctor before use if you have <ul style="list-style-type: none"> ■ deep puncture wounds, animal bites or serious burns Stop use and ask a doctor if <ul style="list-style-type: none"> ■ swelling, rash, or fever develops ■ the condition persists or gets worse ■ sore mouth symptoms do not improve in 7 days ■ irritation, pain, or redness persists or worsens 	
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center (1-800-222-1222) right away.	
Directions For use as a first aid antiseptic: <ul style="list-style-type: none"> ■ clean affected area ■ if bandaged, let dry first ■ apply small amount of product on affected area 1-3 times a day ■ may be covered with a sterile bandage For use as an oral debriding agent (oral rinse): <ul style="list-style-type: none"> ■ adults and children 2 years of age and over: <ul style="list-style-type: none"> ■ mix with an equal amount of water ■ swish around in the mouth over affected area for at least 1 minute and then spit out. Do not swallow. ■ use up to 4 times daily after meals and at bedtime or as directed by a dentist or doctor ■ children under 12 years should be supervised in the use of this product ■ children under 2 years of age: consult a doctor or dentist 	
Other information <ul style="list-style-type: none"> ■ keep tightly closed and in a dark place at room temperature ■ do not shake bottle. Hold away from face when opening. 	
Inactive ingredient purified water	
Questions or comments? Call 1-877-753-3935 Monday-Friday 9AM-5PM EST	

DISTRIBUTED BY CARDINAL HEALTH, DUBLIN, OHIO 43017
www.myleader.com 1-800-200-6313
© 2019 Cardinal Health
PLD-A220C LB006571 **CIN 5511712** REV. 2/19

0 96295 13703 3

LEADER Hydrogen Peroxide 3% USP

HYDROGEN PEROXIDE 3 PERCENT

hydrogen peroxide liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70000-0500
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
HYDROGEN PEROXIDE (UNII: BBX060AN9V) (HYDROGEN PEROXIDE - UNII:BBX060AN9V)	HYDROGEN PEROXIDE	3 mL in 100 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70000-0500-1	946 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/28/2019	
2	NDC:70000-0500-2	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/28/2019	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	02/28/2019	

Labeler - Cardinal Health (Leader) 70000 (063997360)

Revised: 10/2021

Cardinal Health (Leader) 70000