

**ACETAMINOPHEN EXTRA STRENGTH- acetaminophen tablet, coated
FSA STORE INC.**

1170-CRM-2024-0730

Drug Facts

Active ingredient (in each gelcap)

Acetaminophen 500 mg

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - the common cold
 - headache
 - backache
 - minor pain of arthritis
 - toothache
 - muscular aches
 - premenstrual and menstrual cramps
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or

pharmacist.

- if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see overdose warning)**

adults and children 12 years and over	<ul style="list-style-type: none">▪ take 2 gelcaps every 6 hours while symptoms last▪ do not take more than 6 gelcaps in 24 hours, unless directed by a doctor▪ do not use for more than 10 days unless directed by a doctor
children under 12 years	<ul style="list-style-type: none">▪ ask a doctor

Other information

- store between 20-25°C (68-77°F)
- retain carton for complete product information

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, D&C red #33, edible ink, FD&C blue #1, FD&C red #40, FD&C yellow #6, gelatin, hydroxypropyl cellulose, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, polyethylene glycol, povidone,

pregelatinized starch, stearic acid, titanium dioxide

PRINCIPAL DISPLAY PANEL

Caring Mill™

NDC 81522-070-13

†Compare to the active ingredient in Tylenol® Extra Strength Rapid Release Gels

Extra Strength

Acetaminophen

500 mg

Pain Reliever/Fever Reducer

For Adults

225 GELCAPS

500 MG EACH

Actual Size

DO NOT USE IF IMPRINTED SEAL UNDER CAP IS BROKEN OR MISSING

NDC 81522-070-13

caring mill™

Compare to the active ingredient in Tylenol® Extra Strength Rapid Release Gels

**Extra Strength
Acetaminophen
500 mg
Pain Reliever/Fever Reducer**

**225 GELCAPS
500 MG EACH**

For Adults

Actual Size

Drug Facts

Active ingredient (in each gelcap)
Acetaminophen 500 mg

Purpose
Pain reliever/fever reducer

Uses
temporarily relieves minor aches and pains due to:
the common cold ■ headache ■ backache ■ minor pain of arthritis
toothache ■ muscular aches ■ premenstrual and menstrual cramps
temporarily reduces fever

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
more than 4,000 mg of acetaminophen in 24 hours
with other drugs containing acetaminophen
3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:
skin reddening ■ blisters ■ rash
if a skin reaction occurs, stop use and seek medical help right away.

Do not use
with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
if you are allergic to acetaminophen or any of the inactive ingredients in this product

Ask a doctor before use if you have liver disease.

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if
pain gets worse or lasts more than 10 days
fever gets worse or lasts more than 3 days
new symptoms occur
redness or swelling is present
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
do not take more than directed (see overdose warning)
adults and children 12 ■ take 2 gelcaps every 6 hours while symptoms last
years and over ■ do not take more than 6 gelcaps in 24 hours, unless directed by a doctor
children under 12 years ■ do not use for more than 10 days unless directed by a doctor ■ ask a doctor

Other information
store between 20-25°C (68-77°F)

Inactive ingredients
colloidal silicon dioxide, croscarmellose sodium, D&C red #33, edible ink, FD&C blue #1, FD&C red #40, FD&C yellow #6, gelatin, hydroxypropyl cellulose, hypromellose, iron oxide black, iron oxide red, iron oxide yellow, polyethylene glycol, povidone, pregelatinized starch, stearic acid, titanium dioxide

L117035291_01

Manufactured for:
FSA Store Inc.
5712 W. 10th Rd
Suite 100
PMB 24308
Dallas, Texas 75231
Customer Care Help Line
1-844-705-4434
Made in India

This product is not manufactured or distributed in the United States by the manufacturer of Tylenol Extra Strength Rapid Release Gels.

0 199581 191233 18

ACETAMINOPHEN EXTRA STRENGTH

acetaminophen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:81522-070
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
D&C RED NO. 33 (UNII: 9DBA0SBB0L)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GELATIN (UNII: 2G86QN327L)	
HYDROXYPROPYL CELLULOSE (1600000 WAMW) (UNII: RFW2ET671P)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	gray (red and light blue ends)	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	G1
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:81522-070-13	225 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	07/30/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/30/2024	

Labeler - FSA STORE INC. (049283340)