

**NICOTINE- nicotine polacrilex gum, chewing**  
**Rite Aid Corporation**

-----  
**Drug Facts**

**Active ingredient (in each chewing piece)**

Nicotine polacrilex 4 mg

**Purpose**

Stop smoking aid

**Use**

- reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

**Warnings**

**If you are pregnant or breast-feeding,**

**only use this medicine on the advice of your health care provider.** Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

**Ask a doctor before use if you have**

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase your blood pressure.
- stomach ulcer or diabetes
- history of seizures

**Ask a doctor or pharmacist before use if you are**

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

**Stop use and ask a doctor if**

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness,

diarrhea, weakness and rapid heartbeat

- you have symptoms of an allergic reaction (such as difficulty breathing or rash)

### **Keep out of reach of children and pets.**

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### **Directions**

- **if you are under 18 years of age, ask a doctor before use**
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- **if you smoke your first cigarette within 30 minutes of waking up**, use 4 mg nicotine gum according to the following 12 week schedule:
- **if you smoke your first cigarette more than 30 minutes after waking up**, use 2 mg nicotine gum

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat the process until most of the tingle is gone (about 30 minutes)
- do not eat or drink 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause you hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- It is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

### **Other information**

- **each piece contains:** calcium 110 mg, sodium 11 mg
- store at 20-25°C (68-77°F)
- protect from light

### **Inactive ingredients**

acesulfame potassium, butylated hydroxytoluene, carnauba wax, D&C yellow #10 lake, FD&C blue #2 lake, FD&C red #40, FD&C Yellow #6 lake, flavors, gum base, sodium

bicarbonate, sodium carbonate, sorbitol, talc

### **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

Compare to the active ingredient of Nicorette® Gum\*

FREE | SUGAR

FROM | FREE

### **Nicotine Gum**

Nicotine Polacrilex Gum USP, 4 mg (nicotine)

Stop Smoking Aid

ORIGINAL FLAVOR

### **FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE MORE THAN 30 MINUTES OF WAKING UP.**

If you smoke your first cigarette **WITHIN 30 MINUTES** after waking up, use Nicotine Polacrilex Gum USP, 2 mg

### **TO INCREASE YOUR SUCCESS IN QUITTING:**

- 1.** You must be motivated to quit.
- 2.** Use Enough - Chew at least **9** pieces of Nicotine Polacrilex Gum per day during the first six weeks.
- 3.** Use Long Enough - Use Nicotine Polacrilex Gum for the full 12 weeks.
- 4. Use With a Support Program** as directed in the enclosed User's Guide.

To remove the gum, tear off single unit.

Peel off backing starting at corner with loose edge.

Push gum through foil

- **NOT FOR SALE TO THOSE UNDER 18 YEARS OF AGE**
- **PROOF OF AGE REQUIRED**
- **NOT FOR SALE IN VENDING MACHINES OR FROM ANY SOURCE WHERE PROOF OF AGE CANNOT BE VERIFIED**

\*This product is not manufactured or distributed by GlaxoSmithKline Consumer Healthcare, L.P., distributor of Nicorette® Gum

**See inside for more details or visit [www.nitthehabit.com](http://www.nitthehabit.com)**

**BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN**

**DISTRIBUTED BY:**

RITE AID, 30 HUNTER LANE,  
CAMP HILL, PA 17011  
www.riteaid.com

## Package Label

**TO INCREASE YOUR SUCCESS IN QUITTING:**

1. You must be motivated to quit.
2. **Use Enough** - Chew at least **9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks.
3. **Use Long Enough** - Use Nicotine Polacrilex Gum for the full 12 weeks.
4. **Use With a Support Program** as directed in the enclosed User's Guide.



**FREE FROM SUGAR FREE**

NDC 11822-3323-2

Compare to the active ingredient of **Nicorette® Gum\***

# Nicotine Gum

Nicotine Polacrilex Gum USP,  
4 mg (nicotine)  
Stop Smoking Aid

**4 mg**

**FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP**

If you smoke your first cigarette **MORE THAN 30 MINUTES** after waking up, use Nicotine Polacrilex Gum USP, 2 mg sugar free uncoated nicotine gum



ACTUAL SIZE  
ORIGINAL FLAVOR

**170**  
PIECES 4 mg EACH

NDC 11822-3323-2

**FREE FROM SUGAR FREE**

Compare to the active ingredient of **Nicorette® Gum\***

# Nicotine Gum

Nicotine Polacrilex Gum USP,  
4 mg (nicotine)  
Stop Smoking Aid

**4 mg**

If you're not satisfied, we'll happily refund your money.



**Nicotine Gum**  
4 mg (nicotine)  
Nicotine Polacrilex Gum USP, Stop Smoking Aid

**OUR PROMISE:**

Sugar Free  
Recyclable Material

**FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP**

If you smoke your first cigarette within 30 minutes of waking up, you are at a significantly higher risk of heart disease and stroke. Nicotine Polacrilex Gum USP, 4 mg is a proven, effective, safe, and pleasant way to reduce your nicotine craving and help you quit smoking.

**Directions**

If you smoke your first cigarette more than 30 minutes after waking up, you should use Nicotine Polacrilex Gum USP, 4 mg as directed. If you smoke your first cigarette within 30 minutes of waking up, you should use Nicotine Polacrilex Gum USP, 4 mg as directed. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are pregnant or breastfeeding. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: carbamazepine, cefepime, ceftriaxone, cefuroxime, dexamethasone, diltiazem, doxycycline, erythromycin, fluoxetine, gabapentin, gemfibrozil, hydrocodone, levofloxacin, linezolid, meprobamate, nifedipine, nortriptyline, omeprazole, ranitidine, ropinirole, telmisartan, venlafaxine, or zolpidem. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: bupropion, duloxetine, escitalopram, levamisole, norethisterone, or tramadol. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: acetaminophen, aspirin, ibuprofen, or naproxen. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: albuterol, beclomethasone, budesonide, fluticasone, or salmeterol. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: amoxicillin, ampicillin, azithromycin, cefadroxil, cefprozil, cephalexin, clarithromycin, clindamycin, erythromycin, levofloxacin, moxifloxacin, trimethoprim-sulfamethoxazole, or vancomycin. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: amlodipine, diltiazem, felodipine, isradipine, or nifedipine. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: atorvastatin, fluvastatin, or rosuvastatin. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: calcium channel blockers, digoxin, or antiarrhythmics. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: cholinesterase inhibitors, or tricyclic antidepressants. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: cyclosporin, cyclosporine, or tacrolimus. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: diuretics, epinephrine, guanethidine, guanidines, nifedipine, or sodium bicarbonate. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: fentanyl, morphine, or other opioids. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: fentanyl, morphine, or other opioids. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: fentanyl, morphine, or other opioids.

**Warnings**  
If you are pregnant or breastfeeding, only use this medicine on the advice of your health care provider. Nicotine Polacrilex Gum USP, 4 mg may cause dizziness, lightheadedness, or other symptoms of nicotine withdrawal. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: carbamazepine, cefepime, ceftriaxone, cefuroxime, dexamethasone, diltiazem, doxycycline, erythromycin, fluoxetine, gabapentin, gemfibrozil, hydrocodone, levofloxacin, linezolid, meprobamate, nifedipine, nortriptyline, omeprazole, ranitidine, ropinirole, telmisartan, venlafaxine, or zolpidem. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: bupropion, duloxetine, escitalopram, levamisole, norethisterone, or tramadol. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: acetaminophen, aspirin, ibuprofen, or naproxen. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: albuterol, beclomethasone, budesonide, fluticasone, or salmeterol. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: amoxicillin, ampicillin, azithromycin, cefadroxil, cefprozil, cephalexin, clarithromycin, clindamycin, erythromycin, levofloxacin, moxifloxacin, trimethoprim-sulfamethoxazole, or vancomycin. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: amlodipine, diltiazem, felodipine, isradipine, or nifedipine. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: atorvastatin, fluvastatin, or rosuvastatin. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: calcium channel blockers, digoxin, or antiarrhythmics. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: cholinesterase inhibitors, or tricyclic antidepressants. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: cyclosporin, cyclosporine, or tacrolimus. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: diuretics, epinephrine, guanethidine, guanidines, nifedipine, or sodium bicarbonate. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: fentanyl, morphine, or other opioids. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: fentanyl, morphine, or other opioids.

**Use**  
Nicotine Polacrilex Gum USP, 4 mg is a nicotine replacement therapy. It is not a cure for nicotine dependence. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: carbamazepine, cefepime, ceftriaxone, cefuroxime, dexamethasone, diltiazem, doxycycline, erythromycin, fluoxetine, gabapentin, gemfibrozil, hydrocodone, levofloxacin, linezolid, meprobamate, nifedipine, nortriptyline, omeprazole, ranitidine, ropinirole, telmisartan, venlafaxine, or zolpidem. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: bupropion, duloxetine, escitalopram, levamisole, norethisterone, or tramadol. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: acetaminophen, aspirin, ibuprofen, or naproxen. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: albuterol, beclomethasone, budesonide, fluticasone, or salmeterol. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: amoxicillin, ampicillin, azithromycin, cefadroxil, cefprozil, cephalexin, clarithromycin, clindamycin, erythromycin, levofloxacin, moxifloxacin, trimethoprim-sulfamethoxazole, or vancomycin. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: amlodipine, diltiazem, felodipine, isradipine, or nifedipine. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: atorvastatin, fluvastatin, or rosuvastatin. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: calcium channel blockers, digoxin, or antiarrhythmics. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: cholinesterase inhibitors, or tricyclic antidepressants. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: cyclosporin, cyclosporine, or tacrolimus. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: diuretics, epinephrine, guanethidine, guanidines, nifedipine, or sodium bicarbonate. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: fentanyl, morphine, or other opioids. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: fentanyl, morphine, or other opioids.

**Drug Facts (continued)**

Quantity	1 to 2 hours	2 to 4 hours	4 to 8 hours
Nicotine 1 to 2 mg	1 place every 2 hours	1 place every 4 hours	1 place every 8 hours
Nicotine 2 to 4 mg	1 place every 2 hours	1 place every 4 hours	1 place every 8 hours

If you smoke your first cigarette within 30 minutes of waking up, use 4 mg Nicotine Polacrilex Gum USP, 4 mg every 2 to 4 hours, then every 4 to 8 hours. If you smoke your first cigarette more than 30 minutes after waking up, use 2 mg Nicotine Polacrilex Gum USP, 4 mg every 2 to 4 hours, then every 4 to 8 hours. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: carbamazepine, cefepime, ceftriaxone, cefuroxime, dexamethasone, diltiazem, doxycycline, erythromycin, fluoxetine, gabapentin, gemfibrozil, hydrocodone, levofloxacin, linezolid, meprobamate, nifedipine, nortriptyline, omeprazole, ranitidine, ropinirole, telmisartan, venlafaxine, or zolpidem. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: bupropion, duloxetine, escitalopram, levamisole, norethisterone, or tramadol. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: acetaminophen, aspirin, ibuprofen, or naproxen. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: albuterol, beclomethasone, budesonide, fluticasone, or salmeterol. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: amoxicillin, ampicillin, azithromycin, cefadroxil, cefprozil, cephalexin, clarithromycin, clindamycin, erythromycin, levofloxacin, moxifloxacin, trimethoprim-sulfamethoxazole, or vancomycin. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: amlodipine, diltiazem, felodipine, isradipine, or nifedipine. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: atorvastatin, fluvastatin, or rosuvastatin. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: calcium channel blockers, digoxin, or antiarrhythmics. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: cholinesterase inhibitors, or tricyclic antidepressants. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: cyclosporin, cyclosporine, or tacrolimus. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: diuretics, epinephrine, guanethidine, guanidines, nifedipine, or sodium bicarbonate. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: fentanyl, morphine, or other opioids. Do not use Nicotine Polacrilex Gum USP, 4 mg if you are taking any of the following medications: fentanyl, morphine, or other opioids.

**Questions or comments?**  
For more information, call 1-800-845-2333. Visit our website at [www.rteaid.com](http://www.rteaid.com).

**Ingredients**  
Nicotine Polacrilex Gum USP, 4 mg contains the following ingredients: Nicotine Polacrilex Gum USP, 4 mg, Sugarfree Gum Base, Hydroxytoluene, Potassium Acesulfame, Carnauba Wax, Talc, D & C Yellow No. 10, FD & C Blue No. 2, FD & C Red No. 40, FD & C Yellow No. 6, and Nicotine Polacrilex Gum USP, 4 mg.

**Distributed by:**  
RITE AID, 30 HUNTER LANE  
CAMP HILL, PA 17011  
www.rteaid.com

**ARTIFACT**

## RITE AID Nicotine Gum Original Flavor

### NICOTINE

nicotine polacrilex gum, chewing

#### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-3323
Route of Administration	ORAL		

#### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	4 mg

#### Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 230V73Q5G9)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SODIUM BICARBONATE (UNII: 8MDF5V39Q0)	
SODIUM CARBONATE (UNII: 45P3261C7T)	
TALC (UNII: 7SEV7J4R1U)	
SORBITOL (UNII: 506T60A25R)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

**Product Characteristics**

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	SQUARE	<b>Size</b>	14mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-3323-2	170 in 1 CARTON	01/31/2020	
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:11822-3323-1	110 in 1 CARTON	01/31/2020	
2		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA074707	01/31/2020	

**Labeler** - Rite Aid Corporation (014578892)

Revised: 1/2022

Rite Aid Corporation