

**NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate capsule, liquid filled
Lil' Drug Store Products, Inc.**

Nighttime Cold & Flu

Drug Facts

<i>Active ingredients (in each softgel)</i>	<i>Purposes</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 15 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Antihistamine

Uses

- temporarily relieves common cold and flu symptoms:
 - sore throat
 - headache
 - minor aches and pains
 - fever
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks daily while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by

fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

Other information

- store between 15- 30°C (59- 86°F)
- avoid excessive heat

Inactive ingredients

D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or comments?

call toll-free **1-877-507-6516** (M-F 8AM-4:30PM CST)

Distributed by:

Lil' Drug Store Products, Inc.
9300 Earhart Lane SW
Cedar Rapids, IA 52404

PRINCIPAL DISPLAY PANEL - 16 Softgel Blister Pack Carton

QUALITY

* GUARANTEED *

Compare to the Active Ingredients in
Vicks® NyQuil® LiquiCaps®**

Cold & Flu
Nighttime

Acetaminophen, Dextromethorphan HBr, and Doxylamine Succinate
Pain Reliever/Fever Reducer, Cough Suppressant, Antihistamine

Relieves

- Sneezing • Runny Nose
- Cough • Aches
- Fever • Sore Throat

ACTUAL SIZE

16
Softgels

Lil'
DrugStore®

Cold & Flu Nighttime



Compare to the Active Ingredients in
Vicks® NyQuil® LiquiCaps®**

Cold & Flu Nighttime

Acetaminophen, Dextromethorphan HBr, and Doxylamine Succinate
Pain Reliever/Fever Reducer, Cough Suppressant, Antihistamine

Relieves

- Sneezing • Runny Nose
- Cough • Aches
- Fever • Sore Throat



ACTUAL SIZE



16 Softgels

Cold & Flu Nighttime

Acetaminophen, Dextromethorphan HBr, and Doxylamine Succinate
Pain Reliever/Fever Reducer, Cough Suppressant, Antihistamine

Drug Facts (continued)

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

Other information

- store between 15-30°C (59-86°F)
- avoid excessive heat

Inactive ingredients

D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

Questions or comments?

call toll-free 1-877-507-6516
(M-F 8AM-4:30PM CST)

**This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil® and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

16
Softgels



68168C-US-10-19-PA


Distributed by:
Lil' Drug Store Products, Inc.
9300 Earhart Lane SW
Cedar Rapids, IA 52404

Softgels are
liquid-filled
capsules.



PLD-A569B FC006104

Lot No:
Exp Date:



PRINT SIDE GLUE

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Drug Facts		Drug Facts (continued)
Active ingredients (in each softgel)	Purposes	an MAOI, ask a doctor or pharmacist before taking this product.
Acetaminophen 325 mg.....Pain reliever/fever reducer		Ask a doctor before use if you have
Dextromethorphan HBr 15 mg.....Cough suppressant		• liver disease • glaucoma
Doxylamine succinate 6.25 mg.....Antihistamine		• trouble urinating due to an enlarged prostate gland
		• a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema
Uses		• cough that occurs with too much phlegm (mucus)
• temporarily relieves common cold and flu symptoms:		Ask a doctor or pharmacist before use if you are
• sore throat • headache • minor aches and pains		• taking the blood thinning drug warfarin
• fever • runny nose and sneezing		• taking sedatives or tranquilizers
• cough due to minor throat and bronchial irritation		When using this product
Warnings		• do not exceed recommended dosage
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:		• excitability may occur, especially in children
• more than 4,000 mg of acetaminophen in 24 hours		• avoid alcoholic drinks • marked drowsiness may occur
• with other drugs containing acetaminophen		• alcohol, sedatives, and tranquilizers may increase drowsiness
• 3 or more alcoholic drinks daily while using this product		• be careful when driving a motor vehicle or operating machinery
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening		Stop use and ask a doctor if
• blisters • rash If a skin reaction occurs, stop use and seek medical help right away.		• pain or cough gets worse or lasts more than 7 days
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.		• fever gets worse or lasts more than 3 days
Do not use		• redness or swelling is present • new symptoms occur
• with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.		• cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains		If pregnant or breast-feeding, ask a health professional before use.
		Keep out of reach of children.
		Overdose warning: Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

NIGHTTIME COLD AND FLU			
acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate capsule, liquid filled			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66715-6716

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
DOXYLAMINE SUCCINATE (UNII: V9BI9B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

Inactive Ingredients

Ingredient Name	Strength
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
GELATIN, UNSPECIFIED (UNII: 2G86QN327L)	
GLYCERIN (UNII: PDC6A3C0OX)	
MANNITOL (UNII: 3OWL53L36A)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITAN (UNII: 6O92ICV9RU)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	green	Score	no score
Shape	OVAL	Size	20mm
Flavor		Imprint Code	P30
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66715-6716-8	8 in 1 CARTON	02/15/2020	
1		2 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/05/2014	

Revised: 12/2025

Lil' Drug Store Products, Inc.