

JOHNSON FOAM FOR MEN- minoxidil aerosol, foam
JOHNSON BIOPHARMACEUTICAL INC.

JOHNSON FOAM FOR MEN

MINOXIDIL 5%
FINASTERIDE 0.1%

Fight primary cause of hair loss in men. Anti hair loss to help increase hair density and volume.

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Fight primary cause of hair loss in men. Anti hair loss to help increase hair density and volume.

do not apply on other parts of the body.

avoid contact with eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water.

some people have experienced changes in hair color and/or texture.

For external use only.

For use by men only.

Extremely Flammable

Avoid fire, flame, or smoking during and immediately following application.

1. PREP HANDS WITH COLD WATER. Rinse Your Fingers In Cold Water And Dry Them. note: Foam Will Melt On Contact With Warm Surfaces.

2. OPEN CHILD RESISTANT CAP

Be Sure To Align Arrow On The Cap With Arrow On Blue Ring. Tilt Cap Back And Pull Off.

3. DISPENSE VERTICALLY ON TO A COLD SURFACE

Hold The Can Straight Upside Down. If You Hold The Can At An Angle, Foam May Not Dispense Properly. Press Nozzle To Dispense Half A Capful Of Foam Onto Cold Surface (e.g. dish or Cold Hands).

4. APPLY TO SCALP. NOT HAIR

Part Your Hair To Expose Hair Loss Area. Massage Foam Into Scalp Not Hair, Repeat Until All Hair Loss Areas Have Been Covered.

5. CLOSE CAP AND WASH HANDS. Snap cap back into place be sure arrows do not line up, so cap remains child resistant, wash hands and any surface thoroughly after use.

Store at controlled room temperature 20 °C to 25 °C (68° to 77 F). Contents under pressure. Do not puncture or incinerate container. Do not expose to heat or store at temperatures above 120 F (49 °C).

ALCOHOL

GLYCERIN

WATER

CETYL PEG/PPG-10/1 DIMETHICONE (HLB 5)

HYDROLYZED KERATIN

UBIQUINOL
BIOTIN
ZINC PIDOLATE
BUTYLATED HYDROXYTOLUENE
EDETATE DISODIUM
LEVOMENTHOL
Poloxzam
POLYSORBATE 80

包装彩盒:70*65*185mm
工艺:银卡+ 逆向UV

激凸

JOHNSON



DRUG FACTS

Active Ingredient(s)
Minoxidil(5%), Finasteride(0.1%)

Purpose:
Treats primary cause of hair loss in men. Anti hair loss to help increase hair density and volume.

Warnings:
• For external use only.
• For use by men only.
• Pregnancy: Finasteride.
• Avoid fire, flame, or smoking during and immediately following application.

Do not use if:
• you have not family history of hair loss,
• your hair loss is sudden and/or patchy,
• you do not know the reason for your hair loss,
• you are and 18 years of age,
• Do not use on broken or chapped skin,
• your scalp is red, inflamed, infected, irritated, or painful,
• you use other medicines on the scalp.

Ask a doctor before use if you have
heart disease.

When using this product
• do not apply on other parts of the body.
• avoid contact with eyes. In case of accidental contact, rinse eyes with large amounts of cool tap water.
• some people have experienced changes in hair color and/or texture.

Stop use and ask a doctor if:
• chest pain, rapid heartbeat, faintness, or dizziness occurs.
• sudden, unexplained weight gain occurs.
• your hands or feet swell.
• scalp irritation or redness occurs.
• unwanted facial hair growth occurs.
• you do not see hair regrowth in 4 months.

May be harmful if used when pregnant or breast-feeding.
If swallowed, get medical help or contact a Poison Control Center right away.

Company information
Name: Johnson Biopharmaceutical, Inc.
Address: English Park, GreenTree Centre 525 ROUTE 100, FORT LINDSEY, VA 22036
MVA#: 4000170000, Johnson@jbi.com

Finasteride, a specific inhibitor of type II 5 α -reductase, competitively inhibits the conversion of testosterone to dihydrotestosterone in the scalp of men, thereby serving as a treatment for male pattern hair loss. Unlike oral finasteride, topical formulations enable precise and direct application to the scalp surface, ensuring high local drug concentration and minimizing side effects commonly associated with systemic administration. Furthermore, topical formulations reduce systemic absorption of finasteride, limiting exposure outside the targeted area.

Inactive ingredients:
Alcohol, Glycerin, Water, Dimethicone, Hydrolyzed keratin, Ubiquinol, Zinc pidolate, Butylated hydroxytoluene, Edetate disodium, Levomenthol, Poloxzam, Polysorbate 80.

Store at controlled room temperature 20° C to 25° C (68° F to 77° F). Contents under pressure. Do not puncture or incinerate container. Do not expose to heat or store at temperatures above 120° (49° C).

THIS PACKAGE CONTAINS:



One 90g Topical Aerosol



JOHNSON BIOPHARMACEUTICAL

DERMATOLOGIST RECOMMENDED BRAND

FINASTERIDE FOAM WITH 5% MINOXIDIL

Hair Regrowth Treatment



REACTIVATES HAIR FOLLICLES TO STIMULATE REGROWTH

CLINICALLY PROVEN TO REGROW HAIR

Easy-to-use Foam
90 Days Supply
One 90 g (3.03oz) Can

DIRECTIONS FOR USE

See information booklet with capsule directions on how to use and obtain best results.

1. PREP HANDS WITH COLD WATER
Wash Your Hands in Cold Water. And Dry Thoroughly. Foam Will Not Set On Contact With Warm Surface.

2. OPEN CHILD RESISTANT CAP
Be Sure To Align Arrow On The Cap With Arrow On Blue Ring. Lift Cap Back And Pull Off.

3. DISPENSE VERTICALLY ONTO A COLD SURFACE
Hold The Can Straight Up, Push Down. If You Hold The Can At An Angle, Foam May Not Dispense Properly. Press Nozzle To Dispense. Hold A Card Or Paper On Cold Surface (e.g. Dish or Cold Hands).

4. APPLY TO SCALP, NOT HAIR
Part Your Hair To Expose Hair Loss Area. Massage Foam Into Scalp. Not Hair. Repeat Until All Hair Loss Areas Have Been Covered.



5. CLOSE CAP AND WASH HANDS
Always keep cap on. Cap should be kept closed to prevent foam from drying. After use, wash hands and any surface thoroughly after use.

Read and keep package and booklet for full product information

NDC 84539-001-01



丝印:125*130mm
 正反面印 ■印金
 红框丝印范围



瓶盖黑色半透

瓶身黑色

JOHNSON FOAM FOR MEN

minoxidil aerosol, foam

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84539-001
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FINASTERIDE (UNII: 57GNO57U7G) (FINASTERIDE - UNII:57GNO57U7G)	FINASTERIDE	0.1 g in 100 g
MINOXIDIL (UNII: 5965120SH1) (MINOXIDIL - UNII:5965120SH1)	MINOXIDIL	5 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
EDETATE DISODIUM (UNII: 7FLD91C86K)	
ALCOHOL (UNII: 3K9958V90M)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0K00R)	
COCODIMONIUM HYDROXYPROPYL HYDROLYZED KERATIN (1000 MW) (UNII: 8V0I3U3HMO)	
UBIQUINOL (UNII: M9NLOC577Y)	
BIOTIN (UNII: 6S06U10H04)	
ZINC PIDOLATE (UNII: C32PQ86DH4)	
BUTYLATED HYDROXYTOLUENE (UNII: 1P9D0Z171K)	

CETYL PEG/PPG-10/1 DIMETHICONE (HLB 5) (UNII: 035JKJ76MT)

LEVOMENTHOL (UNII: BZ1R15MTK7)

POLOXAMER 407 (UNII: TUF2IVW3M2)

POLYSORBATE 80 (UNII: 6OZP39ZG8H)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84539-001-01	90 g in 1 BOTTLE; Type 0: Not a Combination Product	07/15/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M032	07/15/2024	

Labeler - JOHNSON BIOPHARMACEUTICAL INC. (132195896)

Establishment

Name	Address	ID/FEI	Business Operations
JOHNSON BIOPHARMACEUTICAL INC.		132195896	manufacture(84539-001)

Revised: 7/2024

JOHNSON BIOPHARMACEUTICAL INC.