CITROMA- magesium citrate liquid Wal-Mart Stores, Inc

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active ingredient

Magnesium citrate 1.745g per fl oz

Purpose

Saline laxative

use

for relief of occasional constipation (irregularity). Generally produces bowel movement in 1/2 to 6 hours

Warnings

Warnings

Ask A Doctor

Ask a doctor before use if you have

- kidney disease
- a magnesium or sodium restricted diet
- abdorominal pain, nausea, or vomiting
- noticed a sudden change in bowel habits tat persists over a period of 2 weeks
- already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement afte use. These could be signs of a serious condition.

if pregnant or breast-feeding

If pregnant or breast-feeding, ask help professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- drink a full glass (8 ounces) of liquid with each dose
- may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over - 6.5 to 10 fl oz in 24 hours children 6 to under 12 years of age - 3 to 7 fl oz maximum 7 fl oz in 24 hours children 2 to under 6 years of age - 2 to 3 fl oz children under 2 years of age - ask a doctor discard unused product within 24 hours of opening bottle

other information

Other information

- magnesium content 290 mg per 1 fl oz (30 mL)
- sodium content 48 mg per 1 fl oz (30 mL)
- store at temperatures between 46° and 86° F (8° and 30° C)

Inactive ingredients

citric acid, flavor, sodium bicarbonate, sodium saccharin, water

Questions?

Questions? 1-888-287-1915

Adverse Reaction

Distributed by: Wal-Mart Stores Inc., Bentonville, AR 72716 525.000/525AA

principal display panel

Do not use if tamper evident twist-off cap is missing, broken or separated from neckring

Equate

Sugar Free

Magnesium

Citrate

U.S.P. Oral Solution

- Relief of occasional constipation
- Bowel cleansing for surgery or exam preparation
- Pasteurized

Grape

Flavor

10 FL OZ (296 mL

Do not use if tamper evident twist-off cap is missing, broken or separated from neckring



Sugar Free Magnesium

Citrate **U.S.P. Oral Solution**

Saline Laxative

- Relief of occasional constinution
- Bowel cleansing for surgery or exam preparation
- Pasteurized

10 FL OZ (296 mL)

Drug Facts

Active ingredient

Magnesium citrate 1.745 g per fl oz...

Purpose Saline laxative

Use for relief of occasional constigation (regularity), Generally produces bowel movement in 1/2 to 6 hours.

Ask a doctor before use if you have

🛮 kidney disease 🔳 a magnesium or sodium-restricted diet 🗎 abdominal pain, nausea, or vomiting
roticed a sudden change in bowel habits that persists over a period of 2 weeks 🖪 already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work,

Step use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions adrink a full glass (8 ounces) of liquid with each dose

may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over | 6.5 to 10 fl oz maximum 10 fl oz in 24 hours 3 to 7 fl oz maximum 7 fl oz in 24 hours children 6 to under 12 years of age children 2 to under 6 years of age 2 to 3 flioz maximum 3 flioz in 24 hours children under 2 years of age ask a doctor


discard unused product within 24 hours of opening bottle

Other information m magnesium content 290 mg per 1 fi oz (30 mL)

sodium content 48 mg per 1 fl oz (30 mL)
store at temperatures between 46° and 86° F (8° and 30° C)

Inactive ingredients citric acid, flavor, sodium bicarbonate, sodium saccharin, water

Questions? 1-888-287-1915



CITROMA

magesium citrate liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:49035-525

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM CITRATE (UNII: RHO26O1T9V) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM CITRATE	1.745 g in 29.6 mL

Inactive Ingredients			
Ingredient Name	Strength		
CITRIC ACID MONO HYDRATE (UNII: 2968 PHW8 QP)			
SODIUM BICARBONATE (UNII: 8MDF5V39QO)			
SACCHARIN SO DIUM (UNII: SB8 ZUX40 TY)			
WATER (UNII: 059QF0KO0R)			

Packaging			
# Itam Cada	Dackage Description	Marketing Start	Marketing End

#	# Item Code Package Description		Date	Date
1		296 mL in 1 BOTTLE, GLASS; Type 0: Not a Combination Product	08/29/1995	
Marketing Information				
	Marketing Catego	ry Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
О	TC monograph not f	inal part334	08/29/1995	

Labeler - Wal-Mart Stores, Inc (051957769)

Registrant - Vi-Jon (790752542)

Establishment				
Name	Address	ID/FEI	Business Operations	
Vi-Jon		790752542	manufacture(49035-525)	

Revised: 11/2019 Wal-Mart Stores, Inc