# DG HEALTH CHILDRENS PAIN AND FEVER- acetaminophen suspension Dolgencorp Inc

-----

# Dolgencorp, LLC Children's Pain & Fever Drug Facts

### Active ingredient (in each 5 mL)

Acetaminophen 160 mg

## **Purpose**

Pain reliever/fever reducer

#### Uses

temporarily:

- reduces fever
- relieves minor aches and pains due to:
- the common cold
- flu
- headache
- sore throat
- toothache

# **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

#### Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child has ever had an allergic reaction to this product or any of its ingredients

## Ask a doctor before use if your child has

liver disease

### Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin

# When using this product

do not exceed recommended dose (see overdose warning)

### Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present

These could be signs of a serious condition.

# Keep out of reach of children.

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical even if you do not notice any signs or symptoms.

#### **Directions**

- this product does not contain directions or complete warnings for adult use
- do not give more than directed (see overdose warning)
- shake well before using
- mL = milliliter
- find right dose on chart. If possible, use weight to dose; otherwise, use age.
- remove the child protective cap and squeeze your child's dose into the dosing cup
- repeat dose every 4 hours while symptoms last
- do not give more than 5 times in 24 hours

Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL

ı	-	İ
72-95	11 years	15 mL

<sup>\*</sup>or as directed by a doctor

**Attention:** use only enclosed dosing cup specifically designed for use with this product. Do not use any other dosing device.

#### Other information

- each 5 mL contains: sodium 3 mg
- store at 20-25°C (68-77°F)
- · do not use if printed neckband is broken or missing

### **Inactive ingredients**

anhydrous citric acid, calcium sulfate, carrageenan, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose, tribasic sodium phosphate

#### Questions or comments?

1-888-309-9030

# Package/Label Principal Display Panel

DG® health

Compare to the active ingredient of Children's Tylenol® Oral Suspension

For Ages 2 to 11 Years

Children's Pain & Fever

Acetaminophen 160 mg per 5 mL

Pain Reliever/Fever Reducer

Oral Suspension

Alcohol free

Aspirin free

Ibuprofen free


Paraben Free

Fast, effective relief of children's pain and fever

Dye Free

Cherry Flavor

8 FL OZ (237 mL)



# **DG**\* health

For Ages 2 to 11 Years

# Children's Pain &

Fever

Acetaminophen, 160 mg per 5 mL Pain Reliever/Fever Reducer Oral Suspension

# Drug Facts

Active ingredient (in each 5 mL)

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes
more than 5 doses in 24 hours, which is the maximum daily amount.

amount

with other drugs containing ocetaminophen
Allergy alert. Acetaminophen may cause severe skin reactions.
Symptome may include:

skin reddening blisters Blash
It a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If ore throat is severe, persists for more
ana 2 days, is ecomparied or followed by fever, headache, reah,
nauses, or vomiting, consult a doctor promptly.

The state of the s

Ask a doctor before use if your child has liver disease

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

When using this product do not exceed recommended dose (see overdose warning)

see overtose wammig)

Stop use and ask a doctor if

pain gets worse or lasts more than 5 days
fever gets worse or lasts more than 3 days
new symptoms occur
redness or swelling is present
These could be signs of a serious condition.

Reep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Quick medical attention is critical even if you do not notice any signs or symptoms.

- Directions

   this product does not contain directions or complete warnings for adult use

   do not give more than directed (see overdose warning)

   shake well before using
- man and the second of the
- into the dosing cup

   repeat dose every 4 hours while symptoms last

#### Drug Facts (continued)

o nor give more and	an o annes in 24 nours	
Weight (lb)	Age (yr)	Dose (mL)*
under 24	under 2 years	ask a doctor
24-35	2-3 years	5 mL
36-47	4-5 years	7.5 mL
48-59	6-8 years	10 mL
60-71	9-10 years	12.5 mL
72-95	11 years	15 mL

\*\*ras directed by a doctor

\*\*the transfer of the transfer of

- Other information
   each 5 ml. contains: sodium 3 mg
   store at 20-25°C (68-77°F)
   do not use if printed neckband is broken or missing

Inactive ingredients anhydrous citric acid, calcium sulfate, carrageeran, flavor, glycerin, hydroxyethyl cellulose, microcrystalline cellulose and carboxymethylcellulose sodium propylene glycol, purified water, sodium benzoate, sorbitol sol sucralose, tribasic sodium phosphate

Questions or comments? 1-888-309-9030

\*\*This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., distributor of Children's Tylenol® Oral Suspension.



Compare to the active ingredient of Children's Tylenol® Oral Suspension

For Ages 2 to 11 Years

# Children's

Acetaminophen 160 mg per 5 mL Pain Reliever/Fever Reducer **Oral Suspension** 

Alcohol free Aspirin free lbuprofen free Paraben Free



8 FL 0Z (237 mL)

DIRECTIONS: STEP 1.





STEP 2. Remove child protective cap



STEP 3. Squeeze your child's dose into the




IMPORTANT:

Please see dosing chart or ask a doctor for your child's correct dose

DISTRIBUTED BY OLD EAST MAIN CO. TN 37072

100% Satisfaction Guaranteed! If you're not satisfied with this product for any reason, please call us so we can make you 100% satisfied.

(888) 309-9030





95934 VT C1

### **DG HEALTH CHILDRENS PAIN AND FEVER**

acetaminophen suspension

#### **Product Information**

**Product Type** 

HUMAN OTC DRUG

Item Code (Source)

NDC:55910-738

**Route of Administration** 

**ORAL** 

Active Ingredient/Active Moiety			
Ingredient Name	<b>Basis of Strength</b>	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	160 mg in 5 mL	

Inactive Ingredients	
Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
CALCIUM SULFATE, UNSPECIFIED FORM (UNII: WATODDB505)	
CARRAGEENAN (UNII: 5C69YCD2YJ)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K6790BS311)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
SODIUM PHOSPHATE, TRIBASIC (UNII: A752Q30A6X)	

Product Characteristics			
Color	WHITE (opaque, viscous)	Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

P	Packaging			
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55910-738- 34	1 in 1 CARTON	05/03/2024	
1		237 mL in 1 BOTTLE; Type 0: Not a Combination Product		
2	NDC:55910-738- 26	1 in 1 CARTON	04/30/2024	
2		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	04/30/2024	

# Labeler - Dolgencorp Inc (068331990)

Revised: 5/2024 Dolgencorp Inc