CHILDRENS MUCINEX CHEST CONGESTION- guaifenes in liquid Reckitt Benckis er LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Childrens Mucinex® Chest Congestion

Drug Facts

Active ingredient (in each 5 mL)

Guaifenesin, USP 100 mg

Purpose

Expectorant

Uses

helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Ask a doctor before use if the child has

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- \blacksquare mL = milliliter

Age	Dose
children 6 wears to under 10 wears	5 mL – 10 mL
children 6 years to under 12 years	every 4 hours
abildren 4 veers to under 6 veers	2.5 mL – 5 mL
children 4 years to under 6 years	every 4 hours
children under 4 years	do not use

Other information

- tamper evident: do not use if printed neckband on bottle cap is broken or missing
- store between 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided

Inactive ingredients

acesulfame potassium, anhydrous citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, maltitol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions?

1-866-MUCINEX (1-866-682-4639)

You may also report side effects to this phone number.

PRINCIPAL DISPLAY PANEL - 118 mL Bottle Carton

NDC 63824-280-64

Children's

Mucinex®

Chest

Congestion

Guaifenesin Oral Solution, USP Expectorant

- Chest Congestion
- ☐ Breaks up Mucus
- Alcohol-free

Grape Flavor Liquid

4 FL OZ (118 mL)

Tamper evident: do not use if printed neckband on bottle cap is broken or missing.

Mucinex:

Chest Congestion

Guaifenesin Oral Solution, USP Expectorant


NDC 63824-280-64

Mucinex Mucinex

Chest **Congestion Congestion**

Guaifenesin Oral Solution, USP Expectorant

- ✓ Chest Congestion
- Breaks up Mucus
- Alcohol-free



Chest



DOSING CUP INCLUDED

Each dose of Children's Mucinex® Chest Congestion Liquid:

- ✓ Relieves Chest Congestion
- ✓ Breaks up Mucus
- Alcohol-free

Questions or comments? 1-866-MUCINEX (1-866-682-4639) 118 mL)






or www.mucinex.com

LOT NO.

EXP.

: 5885P M5 C5





Drug Facts

Active Ingredient (In each 5 mL) Purpose

Guaifenesin, USP 100 mg......Expectorant

Uses helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Ask a doctor before use if the child has

- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with asthma

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not use dosing cup with other products
- mL = milliliter

Mucinex:

Chest Congestion

Also look for

Age	Dose
children 6 years to	5 mL - 10 mL
under 12 years	every 4 hours
children 4 years to	2.5 mL – 5 mL
under 6 years	every 4 hours
children under 4 years	do not use

Other Information


- tamper evident: do not use if printed neckband on bottle cap is broken or missing
- store between 20-25°C (68-77°F)
- do not refrigerate
- dosing cup provided

Inactive Ingredients acesulfame potassium, anhydrous citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, maltitol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sorbitol, sucralose, xanthan gum

Questions? 1-866-MUCINEX (1-866-682-4639)
You may also report side effects to this phone number.



Dist. by: ReckittBenckiser, Parsippany, NJ 07054-0224 @RB 2012



CHILDRENS MUCINEX CHEST CONGESTION

guaifenesin liquid

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63824-280
Route of Administration	ORAL		

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
Guaifenesin (UNII: 495W7451VQ) (Guaifenesin - UNII:495W7451VQ)	Guaifenesin	100 mg in 5 mL	

Inactive Ingredients

Ingredient Name	Strength
acesulfame potassium (UNII: 23OV73Q5G9)	
anhydrous citric acid (UNII: XF417D3PSL)	
FD&C blue no. 1 (UNII: H3R47K3TBD)	
FD&C red no. 40 (UNII: WZB9127XOA)	
glycerin (UNII: PDC6A3C0OX)	
maltitol (UNII: D65DG142WK)	
propylene glycol (UNII: 6DC9Q167V3)	
water (UNII: 059QF0KO0R)	
saccharin sodium (UNII: SB8ZUX40TY)	
sodium benzoate (UNII: OJ245FE5EU)	
sorbitol (UNII: 506T60A25R)	
sucralose (UNII: 96K6UQ3ZD4)	
xanthan gum (UNII: TTV12P4NEE)	

Product Characteristics			
Color	PURPLE	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

F	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:63824-280-64	1 in 1 CARTON			
1		118 mL in 1 BOTTLE			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL	part341	07/01/2011	

Labeler - Reckitt Benckiser LLC (094405024)

Revised: 11/2012 Reckitt Benckiser LLC