HEAD CONGESTION PLUS FLU SEVERE- acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated United Natural Foods, Inc. dba UNFI

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

1174-ELN-2023-0517

Drug Facts

Active ingredients (in each caplet)	Purposes
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 100 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these symptoms due to the common cold:
 - nasal congestion
 - headache
 - minor aches and pains
 - cough
 - sore throat
 - sinus congestion and pressure
- helps loosen phlegm (mucus) and thin bronchial secretions to drain bronchial tubes and make coughs more productive
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. The maximum daily dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert

Acetaminophen may cause severe skin reactions. Symptoms may include:

skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over	 take 2 caplets every 4 hours do not take more than 10 caplets in 24 hours
children under 12 years	■ ask a doctor

Other information

- store between 20-25°C (68-77°F) in a dry place
- retain carton for complete product information and warnings

Inactive ingredients

colloidal silicon dioxide, corn starch, croscarmellose sodium, crospovidone, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions or comments?

1-855-423-2630

PRINCIPAL DISPLAY PANEL

EQUALINE®

NDC 41163-713-02

compare to Sudafed PE® Head Congestion + Flu Severe active ingredients†

head congestion plus flu severe PE

acetaminophen (pain reliever/fever reducer)

dextramethorphan HBr (cough suppressant)

quaifenesin (expectorant)

phenylephrine HCI (nasal decongestant)

for relief of

• sinus pressure • headache

- sore throat cough
- chest congestion

24 caplets

actual size

com starch, croscarmellose sodium, crospovidone, Inactive ingredients colloidal silicon dioxide,

 retain carton for complete product information and warnings Other information store between 20-25°C (68-77°F) in a dry place

Drug Facts (continued)

■ IIVer disease

Questions or comments? 1-822-423-5930

povidone, pregelatinized starch, stearic acid, talc, titanium dioxide microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, FD&C yellow #6 aluminum lake, magnesium stearate, maltodextrin,

notice any signs or symptoms.

keep out of reach of children.

■ redness or swelling is present

Stop use and ask a doctor if

These could be signs of a serious condition.

■ Tever gets worse or lasts more than 3 days

■ uervousness, dizziness, or sleeplessness occur

condh that occurs with too much phlegm (mucus)

■ trouble urinating due to an enlarged prostate gland

bersistent or chronic cough such as occurs with smoking,

serums, chronic bronchius, or emphysema

Drug Facts (continued)

children under µiðµ pjoog bressure ■ neart disease Ask a doctor before use if you have 12 years and over

■ if you have ever had an allergic reaction to this product or any

an MAOI, ask a doctor or pharmacist before taking this product. the MAOI drug. If you do not know if your prescription drug contains conditions, or Parkinson's disease), or for 2 weeks after stopping (MAUI) (certain drugs for depression, psychiatric or emotional ■ if you are now taking a prescription monoamine oxidase inhibitor acetaminophen, ask a doctor or pharmacist.

nonprescription). If you are not sure whether a drug contains with any other drug containing acetaminophen (prescription or

esu fon od vomiting, consult a doctor promptly.

days, is accompanied or followed by fever, headache, rash, nausea, or Sore throat warning: If sore throat is severe, persists for more than 2 If a skin reaction occurs, stop use and seek medical help right away. Allergy slert: Acetaminophen may cause severe skin reactions. ■ 3 or more alcoholic drinks every day while using this product

 with other drugs containing acetaminophen ■ more than 4,000 mg of acetaminophen in 24 hours

Severe liver damage may occur if you take

dose of this product is 10 caplets (3,250 mg acetaminophen) in 24 hours. Liver warning: This product contains acetaminophen. The maximum daily **Marnings**

Drug Facts (continued)

Drug Facts

compare to

Sudafed PE®

active ingredients¹

■ do not take more than 10 caplets in 24 hours ■ take 2 caplets every 4 hours adults and children Directions • do not take more than directed (see overdose warning)

attention is critical for adults as well as for children even if you do not

Poison Control Center right away (1-800-222-1222). Quick medical

If pregnant or breast-feeding, ask a health professional before use

condh comes back or occurs with rash or headache that lasts

Overdose warning: In case of overdose, get medical help or contact a

■ pain, nasal congestion, or cough gets worse or lasts more than 7 days

When using this product do not exceed recommended dosage

Ask a doctor or pharmacist before use if you are taking the blood

uew symptoms occur

†This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Sudated PE® Head Congestion + Flu Severe.

NOT USE IF BLISTER UNITS ARE TORN OR BROKEN

Like it or let us make it right
That's our quality promise.
855-423-2630

DISTRIBUTED I PROVIDENCE, I 프망

F117402ELN_R

41163 49694

Nasal decongestant Рћепујерћгіпе НСІ 5 тд. Expectorant Guaifenesin 100 mg Cough suppressant Dextromethorphan HBr 10 mg. Active ingredients (in each caplet)

 femborarily reduces tever ■ minor aches and pains ■ cough common cold:

ussal congestion

Drug Facts (continued)

drain bronchial tubes and make coughs more productive ■ helps loosen phlegm (mucus) and thin bronchial secretions to sinus congestion and pressure

USES - temporarily relieves these symptoms due to the

Drug Facts (continued)

EQUALINE®

Head Congestion + Flu Severe

NDC 41163-713-02

head congestion plus flu severe PE

acetaminophen (pain reliever/fever reducer) dextromethorphan HBr (cough suppressant) guaifenesin (expectorant) phenylephrine HCI (nasal decongestant)

for relief of

- sinus pressure headache
- sore throat cough
- chest congestion

24 caplets

actual size

HEAD CONGESTION PLUS FLU SEVERE

acetaminophen, dextromethorphan hydrobromide, guaifenesin, and phenylephrine hydrochloride tablet, coated

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:41163-713

Route of Administration ORAL

Active Ingredient/Active Moiety			
Ingredient Name	Basis of Strength	Strength	
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg	
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg	
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg	
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg	

Inactive Ingredients	
Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M280L1HH48)	
CROSPOVIDONE (UNII: 2S7830E561)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
ALUMINUM OXIDE (UNII: LMI26O6933)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
STARCH, PREGELATINIZED CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics			
Color	orange	Score	no score
Shape	OVAL	Size	17mm
Flavor		Imprint Code	AAA;1134
Contains			

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date

12 in 1 BLISTER PACK; Type 0: Not a Combination Product	1	NDC:41163- 713-02	2 in 1 CARTON	08/19/2022	
	1				

Marketing Information			
Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
part341	08/19/2022		
	Application Number or Monograph Citation	Application Number or Monograph Marketing Start Citation Date	

Labeler - United Natural Foods, Inc. dba UNFI (943556183)

Revised: 5/2023 United Natural Foods, Inc. dba UNFI