

SEVERE CONGESTION AND COUGH- dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated
Walgreen Company

Walgreens 44-648-SCC

Active ingredients (in each caplet)

Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purpose

Cough suppressant
Expectorant
Nasal decongestant

Uses

- temporarily relieves:
 - cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
 - the intensity of coughing
 - the impulse to cough to help you get to sleep
 - nasal congestion due to a cold
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- high blood pressure
- heart disease
- difficulty in urination due to enlargement of the prostate gland
- diabetes
- thyroid disease
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough that occurs with too much phlegm (mucus)

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache.

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and over: take 2 caplets every 4 hours
- children under 12 years: do not use

Other information

- **each caplet contains:** sodium 3 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal display panel

NDC 0363-6048-19

Walgreens

WALGREENS PHARMACIST RECOMMENDED†

Compare to active ingredients in
Maximum Strength Mucinex®
Fast-Max® Severe Congestion & Cough††

**Severe
Congestion
& Cough**

DEXTROMETHORPHAN HBr / COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength

Multi-Symptom

- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus

20

COATED
CAPLETS

ACTUAL SIZE

**TAMPER EVIDENT: DO NOT USE IF PACKAGE IS
OPENED OR IF BLISTER UNIT IS TORN, BROKEN
OR SHOWS ANY SIGNS OF TAMPERING**


DISTRIBUTED BY: **WALGREEN CO.**
DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2023 Walgreen Co.

†Our pharmacists recommend the Walgreens brand.

We invite you to compare to national brands.

††This product is not manufactured or distributed by RB Health (US) LLC,
owner of the registered trademark Maximum Strength Mucinex®
FAST-MAX® Severe Congestion & Cough.

50844 ORG042364809



NDC 0363-6048-19

Compare to active ingredients in Maximum Strength Mucinex® Fast-Max® Severe Congestion & Cough™


Severe Congestion & Cough

DEXTROMETHORPHAN HBR / COUGH SUPPRESSANT
GUAIFENESIN / EXPECTORANT
PHENYLEPHRINE HCl / NASAL DECONGESTANT

Maximum Strength **Multi-Symptom**

- Controls cough
- Relieves nasal & chest congestion
- Thins & loosens mucus

20 COATED CAPLETS



ACTUAL SIZE

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

W3ORG0523-F
REV0523

Do not print

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

90844 ORG042364809
FAST-MAX® Severe Congestion & Cough™
owner of the registered trademark Maximum Strength Mucinex®
This product is not manufactured or distributed by BB Health (US) LLC.
We invite you to compare to national brands.
Our pharmacists recommend the Walgreens brand.

KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Drug Facts

Active ingredients (in each caplet)	Purpose
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses ■ temporarily relieves:

- cough due to minor throat and bronchial irritation as may occur with the common cold or inhaled irritants
- the intensity of coughing
- nasal congestion due to a cold
- the impulse to cough to help you get to sleep
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- high blood pressure
- heart disease
- diabetes
- thyroid disease
- difficulty in urination due to enlargement of the prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not get better within 7 days or occur with fever
- cough persists more than 7 days, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than directed

- do not take more than 12 caplets in any 24-hour period
- adults and children 12 years and over: take 2 caplets every 4 hours
- children under 12 years: do not use

Other information ■ each caplet contains: sodium 3 mg

TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN

- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Drug Facts (continued)

Inactive ingredients corn starch, FD&C blue #2 aluminum lake, FD&C red #40 aluminum lake, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, polysorbate 80, polyvinyl alcohol, povidone, silicon dioxide, sodium starch glycolate, stearic acid, talc, titanium dioxide

Questions or comments? 1-800-426-9391

ITEM 766690 W00000-0000-0
DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED
walgreens.com
©2023 Walgreen Co.

Walgreens 44-648

SEVERE CONGESTION AND COUGH

dextromethorphan hbr, guaifenesin, phenylephrine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0363-6048
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII: 7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII: 495W7451VQ)	GUAIFENESIN	200 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
FD&C BLUE NO. 2--ALUMINUM LAKE (UNII: 4AQJ3LG584)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	red (maroon)	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	44;648
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-6048-19	2 in 1 CARTON	07/18/2023	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0363-6048-09	2 in 1 PACKAGE	07/20/2015	07/18/2023
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/20/2015	

Labeler - Walgreen Company (008965063)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(0363-6048) , pack(0363-6048)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0363-6048)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(0363-6048)

Revised: 7/2024

Walgreen Company