

**NUMBING PAIN RELIEF- 5% lidocaine cream**  
**Medhome Pharma Inc**

-----

**Active Ingredient**

Lidocaine 5%

**Purpose**

Local Anesthetic

**Use**

For temporary relief of local discomfort, itching, pain, soreness, or burning related to the tattoo process

**Warnings**

For external use only.

**Do not use**

Do not exceed the recommended daily dosage unless directed by a doctor

**Stop Use**

Do not exceed the recommended daily dosage unless directed by a doctor.

Avoid contact with eyes. If this happens, rinse thoroughly with water.

If symptoms persist for more than 7 days or subside but occur again within 3 days, stop use and consult a physician.

Stop use and seek medical attention if signs of an allergy occur, such as redness, irritation, swelling, pain, or other severe symptoms. If symptoms do not subside completely within 7 days, seek medical help.

**Ask Doctor**

Stop use and seek medical attention if signs of an allergy occur, such as redness, irritation, swelling, pain, or other severe symptoms. If symptoms do not subside completely within 7 days, seek medical help.

**Directions**

Before applying the numbing cream, thoroughly wash your hands and wear disposable

tattoo gloves.

Clean the area of the body receiving the tattoo, Dry completely.

Apply a generous amount of numbing cream to the cleaned area and rub it in thoroughly in a circular motion for 30 to 60 seconds, Then. cover with plastic wrap.

Leave the cream and wrap in place for 30 to 45 minutes, Wipe away the cream and thoroughly clean the area to prepare for a pain-free tattoo.

### **Keep Out Of Reach Of Children**

Keep out of reach of children

### **Other information**

Store between 15-30°C (59-86°F). Avoid freezing and excessive heat above 40°C (104°F).

### **Inactive ingredients**

**Water, Carbomer 934, propylene Glycol, soybean Lecithin, .Alpha.-Tocopherol Acetate, Benzyl Alcohol.**

### **Package**



## NUMBING PAIN RELIEF

5% lidocaine cream

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:84007-334
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>LIDOCAINE</b> (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	5 g in 100 g

### Inactive Ingredients

Ingredient Name	Strength
<b>ALPHA-TOCOPHEROL ACETATE</b> (UNII: 9E8X80D2L0)	
<b>BENZYL ALCOHOL</b> (UNII: LKG8494WBH)	
<b>CARBOMER 934</b> (UNII: Z135WT9208)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SOYBEAN LECITHIN</b> (UNII: 1DI56QDM62)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84007-334-01	10 g in 1 BOTTLE; Type 0: Not a Combination Product	07/24/2024	
2	NDC:84007-334-02	20 g in 1 BOTTLE; Type 0: Not a Combination Product	07/24/2024	
3	NDC:84007-334-03	30 g in 1 BOTTLE; Type 0: Not a Combination Product	07/24/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	07/24/2024	

**Labeler** - Medhome Pharma Inc (129936136)

**Registrant** - Medhome Pharma Inc (129936136)

Revised: 7/2024

Medhome Pharma Inc