

WELMATE FAMOTIDINE 10MG- famotidine 10mg tablet
YYBA Corp

Welmate Famotidine 10mg 73581-108-10

Active ingredient (in each tablet)

Famotidine, USP 10 mg

Purpose

Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert

Do not use if you are allergic to famotidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

If pregnant or breast-feeding

ask a health professional before use.

Directions

- adults and children 12 years and over:
 - to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
 - to prevent symptoms, swallow 1 tablet (of 10 mg) with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn
 - to prevent symptoms, swallow 1 tablet (of 20 mg) with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
 - before eating food or drinking beverages that cause heartburn
 - do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Inactive Ingredients

Microcrystalline Cellulose, Pre-gelatinized Starch, Sodium Starch Glycolate, Hydroxypropyl Cellulose, Magnesium Stearate, Hypromellose, Titanium Dioxide, Triacetin, Talc, Macrogol

Other information

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°C to 25°C (68°F to 77°F)
- protect from moisture

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL-300's Count

DRUG FACTS, DIRECTIONS + DOSAGE INFORMATION

Drug Facts	
Active ingredient (in each tablet) Famotidine USP 10 mg.....	Purpose Acid reducer
Uses ■ relieves heartburn associated with acid indigestion and sour stomach ■ prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages	
Warnings Allergy alert: Do not use if you are allergic to famotidine or other acid reducers Do not use ■ if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. ■ with other acid reducers	
Ask a doctor before use if you have ■ had heartburn over 3 months. This may be a sign of a more serious condition. ■ heartburn with lightheadedness, sweating, or dizziness ■ chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness ■ frequent chest pain ■ frequent wheezing, particularly with heartburn ■ unexplained weight loss ■ nausea or vomiting ■ stomach pain ■ kidney disease	
Drug Facts, cont.	
Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.	
Stop use and ask a doctor if ■ your heartburn continues or worsens ■ you need to take this product for more than 14 days	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)	
Directions	
adults and children 12 years and over:	■ to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew. ■ to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn ■ do not use more than 2 tablets in 24 hours
children under 12 years	ask a doctor

DIRECTIONS FOR USE
To relieve symptoms, swallow 1 tablet with water.
To prevent symptoms, swallow 1 tablet with water 15-60 minutes before eating or drinking.
Do not take more than 2 tablets in 24 hours.



PACKAGE LABEL.PRINCIPAL DISPLAY PANEL-90's Count



ACID REDUCER

ORIGINAL STRENGTH

Famotidine USP 10 mg

Prevents & Relieves Heartburn due to Acid Indigestion

90 TABLETS

100% SATISFACTION GUARANTEED

FAST ACTING JUST 1 TABLET



DO NOT USE IF PUNCHED SEAL UNDER CAP IS MISSING OR DAMAGED

Drug Facts

Purpose
Active ingredient
(in each tablet)
Famotidine, USP, 10 mg.....Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to famotidine or other acid reducers.

Do not use: if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. with other acid reducers

Ask a doctor before use if you have had heartburn over 3 months. This may be a sign of a more serious condition heartburn with **lightheadedness, sweating, dizziness** chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness frequent chest pain frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain kidney disease

Ask a doctor or pharmacist before use if you are taking a prescription drug. Acid reducers may interact with certain prescription drugs.



PEEL FOR DIRECTIONS 3 73581-00076 0

LIFT HERE

Drug Facts (continued)

Stop and ask a doctor if your heartburn continues or worsens you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours
- children under 12 years: ask a doctor

Other information

- read the directions and warnings before use
- store at 20°-25°C (68°-77°F)
- protect from moisture

Inactive ingredients

Hydroxypropyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide, triacetin.

Questions?

call toll-free 1-866-933-4337



WELMATE FAMOTIDINE 10MG

famotidine 10mg tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73581-108
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	10 mg

Inactive Ingredients

Ingredient Name	Strength
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ05DW1A)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
TALC (UNII: 7SEV7J4R1U)	
STARCH, CORN (UNII: O8232NY3SJ)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	

Product Characteristics

Color	white (off-white)	Score	no score
Shape	ROUND (debossed with "V" on one side and "21" on the other side)	Size	5mm
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73581-108-10	300 in 1 BOTTLE; Type 0: Not a Combination Product	07/25/2024	
2	NDC:73581-108-09	90 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215822	07/25/2024	

Labeler - YYBA Corp (006339772)

Registrant - YYBA Corp (006339772)