

**BC MAX- aspirin, acetaminophen, caffeine powder
Medtech Products Inc.**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

BC Max 63029-203-16

BC™

MAX STRENGTH

ACETAMINOPHEN & ASPIRIN (NSAID) / CAFFEINE

PAIN RELIEVER / PAIN RELIEVER AID

16 PACKS

LEMONADE FLAVOR

DRUG FACTS

Attention: Contains acetaminophen. Read entire label before use.

Drug Facts

Active ingredients (in each powder):

Purposes

Aspirin (NSAID*) 500 mg.....Pain reliever
Acetaminophen 500mg.....Pain reliever
Caffeine 65 mg.....Pain reliever aid

*nonsteroidal anti-inflammatory drug

Uses

Temporarily relieves minor aches and pains due to:

- headache
- backache
- muscle aches
- toothache
- menstrual cramps
- colds
- minor arthritis pain

Warnings:

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause severe allergic reaction which may include:

- hives • facial swelling • shock • asthma (wheezing).

Acetaminophen may cause severe skin reactions. Symptoms may include

- skin reddening • blisters • rash

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher

if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed.

Caffeine warning:

This product contains caffeine. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

Do not use

- if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

taking

- a prescription drug for diabetes, gout or arthritis
- the blood thinning drug warfarin

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away
- you experience any of the following signs of stomach bleeding
- feel faint • vomit blood • have bloody or black stool

- have stomach pain that does not get better
- redness or swelling is present
- any new symptoms appear
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **Do not take more than directed** (see overdose warning)
- **Adults and children 12 years and over:** place 1 powder on tongue every 6 hours, while symptoms persist. Drink a full glass of water or other liquid.
- Do not take more than 4 powders in 24 hours unless directed by a doctor.
- **Children under 12 years:** ask a doctor.

Other information

Store below 25°C (77°F)

Inactive ingredients

flavor, guar gum, mannitol, microcrystalline cellulose, silicon dioxide, sucralose, xylitol

Questions?

1-866-255-5197 BCpowder.com

TAMPER EVIDENT: DO NOT USE IF PACK IS DAMAGED OR OPEN.

TO OPEN CHILD RESISTENT PACKAGE

1. **SHAKE PACK** so powder settles on bottom
2. Pinch and tear at either arrow or use scissors

PRINCIPAL DISPLAY PANEL

BC

MAX STRENGTH

ACETAMINOPHEN & ASPIRIN (NSAID) / CAFFEINE

PAIN RELIEVER / PAIN RELIEVER AID

16 PACKS

LEMONADE FLAVOR

NEW! WITH
ACETAMINOPHEN



MAX STRENGTH

FAST PAIN RELIEF POWDER

WITH **ACETAMINOPHEN & ASPIRYN (NSAID)**
PAIN RELIEVER, CAFFEINE, PAIN RELIEVER AID.



16 PACKS

LEMONADE
FLAVOR

Drug Facts (continued)

(prescription or nonprescription). If you are not sure whether a drug contains acetaminophen ask a doctor or pharmacist.

Ask a doctor before use if

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking

- a prescription drug for diabetes, gout or arthritis
- the blood thinning drug warfarin

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- redness or swelling is present
- any new symptoms appear
- ringing in the ears or a loss of hearing occurs

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- **Do not take more than directed** (see overdose warning).
- **Adults and children 12 years and over:** place 1 powder on tongue every 6 hours, while symptoms persist. Drink a full glass of water or other liquid. ■ **Do not take more than 4 powders in 24 hours unless directed by a doctor.**
- **Children under 12 years:** ask a doctor.

Other information store below 25°C (77°F)

Inactive ingredients flavor, guar gum, mannitol, microcrystalline cellulose, silicon dioxide, sucralose, xylitol

Questions? 1-866-255-5197 BCpowder.com

NO COAT
NO COPY

NO COAT
NO COPY

TAMPER EVIDENT: DO NOT USE IF PACK IS DAMAGED OR OPEN.

ATTENTION: CONTAINS ACETAMINOPHEN. READ ENTIRE LABEL BEFORE USE.

TO OPEN CHILD-RESISTANT PACKAGE



1. SHAKE PACK so powder settles on bottom.



2. Pinch and tear at either arrow or use scissors.

Drug Facts

Active ingredients (in each powder)	Purposes
Acetaminophen 500 mg	Pain reliever
Aspirin (NSAID*) 500 mg	Pain reliever
Caffeine 65 mg	Pain reliever aid
*nonsteroidal anti-inflammatory drug	

Uses

Temporarily relieves minor aches and pains due to:

- headache
- backache
- muscle ache
- toothache
- menstrual cramps
- colds
- minor arthritis pain

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause severe allergic reaction which may include: ■ hives ■ facial swelling ■ shock ■ asthma (wheezing). Acetaminophen may cause severe skin reactions. Symptoms may include ■ skin reddening ■ blisters ■ rash

Drug Facts (continued)

If a skin reaction occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you ■ are age 60 or older ■ have had stomach ulcers or bleeding problems ■ take a blood thinning (anticoagulant) or steroid drug ■ take other drugs containing prescription or non-prescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ have 3 or more alcoholic drinks every day while using this product ■ take more or for a longer time than directed

Caffeine warning: This product contains caffeine. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

Do not use ■ if you have ever had an allergic reaction to aspirin or any other pain reliever/fever reducer ■ with any other drug containing acetaminophen

BC MAX

aspirin, acetaminophen, caffeine powder

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63029-203
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	500 mg
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)		CAFFEINE	65 mg	
Inactive Ingredients				
Ingredient Name		Strength		
GUAR GUM (UNII: E89I1637KE)				
MANNITOL (UNII: 3OWL53L36A)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
XYLITOL (UNII: VCQ006KQ1E)				
Product Characteristics				
Color	WHITE	Score		
Shape		Size		
Flavor	LEMON	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63029-203-04	4 in 1 CARTON; Type 0: Not a Combination Product	10/01/2020	
2	NDC:63029-203-16	16 in 1 CARTON; Type 0: Not a Combination Product	03/15/2020	
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC MONOGRAPH NOT FINAL	part343	03/15/2020		

Labeler - Medtech Products Inc. (122715688)

Revised: 4/2020

Medtech Products Inc.