

NAPROXEN- naproxen sodium tablet, film coated
Amerisource Bergen

Good Neighbor Pharmacy 44-604

Active ingredient (in each caplet)

Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)*

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

- temporarily relieves minor aches and pains due to:
 - headache
 - muscular aches
 - the common cold
 - toothache
 - menstrual cramps
 - backache
 - minor pain of arthritis
- temporarily reduces fever

Warnings

Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- asthma (wheezing)
- skin reddening
- shock
- blisters
- rash
- facial swelling
- hives

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]

- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- the stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you are taking a diuretic
- you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
- you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- you have symptoms of heart problems or stroke:
 - leg swelling
 - chest pain
 - slurred speech
 - trouble breathing
 - weakness in one part or side of body
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- you have difficulty swallowing
- it feels like the pill is stuck in your throat
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding,

ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- **the smallest effective dose should be used**
- drink a full glass of water with each dose
- adults and children 12 years and older
 - take 1 caplet every 8 to 12 hours while symptoms last
 - for the first dose you may take 2 caplets within the first hour
 - do not exceed 2 caplets in any 8- to 12-hour period
 - do not exceed 3 caplets in a 24-hour period
- children under 12 years: ask a doctor

Other information

- **each caplet contains:** sodium 20 mg
- store at 20°-25°C (68°-77°F). Avoid high humidity and excessive heat above 40°C (104°F)
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, FD&C blue #2 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, talc, titanium dioxide

Questions or comments?

Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Principal display panel

**GOOD
NEIGHBOR
PHARMACY®**

Compare to ALEVE® active ingredient†

NDC 46122-564-71

Naproxen

naproxen sodium tablets 220 mg **(NSAID)**

Pain Reliever/Fever Reducer

Strength to last 12 hours

50 Caplets

220 mg Each

Actual Size

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED
SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Aleve®.

50844 REV1221B60415

Distributed By

AmerisourceBergen

1 West First Avenue

Conshohocken, PA 19428

Questions or Concerns?

www.mygnp.com

GOOD NEIGHBOR PHARMACY

SATISFACTION

GUARANTEED

BRAND PRODUCTS

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions ■ do not take more than directed
 ■ the smallest effective dose should be used
 ■ drink a full glass of water with each dose
 ■ adults and children 12 years and older
 ■ take 1 caplet every 8 to 12 hours while symptoms last
 ■ for the first dose you may take 2 caplets within the first hour
 ■ do not exceed 2 caplets in any 8- to 12-hour period
 ■ do not exceed 3 caplets in a 24-hour period
 ■ children under 12 years: ask a doctor

Other information
 ■ each caplet contains sodium 20 mg
 ■ store at 20-25°C (68-77°F). Avoid high humidity and excessive heat above 40°C (104°F)
 ■ see end flap for expiration date and lot number

Inactive ingredients: croscarmellose sodium, FD&C blue #2 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, talc, titanium dioxide

Questions or comments?
 Call 1-800-426-9391 8:30 AM-4:00 PM ET, Monday-Friday

Compare to ALEVE® active ingredient*

NDC 46122-564-71

GOOD NEIGHBOR PHARMACY®

Naproxen
 naproxen sodium tablets 220 mg (NSAID)

Pain Reliever/Fever Reducer

Strength to last 12 hours

50 Caplets | 220 mg Each

Actual Size

This product is not manufactured or distributed by Bayer HealthCare LLC, owner of the registered trademark Aleve®. 50844 REV12/1800415

SATISFACTION GUARANTEED

Distributed By
 AmerisourceBergen
 1 West First Avenue
 Conshohocken, PA 19428
www.mygnp.com

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

ABC#: 10191501

0 87701 43161 6

No Print / No Varnish
 Lot no. & Exp. date

Drug Facts **KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION**

Active ingredient (in each caplet) Purpose
 Naproxen sodium 220 mg (naproxen 200 mg) (NSAID)* Pain reliever/fever reducer
 *nonsteroidal anti-inflammatory drug

Uses
 ■ temporarily relieves minor aches and pains due to:
 ■ headache ■ muscle aches ■ the common cold
 ■ toothache ■ menstrual cramps
 ■ backache ■ minor pain of arthritis
 ■ temporarily reduces fever

Warnings
Allergy alert: Naproxen sodium may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ asthma (wheezing) ■ skin redness ■ shock
 ■ hives ■ rash ■ facial swelling
 ■ blisters ■ If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause serious stomach bleeding. The chance is higher if you ■ are age 60 or older
 ■ have had stomach ulcers or bleeding problems
 ■ take a blood thinning (anticoagulant) or steroid drug
 ■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) ■ take more or for a longer time than directed
 ■ have 3 or more alcoholic drinks every day while using this product
Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

B-0045-604-15
 REV12/1800415

Drug Facts (continued)

Do not use ■ if you have ever had an allergic reaction to any other pain reliever/fever reducer
 ■ right before or after heart surgery

Ask a doctor before use if
 ■ the stomach bleeding warning applies to you
 ■ you have a history of stomach problems, such as heartburn ■ you are taking a diuretic
 ■ you have high blood pressure, heart disease, liver cirrhosis, kidney disease, asthma, or had a stroke
 ■ you have problems or serious side effects from taking pain relievers or fever reducers

Ask a doctor or pharmacist before use if you are
 ■ under a doctor's care for any serious condition
 ■ taking aspirin for heart attack or stroke, because naproxen may decrease this benefit of aspirin
 ■ taking any other drug

When using this product
 ■ take with food or milk if stomach upset occurs
 ■ Stop use and ask a doctor if ■ you experience any of the following signs of stomach bleeding:
 ■ feel faint ■ vomit blood
 ■ have bloody or black stools
 ■ have stomach pain that does not get better
 ■ you have symptoms of heart problems or stroke:
 ■ leg swelling ■ chest pain
 ■ slurred speech ■ trouble breathing
 ■ weakness in one part or side of body
 ■ pain gets worse or lasts more than 10 days
 ■ fever gets worse or lasts more than 3 days
 ■ you have difficulty swallowing
 ■ it feels like the pill is stuck in your throat
 ■ redness or swelling is present in the painful area
 ■ any new symptoms appear

GNP 44-604

NAPROXEN			
naproxen sodium tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46122-564
Route of Administration	ORAL		
Active Ingredient/Active Moiety			

Ingredient Name		Basis of Strength	Strength	
NAPROXEN SODIUM (UNII: 9TN87S3A3C) (NAPROXEN - UNII:57Y76R9ATQ)		NAPROXEN SODIUM	220 mg	
Inactive Ingredients				
Ingredient Name			Strength	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)				
FD&C BLUE NO. 2 ALUMINUM LAKE (UNII: 4AQJ3LG584)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)				
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
TALC (UNII: 7SEV7J4R1U)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
Product Characteristics				
Color	blue	Score	no score	
Shape	OVAL	Size	12mm	
Flavor		Imprint Code	44;604	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46122-564-78	1 in 1 CARTON	02/01/2019	
1		100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
2	NDC:46122-564-81	200 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	02/01/2019	
3	NDC:46122-564-71	1 in 1 CARTON	02/01/2019	
3		50 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
ANDA	ANDA204872		02/01/2019	

Labeler - Amerisource Bergen (007914906)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(46122-564)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(46122-564) , pack(46122-564)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(46122-564)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(46122-564)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(46122-564)