LIDOCAINE PAIN RELIEVE GEL-PATCH- pain relieve gel-patch patch Henan Kangdi Medical Devices Co., Ltd

69561-005-01

Active Ingredient

LIDOCAINE 4%

Purpose

Topical Analgesic

Use

For temporary relief of pain

Warnings

For external use only

Do not use

- more than 1 patch at a time
- on wounds or damaged skin
- with a heating pad
- if you are allergic to any ingredients of this product

When Using

- use only as directed
- avoid contact with the eyes, mucous membranes, or rashes
- do not bandage tightly

Stop Use

- localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering
- condition worsen
- symptoms persist far more than 7 days
- symptoms clear up and occur again within a few days

Ask Doctor

Keep Out Of Reach Of Children

If swallowed, get medical help or contact a Poison Control Centerright away.

Directions

Adult and children 12 years of age and over:

- clean and dry affected area
- remove film from patch and apply to the skin (see illustration)
- apply 1 patch at a time to affected area, not more than 3 to 4 times daily
- remove patch from the skin after at most 8-hour application
- Children under 12 years of age: consult a doctor

Other information

- Avoid storing product in direct sunlight
- Protect product from excessive moisture

Inactive ingredients

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Hydroxyacetophenone, Kaolin, L(+)-Tartaric Acid, Mineral oil. Polysorbate 80, Polyacrylic Acid, Polyvinylpyrrolidone K90, Propylene Glycol, Sodium Polyacrylate, Titanium Dioxide, Water.

Questions

Healthpatchofficial@gmail.com

PRINCIPAL DISPLAY PANEL



No Mess | Easy to Apply and Remove

Temporary relief of minor aches & pains

- Back
- Neck
- Shoulders
- Knee & Elbows

Lidocaine 4% 10 PATCHES 5.5"X 3.94" (10cm×14cm) EACH INDIVIDUALLY PACKAGED





HOW TO APPLY







8H

Active ingredient

Drug Facts

Purpose . Topical anesthetic

Uses For temporary relief of pain

Warnings (For external use only)

Do not use ■ more than I patch at a time ■ on wounds or damaged skin ■ with a heating pad ■ if you are allergic to any ingredients of this product

When using this product use only as directed avoid contact with the eyes, mucous membranes or rashes do not bandage tightly

Stop use and ask a doctor if

■localized skin reactions occur, such as rash, itching, redness, irritation, pain, sw
and blistering _condition worsen _symptoms persist far more than 7 days
_symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison
Control Centerright away.

Adult and children 12 years of age and over:

• clean and dry affected area
• remove film from patch and apply to the skin (see illustration)
• apply 1 patch at a time to affected area, not more than 3 to 4 times daily
• remove patch from the skin after at most 8-hour application
Children under 12 years of age; consult a doctor

Other information

Avoid storing product in direct sunlight Protect product from excessive moisture

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Hydroxyacetophenone, Kaolin, L(+)-Tartaric Acid, Mineral oil. Polysorbate 80, Polyacrylic Acid, Polyvinylpyrrolid one K90, Propylene Glycol, Sodium Polyacrylate, Titanium Dioxide, Water.

QUESTIONS or COMMENTS Healthpatchofficial@gmail.com

Manufacturer by Henan Kangdi Medical Devices Co.,Ltd No.4, 2nd Area, SME Pioneer Park, Industry Cluster Zone, Huaiyang County, Zhoukou, Henan

Made in China

MAXIMUM STRENGTH LIDOCAINE

MAXIMUM STRENGTH LIDOCAINE

LIDOCAINE PAIN RELIEVE GEL-PATCH IDOCAINE PAIN RELIEVE GEL-PATCH

No Mess | Easy to Apply and Remove Temporary relief of minor

aches & pains

Back

LIDOCAINE 4% 18LARGE PATCHES

- Neck
- Shoulders • Knee & Elbows

18 PATCHES 5.5"X 3.94" (10cm×14cm) EACH INDIVIDUALLY PACKAGED



HOW TO APPLY









Uses For temporary relief of pain

Warnings (For external use only)

Do not use ■ more than 1 patch at a time ■ on wounds or damaged skin ■ with a heating pad ■ if you are allergic to any ingredients of this product

When using this product use only as directed avoid contact with the eyes, mucous membranes or rashes do not bandage tightly

Stop use and ask a doctor if

| localized skin reactions occur, such as rash, itching, redness, irritation, pain, swelling and blistering _condition worsen _ symptoms persist far more than 7 days _ symptoms clear up and occur again within a few days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. If swallowed, get medical help or contact a Poison
Control Centerright away.

- Directions
 Adult and children 12 years of age and over:

 clean and dry affected area

 clean and dry affected area

 remove film from patch and apply to the skin (see illustration)

 apply 1 patch at a time to affected area, not more than 3 to 4 times daily

 remove patch from the skin after at most 8-hour application

 Children under 12 years of age: consult a doctor

Other information

#Avoid storing product in direct sunlight # Protect product from excessive moisture

Inactive ingredients

Dihydroxyaluminum Aminoacetate, Edetate Disodium, Glycerin, Hydroxyacetophenone,

Kadolin, Lei-Partari Acid, Mineral Di Polysorbate 80, Polyacrylic Acid, Polyvinylpyrrolid


one K9D, Propylene Glycol, Sodium Polyacrylate, Titanium Dioxide, Water.

QUESTIONS or COMMENTS Healthpatchofficial@gmail.com

Manufacturer by
Henan Kangdi Medical Devices Co.,Ltd
No.4, 2nd Area, SME Pioneer Park, Industry
Cluster Zone, Huaiyang County, Zhoukou, Henan

Made in China

MAXIMUM STRENGTH LIDOCAINE



LIDOCAINE PAIN RELIEVE GEL-PATCH

pain relieve gel-patch patch

Duas	l _	Inform	
Proc	W		lalion

Product Type HUMAN OTC DRUG Item Code (Source) NDC:69561-005

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength

LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987) LIDOCAINE 4 g in 100 g

Inactive Ingredients

Ingredient Name Strength

SODIUM POLYACRYLATE (2500000 MW) (UNII: 05|15|N|2|)

POLYACRYLIC ACID (250000 MW) (UNII: 9G2MAD7J6W)	
GLYCERIN (UNII: PDC6A3C0OX)	
TARTARIC ACID (UNII: W4888I119H)	
POVIDONE K90 (UNII: RDH86HJV5Z)	
WATER (UNII: 059QF0KO0R)	
KAOLIN (UNII: 24H4NWX5CO)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
MINERAL OIL (UNII: T5L8T28FGP)	
POLYSORBATE 80 (UNII: 60ZP39ZG8H)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
DIHYDROXYALUMINUM AMINOACETATE (UNII: DO250MG0W6)	
HYDROXYACETOPHENONE (UNII: G1L3HT4CMH)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69561-005- 01	10 in 1 BOX	07/27/2024	
1		12 g in 1 PATCH; Type 0: Not a Combination Product		
2	NDC:69561-005- 02	18 in 1 BOX	07/27/2024	
2		12 g in 1 PATCH; Type 0: Not a Combination Product		
3	NDC:69561-005- 03	30 in 1 BOX	07/27/2024	
3		12 g in 1 PATCH; Type 0: Not a Combination Product		

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	07/27/2024	

Labeler - Henan Kangdi Medical Devices Co., Ltd (421288241)

Establishment			
Name	Address	ID/FEI	Business Operations
Henan Kangdi Medical Devices Co., Ltd		421288241	manufacture(69561-005)