

**DYNAREX ANTIFUNGAL POWDER- miconazole powder**  
**Dynarex Corporation**

-----  
**1236 Dynarex Antifungal Powder NDC 67777-316-01**

**Active Ingredient**

Active Ingredient	Purpose
Miconazol 2%	Antifungal

**Purpose**

Uses

For external treatment of fungal infections such as athlete's foot, jock itch, sweat rash, infected diaper rash and fungal infections affecting the skin folds e.g. armpits, groin, or under the breasts.

**Warnings**

**For external use only.**

Consult a doctor or pharmacist before using this product if you:

■ Are sensitive to the listed ingredients or any similar medications ■ Are pregnant, planning a pregnancy or breastfeeding

**When using this product:**

You may occasionally experience some side effects. These are rare and consist of local skin irritation or rashes, which occur if you are unusually sensitive to the ingredients listed.

**Do Not Use**

Do not use:

■ if you are sensitive to any of the active or inactive ingredients listed ■ if the infection is on your scalp or nails.

**Cautions**

**Please read this label carefully before use.**

- Dynarex Antifungal Powder is not suitable for the treatment of fungal infections of the scalp or nails, and you should talk to your doctor or pharmacist for alternative treatments for these conditions. In order to ensure successful treatment, it is important to use the powder regularly and to continue for at least 10 days after the disappearance of symptoms. This prevents the infection from reoccurring.

- Remember that most fungal infections are very infectious and can easily be passed on to other family members. In order to prevent this happening, it is important to ensure that anyone who has an infection avoids sharing clothes, towels, or shoes with other family members. After applying the powder, wash and dry your hands thoroughly. If your skin condition does not improve after one week's use, please consult your doctor or pharmacist.
- If you forget to apply the powder, do not apply the missed dose, but apply the next dose as usual and continue as normal. Do not apply two doses at the same time.

## **Interactions**

If you are taking oral anticoagulants (drugs used to thin the blood, such as Warfarin), talk to your doctor or pharmacist before use.

## **Overdose**

Excessive use can result in skin irritation, which usually disappears after discontinuation of therapy. In case of accidental ingestion of powder go to the hospital immediately.

## **Directions**

Sprinkle onto the affected area twice daily. It can be safely applied to broken skin and may also be sprinkled onto clothes and footwear, which come into contact with the infected area.

## **Other information**

Keep the medicine in the original packaging in a dry place, at a temperature not exceeding 30°C/86°F. Protect from light. Do not use after the expiry date shown on the pack.

## **Keep Out Of Reach Of Children**

## **Keep Out Of Reach Of Children**

Do not let the powder get into your eyes. This product contains talc. Do not breathe in the powder as this may cause irritation of the airways. This is especially important for children and infants. If swallowed, contact a poison control center immediately.

## **Inactive ingredients**

Colloidal silicon dioxide, talc.

## **Principal Display panel**

Dynarex Antifungal Powder 1236

AntiPowGP.jpg

**Drug Facts:**

Active Ingredient	Purpose
Miconazole, 2%	Antifungal

**Uses** For external treatment of fungal infections such as athlete's foot, jock itch, sweat rash, infected nappy rash and fungal infections affecting the skin folds e.g. armpits, groin, or under the breasts.

**Warnings**  
For external use only  
Consult a doctor or pharmacist before using this product if you  
 ■ Are you sensitive to the active ingredient or any similar medicine ■ Are you pregnant, planning a pregnancy or breast-feeding

**Do not use**  
 ■ if you are sensitive to any of the active or inactive ingredients listed ■ if the infection is on your scalp or nails.

**When using this product**  
You may occasionally experience some side effects. These are rare and comprise of local skin irritation or rashes, which occur if you are unusually sensitive to the active ingredient

**Keep out of reach of children**  
Do not let the powder get into your eyes. This product contains talc. Do not breathe in the powder as this may cause irritation of the airways. This is especially important for children and infants. If swallowed, contact a poison control center immediately

**Directions** Sprinkle onto the affected area twice daily. It can be safely applied to broken skin and may also be sprinkled onto clothes and footwear, which come into contact with the infected area.

**Other Information** Keep the medicine in the original packaging in a dry place, at a temperature not exceeding 30°C/86°F. Protect from light. Do not use after the expiry date shown on the pack.

**Inactive Ingredients** Colloidal Silicon Dioxide and Talc.

**Reorder No. 1236**

NDC # 67777-316-01



**Anti-Fungal Powder**

Manufactured for:  
Dynarex Corporation  
Orangeburg, NY 10962  
www.dynarex.com  
Made in India

Net Wt. 3 oz. (85 g)

**Caution**  
Please read this label carefully before use.

- Dynarex Anti-Fungal Powder is not suitable for the treatment of fungal infections of the scalp or nails, and you should talk to your doctor or pharmacist for alternative treatments for these conditions. In order to ensure successful treatment, it is important to use the powder regularly and to continue for at least 10 days after the disappearance of symptoms. This prevents the infection from re-occurring.
- Remember that most fungal infections are very infectious and can easily be passed on to other family members. In order to prevent this happening, it is important to ensure that anyone who has an infection avoids sharing clothes, towels, or shoes with other family members. After applying the powder, wash and dry your hands thoroughly. If your skin condition does not improve after one week's use, please consult your doctor or pharmacist.
- If you forget to apply the powder, do not apply the missed dose, but apply the next dose as usual and continue as normal. Do not apply two doses at the same time.

**Interactions**  
If you are taking oral anticoagulants (drugs used to thin the blood, such as Warfarin), talk to your doctor or pharmacist before use.

**Overdose**  
Excessive use can result in skin irritation, which usually disappears after discontinuation of therapy. In case of accidental ingestion of powder go an emergency room immediately.

Lot No. \_\_\_\_\_ Exp. \_\_\_\_\_



6 16784 12361 5

6070850XXX

## DYNAREX ANTIFUNGAL POWDER

miconazole powder

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:67777-316
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>MICONAZOLE NITRATE</b> (UNII: VW4H1CYW1K) (MICONAZOLE - UNII: 7NNO0D7S5M)	MICONAZOLE NITRATE	20 mg in 1 g

### Inactive Ingredients

Ingredient Name	Strength
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	

### Packaging

#	Item Code	Package Description	Marketing Start	Marketing End
---	-----------	---------------------	-----------------	---------------

#	Item Code	Package Description	Date	Date
1	NDC:67777-316-02	24 in 1 CASE	05/19/2016	
1	NDC:67777-316-01	85 g in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug		M005	05/19/2016	

**Labeler** - Dynarex Corporation (008124539)

**Registrant** - Dynarex Corporation (008124539)