# MIDWAY SELECT TOPICAL ANESTHETIC- benzocaine gel Midway Dental Supply, LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

#### **Active Ingredients**

Benzocaine 200 mg ( in each g)

#### **Purpose**

Oral Anesthetic

#### Use

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due to minor dental procedures.

#### Warnings

**Methemoglobinemia warning:** Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in the blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy alert:** Do not use on patients with a history of allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" anesthetics.

#### Do not use

- for more than 7 days unless directed by a physician. If sore mouth symptoms do not improve in 7 days; irritation, pain, or redness persists or worsens; or if swelling, rash or fever develops, see your physician promptly.
- for teething
- in children under 2 years of age

When using this product avoid contact with eyes. If contact occurs, flush with water.

**Do not exceed recommended dosage.** If more than used for pain is accidentally swallowed, get medical help or contact a Poison Control Center right away.

If pregnant or breastfeeding, ask a health professional before use.

Keep out of reach of children.

#### **Directions**

- Apply only amount needed to the oral mucosa to prevent or relieve pain.
- children under 2 years of age: do not use

#### Other Information

Store at 59°-86°F (15°-30°C). Protect from freezing.

#### **Inactive Ingredients**

flavoring, PEG 3350, PEG 400, sodium saccharin. May contain blue #1, green #3, green #5, red #3, red #28, red #40, yellow #5, (tartrazine), yellow #6, as a color additive.

#### **Questions or comments?**

800.474.6111

midwaydental.com

#### Distributed By:

Midway Dental Supply, LLC 32553 Schoolcraft Road Livonia, MI 48150



#### TOPICAL ANESTHETIC GEL

BENZOCAINE 20%



99095811 Rev 06/2018

GLUTEN FREE 1 OZ (30ML)

mouth symptoms do not improve if swelling, rash or fever develops, redness persists or worsens; or directed by a physician. If sore in 7 days; irritation, pain, or

see your physician promptly for teething

in children under 2 years of age Do not exceed recommended contact occurs, flush with water. dosage. If more than used for When using this product Avoid contact with eyes. If

If pregnant or breast feeding, Poison Control Center right away. pain is accidentally swallowed get medical help or contact a ask a physician before use.

Keep out of reach of children. Directions

 Apply only amount needed to the oral mucosa to prevent or relieve pain.

children under 2 years of age: do not use

Store at 59° - 86°F (15 - 30°C). Protect from freezing.

Other information

blue #1, green #3, green #5, red (tartrazine), yellow #6, as a colo #3, red #28, red #40, yellow #5, sodium saccharin. May contain flavoring, PEG 3350, PEG 400, Inactive ingredients

Questions or comments?

800.474.6111

midwaydental.com

# additive.

Active ingredients **Oruq Facts** 

Benzocaine 200 mg *(in each g)* Purpose

Oral Anesthetic

For oral mucosal use only, as directed by dentist. For the temporary relief of pain due

to minor dental procedures.

warning: Use of this product Methemoglobinemia **Marnings** 

may cause methemoglobinemia a serious condition that must be attention if you or a child in your occur even if you have used this reduces the amount of oxygen product before. Stop use and carried in the blood. This can treated promptly because it seek immediate medical

 pale, gray or blue colored skin care develops: cyanosis)

headache

shortness of breath rapid heart rate

dizziness or lightheadedness fatique or lack of energy

**Allergy Alert:** Do not use on allergies to local anesthetics such as procaine, butacaine, benzocaine or other "caine" patients with a history of

anesthetics.

for more than 7 days unless Do not use

#### MIDWAY SELECT TOPICAL ANESTHETIC

benzocaine gel

**Product Information** 

**HUMAN OTC DRUG** Item Code (Source) NDC:72304-005 **Product Type** 

**Route of Administration DENTAL** 

**Active Ingredient/Active Moiety** 

**Basis of Strength Ingredient Name** Strength BENZOCAINE (UNII: U3RSY48JW5) (BENZOCAINE - UNII:U3RSY48JW5) **BENZOCAINE** 200 mg in 1 g

**Inactive Ingredients** Strength **Ingredient Name** POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P) POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ) **SACCHARIN SODIUM** (UNII: SB8ZUX40TY) FD&C YELLOW NO. 5 (UNII: I753WB2F1M)

FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
FD&C RED NO. 3 (UNII: PN2ZH5LOQY)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C GREEN NO. 3 (UNII: 3P3ONR6O1S)	
D&C GREEN NO. 5 (UNII: 8J6RDU8L9X)	

Product Characteristics			
Color	red	Score	
Shape		Size	
Flavor	STRAWBERRY	Imprint Code	
Contains			

Packaging			
# Item Code	Package Description	Marketing Start Date	Marketing End Date
1 NDC:72304-005-	30 g in 1 JAR; Type 0: Not a Combination Product	06/01/2018	01/18/2025

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part356	06/01/2018	01/18/2025

### Labeler - Midway Dental Supply, LLC (102354446)

## Registrant - Midway Dental Supply, LLC (102354446)

Establishment				
Name	Address	ID/FEI	<b>Business Operations</b>	
Midway Dental Supply, LLC		102354446	manufacture(72304-005)	

Revised: 3/2023 Midway Dental Supply, LLC