

NERVIVE PAIN RELIEVING LIQUID ROLL-ON- lidocaine hcl and menthol liquid
The Procter & Gamble Manufacturing Company

NERVIVE™ PAIN RELIEVING Liquid Roll-on

Drug Facts

Active ingredients

Lidocaine HCl 4%

Menthol 1%

Active ingredients Purpose

Topical anesthetic

Topical analgesic

Use

temporarily relieves minor pain

Warnings

For external use only

Flammable

- keep away from heat and open flame

Do not use

- If you have had an allergic reaction to lidocaine or other local anesthetics
- on large areas of the body or on cut, irritated, blistered, or swollen skin
- on puncture wounds
- for more than one week without consulting a doctor

When using this product

- use only as directed. Read and follow all directions and warnings on this carton.
- avoid contact with eyes and mucous membranes
- rare cases of serious burns have been reported with products of this type

- do not apply to wounds or damaged, broken or irritated skin
- do not bandage tightly or apply local heat (such as heating pads) or a medicated patch to the area of use
- a transient burning sensation may occur upon application but generally disappears in several days
- avoid applying into skin folds

Stop use and ask a doctor if

- condition worsens
- severe burning sensation, redness, rash or irritation develops
- symptoms persist for more than 7 days or clear up and occur again within a few days
- you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children and pets.

If swallowed, get medical help or contact a Poison Control Center right away.

Directions

Shake well before use

adults and children over 12 years:

- apply a thin layer to affected area every 6 to 8 hours
- do not exceed 3 applications in a 24 hour period
- massage into painful area until thoroughly absorbed into skin

AFTER APPLYING, WASH HANDS WITH SOAP AND WATER

children 12 years or younger: ask a doctor

Store at no greater than 25°C (77°F).

Inactive ingredients

alcohol, aminomethyl propanol, C30-45 alkyl cetearyl dimethicone crosspolymer, caprylyl methicone, carbomer homopolymer, ceteth-20 phosphate, cetearyl alcohol, dicetyl phosphate, dimethicone, disodium EDTA, ethylhexyl glycerin, glyceryl stearate, hydroxyethyl acrylate/sodium acryloyldimethyl taurate copolymer, isohexadecane, phenoxyethanol, polysorbate 60, sorbitan isostearate, steareth-2, steareth-21, tocopherol, water

Questions?

1-855-446-4345

DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202

PRINCIPAL DISPLAY PANEL - 2.5 OZ

NERVIVE™

PAIN RELIEVING

LIQUID ROLL-ON

LIDOCAINE HCl & MENTHOL

TARGETS MULTIPLE

NERVE PAIN RECEPTORS

MASSAGING ROLL-ON


NET WT 2.5 OZ (70.9 g)



LIDOCAINE HCl & MENTHOL
TARGETS MULTIPLE
NERVE PAIN RECEPTORS
MASSAGING ROLL-ON



NET WT
2.5 OZ (70.9 g)



Blocks nerve
pain signals

Penetrates nerves &
relieves pain in toes,
feet, & legs

Penetrates nerves &
relieves pain in fingers,
hands, & arms



from the global makers of
nerve care products for
over 50 years.

Removal:

1. Grip the bottle firmly with one hand.
2. Apply your thumb and bent pointer finger to the tabs at the base of the cap and squeeze.
3. Turn the cap while maintaining pressure with your fingers. Remove cap from bottle.

Application:

1. To re-apply, place cap directly onto bottle.
2. Turn until you hear a "click" and continue until cap is snug.

ACTUAL SIZE

Drug Facts	
Active ingredients	Purpose
Lidocaine HCl 4%.....	Topical anesthetic
Menthol 1%.....	Topical analgesic
Use temporarily relieves minor pain	
Warnings	
For external use only	
Flammable • Keep away from heat and open flame	
Do not use	
• If you have had an allergic reaction to lidocaine or other local anesthetics	
• on large areas of the body or on cut, irritated, blistered, or swollen skin	
• on puncture wounds	
• for more than one week without consulting a doctor	
When using this product	
• use only as directed. Read and follow all directions and warnings on this label.	
• avoid contact with eyes and mucous membranes	
• rare cases of serious burns have been reported with products of this type	
• do not apply to wounds or damaged, broken or irritated skin	
• do not bandage tightly or apply local heat (such as heating pads) or a medicated patch to the area of use	
• a transient burning sensation may occur upon application but generally disappears in several days	
• avoid applying into skin folds	
Stop use and ask a doctor if	
• condition worsens • severe burning sensation, redness, rash or irritation develops	
• symptoms persist for more than 7 days or clear up and occur again within a few days	
• you experience signs of skin injury, such as pain, swelling, or blistering where the product was applied	
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children and pets. If swallowed, get medical help or contact a Poison Control Center right away.	
Directions	
Shake well before use	
adults and children over 12 years:	
• apply a thin layer to affected area every 6 to 8 hours	
• do not exceed 3 applications in a 24 hour period	
• massage into painful area until thoroughly absorbed into skin	
AFTER APPLYING, WASH HANDS WITH SOAP AND WATER	
children 12 years or younger: ask a doctor	
Store at no greater than 25°C (77°F)	
Inactive ingredients alcohol, aminomethyl propanol, C30-45 alkyl cetearyl dimethicone crosspolymer, caprylyl methicone, carbomer homopolymer, cetyl-20 phosphate, cetearyl alcohol, dicetyl phosphate, dimethicone, disodium EDTA, ethylhexyl glycerin, glyceryl stearate, hydroxyethyl acrylate/sodium acryloyldimethyl tartrate copolymer, isohexadecane, phenoxyethanol, polysorbate 60, sorbitan isostearate, steareth-2, steareth-21, tocopherol, water	
QUESTIONS? 1-855-446-4345	

DIST. BY: PROCTER & GAMBLE,
CINCINNATI, OH 45202

P&G
Patents: www.pg.com/patents



21129666

NERVIVE PAIN RELIEVING LIQUID ROLL-ON

lidocaine hcl and menthol liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84126-317
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE HYDROCHLORIDE ANHYDROUS (UNII: EC2CNF7XFP) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE HYDROCHLORIDE ANHYDROUS	4 g in 100 g
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	1 g in 100 g

Inactive Ingredients

Ingredient Name	Strength
DIMETHICONE (UNII: 92RU3N3Y1O)	
CAPRYLYL TRISILOXANE (UNII: Q95M2P1KJL)	
CARBOMER INTERPOLYMER TYPE B (ALLYL PENTAERYTHRITOL CROSSLINKED) (UNII: 132584PQMO)	
AMINOMETHYLPROPANOL (UNII: LU49E6626Q)	
PHENOXYETHANOL (UNII: HIE492ZZ3T)	
STEARETH-21 (UNII: 53J3F32P58)	
HYDROXYETHYL ACRYLATE/SODIUM ACRYLOYLDIMETHYL TAURATE COPOLYMER (45000 MPA.S AT 1%) (UNII: 86FQE96TZ4)	
ISOHEXADECANE (UNII: 918X1OUF1E)	
ETHYLHEXYLGLYCERIN (UNII: 147D247K3P)	
CETETH-20 PHOSPHATE (UNII: 921FTA1500)	
GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4)	
ALCOHOL (UNII: 3K9958V90M)	
CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)	
SORBITAN ISOSTEARATE (UNII: 01S2G2C1E4)	
POLYSORBATE 60 (UNII: CAL22UVI4M)	
TOCOPHEROL (UNII: R0ZB2556P8)	
C30-45 ALKYL CETEARYL DIMETHICONE CROSSPOLYMER (UNII: 4ZK9VP326R)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
WATER (UNII: 059QF0KO0R)	
DIHEXADECYL PHOSPHATE (UNII: 2V6E5WN99N)	
STEARETH-2 (UNII: V56DFE46J5)	

Product Characteristics

Color	white	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84126-317-03	1 in 1 CARTON	08/12/2024	
1		70.9 g in 1 BOTTLE; Type 0: Not a Combination Product		
	NDC:84126-317			

2	NDC:84126-317-05	1 in 1 CARTON	03/18/2025	
2		130.4 g in 1 BOTTLE; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017		08/12/2024	

Labeler - The Procter & Gamble Manufacturing Company (004238200)