

**WALGREENS COLD COUGH AND FLU RELIEF- acetaminophen,
dextromethorphan hydrobromide liquid
WALGREENS CO.**

Walgreen Cold , Cough & FluRelief

Active Ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Purpose

Pain reliever/fever reducer

Cough Suppressant

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - minor aches and pains
 - headache
 - cough
 - sore throat
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions.

Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that is accompanied with excessive phlegm (mucus)
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than recommended dose
- **adults and children 12 years and over:** take 30 mL in the dosing cup provided every 4 hours. Do not exceed 150 mL (5 doses) in 24 hours or as directed by a doctor.
- **children under 12 years of age:** do not use

Other information

- **each 30 mL contains:** sodium 38 mg
- store between 20° to 25°C (68° to 77°F)

Inactive ingredients

anhydrous citric acid, Edetate Disodium, FD&C blue #1, FD&C red # 40, flavor, glycerin, Poloxamer, propylene glycol, purified water, Sodium benzoate, potassium Citrate, sorbitol liquid, xanthan gum

Questions or comments? 1-866-467-2748

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

Stop use and ask a doctor if • pain or cough get worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions • do not take more than recommended dose • **adults and children 12 years and over**: take 30 mL in the dosing cup provided every 4 hours. Do not exceed 150 mL (5 doses) in 24 hours or as directed by a doctor.
• **children under 12 years of age**: do not use

Other information • each 30 mL contains: sodium 12 mg • store at room temperature.

Inactive ingredients citric acid anhydrous, disodium edetate, FD&C Blue No.1, FD&C Red No. 40, flavors, glycerin, poloxamer, propylene glycol, purified water, sodium benzoate, potassium citrate, sorbitol liquid, sucralose, xanthan gum.

Questions or comments? 1-866-467-2748

Maximum Strength per 4-hour dose. *Decongestant free. W3ORG0325-F

LOT: 81012WAGLR

EXP: 3 11917 06510 6

DISTRIBUTED BY: WALGREEN CO., DEERFIELD, IL 60015
100% SATISFACTION GUARANTEED walgreens.com
©2025 Walgreen Co. MADE IN THE U.S.A.
WITH DOMESTIC AND IMPORTED INGREDIENTS
ITEM 449309 W000000-0000-0



Cold, Cough & Flu Relief

ACETAMINOPHEN 650 mg / PAIN RELIEVER / FEVER REDUCER
DEXTROMETHORPHAN HBr 20mg / COUGH SUPPRESSANT

Maximum Strength

- For relief of sore throat, fever, headache, cough, body aches & pains
- Cold relief for people with high blood pressure*
- Sugar free
- No added alcohol



12 FL OZ (355 mL) Natural Cherry flavor

Drug Facts Tamper Evident: Do not use if imprinted shrink band is missing or broken

Active ingredients (in each 30 mL)	Purposes
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan hydrobromide 20 mg	Cough suppressant

Uses

- temporarily relieves these symptoms due to a cold or flu: • minor aches & pains • headache • cough • sore throat • temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 4,000 mg of acetaminophen in 24 hours • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product

Allergy Alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include: • skin reddening • blisters • rash • hives • facial swelling • asthma (wheezing) • shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients • in children under 12 years of age

Ask a doctor before use if you have • liver disease • persistent or chronic cough such as occurs with smoking, asthma, or emphysema • cough that is accompanied by excessive phlegm (mucus) • a sodium-restricted diet

NDC 0363-8100-12

Walgreens

Compare to the active ingredients in Coricidin® HBP **Maximum Strength††**

NDC# 0363-8100-12

Cold, Cough & Flu Relief

ACETAMINOPHEN 650 mg /PAIN RELIEVER/FEVER REDUCER

DEXTROMETHORPHAN HBr 20 mg /COUGH SUPPRESSANT

Maximum Strength

- For relief of sore throat, fever, headache, cough, body aches & pains
- Cold relief for people with high blood pressure*
- sugar free
- Alcohol Free

Natural Cherry flavor

*Decongestant free

Maximum Strength per 4-hour dose

12 FL OZ (355 mL)

†Our pharmacist recommend the Walgreens brand. We invite you to compare to national brands.

††This product is not manufactured or distributed by Bayer Healthcare LLC, owner of the registered trademark Coricidin® HBP.

DISTRIBUTED BT: **WALGREEN CO.,**

DEERFIELD, IL 60015

100% SATISFACTION GUARANTEED

Walgreens.com

WALGREENS COLD COUGH AND FLU RELIEF				
acetaminophen, dextromethorphan hydrobromide liquid				
Product Information				
Product Type		HUMAN OTC DRUG	Item Code (Source)	
Route of Administration		ORAL	NDC:0363-8100	
Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)			ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)			DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
Inactive Ingredients				
Ingredient Name				Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)				
EDETATE DISODIUM (UNII: 7FLD91C86K)				
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
GLYCERIN (UNII: PDC6A3C0OX)				
POLOXAMER 105 (UNII: XPP85ZHV76)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
WATER (UNII: 059QF0KO0R)				
SODIUM BENZOATE (UNII: OJ245FE5EU)				
POTASSIUM CITRATE (UNII: EE90ONI6FF)				
SORBITOL (UNII: 506T60A25R)				
SUCRALOSE (UNII: 96K6UQ3ZD4)				
XANTHAN GUM (UNII: TTV12P4NEE)				
Product Characteristics				
Color		red	Score	
Shape			Size	
Flavor		CHERRY	Imprint Code	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-8100-	355 mL in 1 BOTTLE; Type 0: Not a Combination	04/01/2025	

12	Product	04/01/2025	
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/01/2025	

Labeler - WALGREENS CO. (008965063)

Revised: 4/2025

WALGREENS CO.