

**NUVICARE DAYTIME COLD AND FLU SOFTGEL- acetaminophen,
dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid
filled
NUVICARE LLC**

**Nuicare DayTime Cold and Flu Softgel (Acetaminophen 325mg,
Dextromethorphan HBr 10mg, Phenylephrine HCl 5mg)**

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325mg

Dextromethorphan HBr 10mg

Phenylephrine HCl 5mg

Purpose

Acetaminophen 325mg.....Pain reliever/fever reducer
Dextromethorphan HBr 10mg..... Cough suppressant
Phenylephrine HCl 5mg Nasal decongestant

Uses:

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

Warning

Allergy Alert: acetamenophen may cause severe skin recations. Symptoms may include:

■ skin reddening ■ blisters ■ rash. if a skin reactions occurs, stop use and seek medical help right away.

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

■ more than 4 doses in 24 hrs, which is the maximum daily amount for this product ■

with other drugs containing acetaminophen ■ 3 or more alcoholic drinks daily while using this product.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use:

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease
- heart disease
- thyroid disease
- diabetes
- high blood pressure
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin

When using this product, do not exceed recommended dosage.

Stop use and ask a doctor if

- pain, cough or nasal congestion get worse or last more than 7 days
- new symptoms occur
- nervousness, dizziness or sleeplessness occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

■ take only as directed- see **OVERDOSE WARNING**

■ do not exceed 8 softgels per 24 hrs.

Adults and children 12 yrs and over	take 2 softgels with water every 4 hrs.
Children 4 to under 12 yrs	ask a doctor
Children under 4 yrs	do not use

Other information

■ store at room temperature between 20-25°C (68-77 °F)

■ avoid excessive heat, cold and humidity

Inactive ingredients

Polyethylene Glycol400, Propylene Glycol, Povidone (PVP K-30), FD & C Red No. 40, FD & C Yellow No. 6, purified water, Gelatin, Glycerin, Sorbitol Sorbitan solution.

Questions or Comments?

Call 1-718-337-8733 or email: support@nuvicare.com

This product is not manufactured or distributed by PROCTER & GAMBLE, owner of the registered trademark Vicks® Dayquil™ LiquiCaps™.

Distributed by: Nuvicare LLC, Brooklyn, NY 11211, USA



Drug Facts (continued)

When using this product, do not exceed recommended dosage.

Ask a doctor or pharmacist before use if you are ■ taking the blood thinning drug warfarin.

Stop use and ask a doctor if ■ pain, cough, or nasal congestion gets worse or lasts more than 7 days ■ new symptoms occur ■ nervousness, dizziness or sleeplessness occur ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions: Take only as directed – see OVERDOSE WARNING.

■ Do not exceed 8 softgels per 24 hrs.

adults & children age 12 yrs & over ■ take 2 softgels with water every 4 hrs

children 4 to under 12 yrs ■ ask a doctor

children under 4 yrs ■ do not use

Other information: ■ store at room temperature between 20° - 25°C (68°-77°F) ■ avoid excessive heat, cold and humidity.

Inactive ingredients: Polyethylene Glycol -400, Propylene Glycol, Povidone (PVP-K-30), FD & C Red No.40, FD & C Yellow No.6, Purified Water, Gelatin, Glycerin, Sorbitol Sorbitan Solution

Questions or comments?

Call 1 (718) 337-8733 or email: support@nuvicare.com

Drug Facts

Active ingredients (in each softgel)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Nasal decongestant

Uses:

temporarily relieves common cold and flu symptoms: ■ minor aches and pains ■ headache ■ sore throat ■ nasal congestion ■ fever

Warnings:

■ Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash. If a skin reaction occurs, stop use and seek medical help right away. ■ Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hours, which is the maximum daily amount for this product. ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, muscle, or vomiting, consult a doctor promptly.

Do not use: with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if you are taking this product, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have: ■ liver disease ■ diabetes ■ heart disease ■ thyroid disease ■ high blood pressure ■ trouble urinating due to an enlarged prostate gland ■ persistent or chronic cough such as much phlegm (mucus)

occurs with smoking, asthma, or emphysema ■ cough that occurs with too



TAMPER EVIDENT: Do not use if package is opened or if blister unit is torn, broken or shows any sign of tampering.

READ AND KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

nuvi+care®
multi-symptom

NON-DROWSY

Daytime Cold & Flu

Acetaminophen 325mg, Dextromethorphan HBr 10mg
Phenylephrine HCl 5mg

✓ Fever & Headache
✓ Sore Throat & Coughing
✓ Minor Aches & Pains
✓ Nasal Congestion

10 softgels

NDC 84324-018-01

not actual size

†compare to VICKS® DAYQUIL™ COLD & FLU Liquicaps™ active ingredients.

nuvi+care®
multi-symptom

NON-DROWSY

Daytime Cold & Flu

Acetaminophen 325mg, Dextromethorphan HBr 10mg, Phenylephrine HCl 5mg

10 softgels

not actual size

†compare to VICKS® DAYQUIL™ COLD & FLU Liquicaps™ active ingredients.

COATING FREE AREA

Lot No:
Exp Dc:

nuvi+care®
Distributed by: Nuvicare LLC, Brooklyn, NY 11211, U.S.A.
www.nuvicare.com
†This product is not manufactured or distributed by Procter & Gamble® owner of the registered trademark Vicks® DayQuil™ Liquicaps™.

PP0125-01

NUVICARE DAYTIME COLD AND FLU SOFTGEL

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84324-018
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients

Ingredient Name	Strength
POLYETHYLENE GLYCOL 400 (UNII: B697894SGQ)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
POVIDONE (UNII: FZ989GH94E)	
Gelatin (UNII: 2G86QN327L)	
Sorbitol (UNII: 506T60A25R)	
GLYCERIN (UNII: PDC6A3C0OX)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	

Product Characteristics

Color	red	Score	no score
Shape	capsule (Oblong)	Size	18mm
Flavor		Imprint Code	SD14
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84324-018-01	10 in 1 BLISTER PACK; Type 0: Not a Combination Product	05/19/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/19/2025	

Labeler - NUVICARE LLC (119257565)

Revised: 5/2025

NUVICARE LLC