

**WELMATE FAMOTIDINE 20MG- famotidine 20mg tablet**  
**YYBA Corp**

-----  
**Welmate Famotidine 20mg 73581-109**

**Active ingredient (in each tablet)**

Famotidine, USP 20 mg

**Purpose**

Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

**Warnings**

**Allergy alert**

Do not use if you are allergic to famotidine or other acid reducers

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

**Ask a doctor before use if you have**

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with lightheadedness, sweating, or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Acid reducers may interact with certain prescription drugs.

**Stop use and ask a doctor if**

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

**If pregnant or breast-feeding**

ask a health professional before use.

**Directions****adults and children 12 years and over:**

- to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.
- to prevent symptoms, swallow 1 tablet (of 10 mg) with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn
- to prevent symptoms, swallow 1 tablet (of 20 mg) with a glass of water at any time from 10 to 60 minutes before eating food or drinking beverages that cause heartburn
- before eating food or drinking beverages that cause heartburn
- do not use more than 2 tablets in 24 hours

**children under 12 years:**

- ask a doctor

**Inactive Ingredients**

Microcrystalline Cellulose, Pre-gelatinized Starch, Sodium Starch Glycolate, Hydroxypropyl Cellulose, Magnesium Stearate, Hypromellose, Titanium Dioxide, Triacetin, Talc, Macrogol

**Other information**

- read the directions and warnings before use
- keep the carton. It contains important information.
- store at 20°C to 25°C (68°F to 77°F)
- protect from moisture

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL-300's Count**

# DRUG FACTS, DIRECTIONS + DOSAGE INFORMATION

## Drug Facts

Active ingredient (in each tablet)	Purpose
Famotidine USP 20 mg.....	Acid reducer

**Uses**

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

### Warnings

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers.

**Do not use**

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers

### Ask a doctor before use if you have

- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating, or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness
- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- kidney disease

## Drug Facts, cont.

**Ask a doctor or pharmacist before use if you are taking a prescription drug.** Acid reducers may interact with certain prescription drugs.

### Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

## Directions

<b>adults and children 12 years and over:</b>	<ul style="list-style-type: none"> <li>to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew.</li> <li>to prevent symptoms, swallow 1 tablet with a glass of water at any time from <b>15 to 60 minutes</b> before eating food or drinking beverages that cause heartburn</li> <li>do not use more than 2 tablets in 24 hours</li> </ul>
<b>children under 12 years</b>	ask a doctor

## DIRECTIONS FOR USE

To relieve symptoms, swallow 1 tablet with water.  
To prevent symptoms, swallow 1 tablet with water 15-60 minutes before eating or drinking.  
Do not take more than 2 tablets in 24 hours.



 welmate

**PACKAGE LABEL.PRINCIPAL DISPLAY PANEL-100's Count**



100% SATISFACTION GUARANTEED

# ACID REDUCER

MAXIMUM STRENGTH OTC\*

## Famotidine USP 20 mg

Prevents & Relieves Heartburn due to Acid Indigestion

100 TABLETS



FAST ACTING JUST 1 TABLET

DO NOT USE IF PRINTED SEAL UNDER CAP IS MISSING OR DAMAGED

### Drug Facts

**Active ingredient** **Purpose**  
(in each tablet) Famotidine, USP 20 mg.....Acid reducer

### Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain foods and beverages

### Warnings

**Allergy alert:** Do not use if you are allergic to famotidine or other acid reducers.

**Do not use:** if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor. with other acid reducers

**Ask a doctor before use if you have** had heartburn over 3 months. This may be a sign of a more serious condition. heartburn with **lightheadedness, sweating, or dizziness** chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or **lightheadedness** frequent chest pain frequent wheezing, particularly with heartburn unexplained weight loss nausea or vomiting stomach pain kidney disease

**Ask a doctor or pharmacist before use if you are** taking a prescription drug. Acid reducers may interact with certain prescription drugs.



### Drug Facts (continued)

**Stop and ask a doctor if** your heartburn continues or worsens you need to take this product for more than 14 days

**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

### Directions

- adults and children 12 years and over:
  - to relieve symptoms, swallow 1 tablet with a glass of water. Do not chew,
  - to prevent symptoms, swallow 1 tablet with a glass of water at any time from 15 to 60 minutes before eating food or drinking beverages that cause heartburn
  - do not use more than 2 tablets in 24 hours
  - children under 12 years: ask a doctor

### Other information

- read the directions and warnings before use
- store at 20°-25°C (68°-77°F)
- protect from moisture

### Inactive ingredients

Hydroxypropyl cellulose, hydroxymethylcellulose, magnesium stearate, microcrystalline cellulose, polyethylene glycol, pregelatinized starch, sodium starch glycolate, talc, titanium dioxide, triacetin.

### Questions?

call toll-free 1-866-933-6337

\*maximum strength without a prescription

## WELMATE FAMOTIDINE 20MG

famotidine 20mg tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73581-109
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
FAMOTIDINE (UNII: 5QZO15J2Z8) (FAMOTIDINE - UNII:5QZO15J2Z8)	FAMOTIDINE	20 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>HYDROXYPROPYL CELLULOSE, UNSPECIFIED</b> (UNII: 9XZ8H6N6OH)	
<b>TRIACETIN</b> (UNII: XHX3C3X673)	
<b>TALC</b> (UNII: 7SEV7J4R1U)	
<b>SODIUM STARCH GLYCOLATE TYPE A CORN</b> (UNII: AG9B65PV6B)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	

### Product Characteristics

<b>Color</b>	white (OFF-WHITE)	<b>Score</b>	no score
<b>Shape</b>	ROUND (DEBOSED WITH "V" ON ONE SIDE AND "21" ON THE OTHER SIDE)	<b>Size</b>	7mm
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73581-109-20	300 in 1 BOTTLE; Type 0: Not a Combination Product	08/01/2024	
2	NDC:73581-109-01	100 in 1 BOTTLE; Type 0: Not a Combination Product	06/01/2025	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA215822	08/01/2024	

**Labeler** - YYBA Corp (006339772)

**Registrant** - YYBA Corp (006339772)