

CLONIDINE HYDROCHLORIDE- clonidine hydrochloride tablet
NCS HealthCare of KY, Inc dba Vanguard Labs

Clonidine Hydrochloride Tablets, USP 0.1 mg, 0.2 mg, 0.3 mg


Rx only

DESCRIPTION

Clonidine hydrochloride, USP is a centrally acting alpha-agonist hypotensive agent available as tablets for oral administration in three dosage strengths: 0.1 mg, 0.2 mg and 0.3 mg. The 0.1 mg tablet is equivalent to 0.087 mg of the free base.

The inactive ingredients are colloidal silicon dioxide, corn starch, dibasic calcium phosphate, sodium starch glycolate, glycerin, lactose monohydrate, magnesium stearate, povidone.

Clonidine hydrochloride is an imidazoline derivative and exists as a mesomeric compound. The chemical name is 2-(2,6-dichlorophenylamino)-2-imidazoline hydrochloride. The following is the structural formula:



Clonidine hydrochloride is an odorless, bitter, white, crystalline substance soluble in water and alcohol.

CLINICAL PHARMACOLOGY

Clonidine stimulates alpha-adrenoreceptors in the brain stem. This action results in reduced sympathetic outflow from the central nervous system and in decreases in peripheral resistance, renal vascular resistance, heart rate, and blood pressure. Clonidine hydrochloride tablets acts relatively rapidly. The patient's blood pressure declines within 30 to 60 minutes after an oral dose, the maximum decrease occurring within 2 to 4 hours. Renal blood flow and glomerular filtration rate remain essentially unchanged. Normal postural reflexes are intact; therefore, orthostatic symptoms are mild and infrequent.

Acute studies with clonidine hydrochloride in humans have demonstrated a moderate reduction (15% to 20%) of cardiac output in the supine position with no change in the peripheral resistance: at a 45° tilt there is a smaller reduction in cardiac output and a decrease of peripheral resistance. During long term therapy, cardiac output tends to return to control values, while peripheral resistance remains decreased. Slowing of the pulse rate has been observed in most patients given clonidine, but the drug does not alter normal hemodynamic response to exercise.

Tolerance to the antihypertensive effect may develop in some patients, necessitating a reevaluation of therapy.

Other studies in patients have provided evidence of a reduction in plasma renin activity and in the excretion of aldosterone and catecholamines. The exact relationship of these pharmacologic actions to the antihypertensive effect of clonidine has not been fully elucidated.

Clonidine acutely stimulates growth hormone release in both children and adults, but does not produce a chronic elevation of growth hormone with long-term use.

Pharmacokinetics

The pharmacokinetics of clonidine is dose-proportional in the range of 100 to 600 µg. The absolute bioavailability of clonidine on oral administration is 70% to 80%. Peak plasma clonidine levels are attained in approximately 1 to 3 hours.

Following intravenous administration, clonidine displays biphasic disposition with a distribution half-life of about 20 minutes and an elimination half-life ranging from 12 to 16 hours. The half-life increases up to 41 hours in patients with severe impairment of renal function. Clonidine crosses the placental barrier. It has been shown to cross the blood-brain barrier in rats.

Following oral administration about 40% to 60% of the absorbed dose is recovered in the urine as unchanged drug in 24 hours. About 50% of the absorbed dose is metabolized in the liver. Neither food nor the race of the patient influences the pharmacokinetics of clonidine.

The antihypertensive effect is reached at plasma concentrations between about 0.2 and 2.0 ng/mL in patients with normal excretory function. A further rise in the plasma levels will not enhance the antihypertensive effect.

INDICATIONS AND USAGE

Clonidine hydrochloride tablets are indicated in the treatment of hypertension. Clonidine hydrochloride tablets may be employed alone or concomitantly with other antihypertensive agents.

CONTRAINDICATIONS

Clonidine hydrochloride tablets should not be used in patients with known hypersensitivity to clonidine (see **PRECAUTIONS**).

WARNINGS

Withdrawal

Patients should be instructed not to discontinue therapy without consulting their physician. Sudden cessation of clonidine treatment has, in some cases, resulted in symptoms such as nervousness, agitation, headache, and tremor accompanied or followed by a rapid rise in blood pressure and elevated catecholamine concentrations in the plasma. The likelihood of such reactions to discontinuation of clonidine therapy appears to be greater after administration of higher doses or continuation of concomitant beta-blocker treatment and special caution is therefore advised in these situations. Rare instances of hypertensive encephalopathy, cerebrovascular accidents and death have been reported after clonidine withdrawal. When discontinuing therapy with clonidine hydrochloride tablets, the physician should reduce the dose gradually over 2 to 4 days to avoid withdrawal symptomatology.

An excessive rise in blood pressure following discontinuation of clonidine hydrochloride tablets therapy can be reversed by administration of oral clonidine hydrochloride or by intravenous phentolamine. If therapy is to be discontinued in patients receiving a beta-blocker and clonidine concurrently, the beta-blocker should be withdrawn several days before the gradual discontinuation of

clonidine hydrochloride tablets.

Because children commonly have gastrointestinal illnesses that lead to vomiting, they may be particularly susceptible to hypertensive episodes resulting from abrupt inability to take medication.

PRECAUTIONS

General

In patients who have developed localized contact sensitization to clonidine transdermal system, continuation of clonidine transdermal system or substitution of oral clonidine hydrochloride therapy may be associated with the development of a generalized skin rash.

In patients who develop an allergic reaction to clonidine transdermal system, substitution of oral clonidine hydrochloride may also elicit an allergic reaction (including generalized rash, urticaria, or angioedema).

The sympatholytic action of clonidine may worsen sinus node dysfunction and atrioventricular (AV) block, especially in patients taking other sympatholytic drugs. There are post-marketing reports of patients with conduction abnormalities and/or taking other sympatholytic drugs who developed severe bradycardia requiring IV atropine, IV isoproterenol and temporary cardiac pacing while taking clonidine.

In hypertension caused by pheochromocytoma, no therapeutic effect of clonidine hydrochloride tablets can be expected.

Perioperative Use

Administration of clonidine hydrochloride tablets should be continued to within four hours of surgery and resumed as soon as possible thereafter. Blood pressure should be carefully monitored during surgery and additional measures to control blood pressure should be available if required.

Information for Patients

Patients should be cautioned against interruption of clonidine hydrochloride tablets therapy without their physician's advice.

Since patients may experience a possible sedative effect, dizziness, or accommodation disorder with use of clonidine, caution patients about engaging in activities such as driving a vehicle or operating appliances or machinery. Also, inform patients that this sedative effect may be increased by concomitant use of alcohol, barbiturates, or other sedating drugs.

Patients who wear contact lenses should be cautioned that treatment with clonidine hydrochloride tablets may cause dryness of eyes.

Drug Interactions

Clonidine may potentiate the CNS-depressive effects of alcohol, barbiturates or other sedating drugs. If a patient receiving clonidine hydrochloride is also taking tricyclic antidepressants, the hypotensive effect of clonidine may be reduced, necessitating an increase in the clonidine dose. If a patient receiving clonidine is also taking neuroleptics, orthostatic regulation disturbances (e.g., orthostatic hypotension, dizziness, fatigue) may be induced or exacerbated.

Monitor heart rate in patients receiving clonidine concomitantly with agents known to affect sinus node function or AV nodal conduction, e.g., digitalis, calcium channel blockers and beta-blockers. Sinus bradycardia resulting in hospitalization and pacemaker insertion has been reported in association with the use of clonidine concomitantly with diltiazem or verapamil.

Amitriptyline in combination with clonidine enhances the manifestation of corneal lesions in rats (see

Toxicology).

Based on observations in patients in a state of alcoholic delirium it has been suggested that high intravenous doses of clonidine may increase the arrhythmogenic potential (QT- prolongation, ventricular fibrillation) of high intravenous doses of haloperidol. Causal relationship and relevance for clonidine oral tablets have not been established.

Toxicology

In several studies with oral clonidine hydrochloride, a dose-dependent increase in the incidence and severity of spontaneous retinal degeneration was seen in albino rats treated for six months or longer. Tissue distribution studies in dogs and monkeys showed a concentration of clonidine in the choroid.

In view of the retinal degeneration seen in rats, eye examinations were performed during clinical trials in 908 patients before, and periodically after, the start of clonidine therapy. In 353 of these 908 patients, the eye examinations were carried out over periods of 24 months or longer. Except for some dryness of the eyes, no drug-related abnormal ophthalmological findings were recorded and, according to specialized tests such as electroretinography and macular dazzle, retinal function was unchanged.

In combination with amitriptyline, clonidine hydrochloride administration led to the development of corneal lesions in rats within 5 days.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Chronic dietary administration of clonidine was not carcinogenic to rats (132 weeks) or mice (78 weeks) dosed, respectively, at up to 46 or 70 times the maximum recommended daily human dose as mg/kg (9 or 6 times the MRDHD on a mg/m² basis). There was no evidence of genotoxicity in the Ames test for mutagenicity or mouse micronucleus test for clastogenicity.

Fertility of male or female rats was unaffected by clonidine doses as high as 150 µg/kg (approximately 3 times MRDHD). In a separate experiment, fertility of female rats appeared to be affected at dose levels of 500 to 2000 µg/kg (10 to 40 times the oral MRDHD on a mg/kg basis; 2 to 8 times the MRDHD on a mg/m² basis).

Pregnancy

Teratogenic Effects: Pregnancy Category C

Reproduction studies performed in rabbits at doses up to approximately 3 times the oral maximum recommended daily human dose (MRDHD) of clonidine hydrochloride tablets produced no evidence of a teratogenic or embryotoxic potential in rabbits. In rats, however, doses as low as 1/3 the oral MRDHD (1/15 the MRDHD on a mg/m² basis) of clonidine were associated with increased resorptions in a study in which dams were treated continuously from 2 months prior to mating. Increased resorptions were not associated with treatment at the same time or at higher dose levels (up to 3 times the oral MRDHD) when the dams were treated on gestation days 6 to 15. Increases in resorption were observed at much higher dose levels (40 times the oral MRDHD on a mg/kg basis; 4 to 8 times the MRDHD on a mg/m² basis) in mice and rats treated on gestation days 1 to 14 (lowest dose employed in the study was 500 µg/kg).

No adequate, well-controlled studies have been conducted in pregnant women. Clonidine crosses the placental barrier (see **CLINICAL PHARMACOLOGY, Pharmacokinetics**). Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Nursing Mothers

As clonidine hydrochloride is excreted in human milk, caution should be exercised when clonidine hydrochloride tablets are administered to a nursing woman.

Pediatric Use

Safety and effectiveness in pediatric patients have not been established in adequate and well-controlled trials (see **WARNINGS, Withdrawal**).

ADVERSE REACTIONS

Most adverse effects are mild and tend to diminish with continued therapy. The most frequent (which appear to be dose-related) are dry mouth, occurring in about 40 of 100 patients; drowsiness, about 33 in 100; dizziness, about 16 in 100; constipation and sedation, each about 10 in 100.

The following less frequent adverse experiences have also been reported in patients receiving clonidine hydrochloride tablets, but in many cases patients were receiving concomitant medication and a causal relationship has not been established.

Body as a Whole: Fatigue, fever, headache, pallor, weakness, and withdrawal syndrome. Also reported were a weakly positive Coombs' test and increased sensitivity to alcohol.

Cardiovascular: Bradycardia, congestive heart failure, electrocardiographic abnormalities (*i.e.*, sinus node arrest, junctional bradycardia, high degree AV block and arrhythmias), orthostatic symptoms, palpitations, Raynaud's phenomenon, syncope, and tachycardia. Cases of sinus bradycardia and atrioventricular block have been reported, both with and without the use of concomitant digitalis.

Central Nervous System: Agitation, anxiety, delirium, delusional perception, hallucinations (including visual and auditory), insomnia, mental depression, nervousness, other behavioral changes, paresthesia, restlessness, sleep disorder, and vivid dreams or nightmares.

Dermatological: Alopecia, angioneurotic edema, hives, pruritus, rash, and urticaria.

Gastrointestinal: Abdominal pain, anorexia, constipation, hepatitis, malaise, mild transient abnormalities in liver function tests, nausea, parotitis, pseudo-obstruction (including colonic pseudo-obstruction), salivary gland pain, and vomiting.

Genitourinary: Decreased sexual activity, difficulty in micturition, erectile dysfunction, loss of libido, nocturia, and urinary retention.

Hematologic: Thrombocytopenia.

Metabolic: Gynecomastia, transient elevation of blood glucose or serum creatine phosphokinase, and weight gain.

Musculoskeletal: Leg cramps and muscle or joint pain.

Oro-otolaryngeal: Dryness of the nasal mucosa.

Ophthalmological: Accommodation disorder, blurred vision, burning of the eyes, decreased lacrimation, and dryness of eyes.

OVERDOSAGE

Hypertension may develop early and may be followed by hypotension, bradycardia, respiratory depression, hypothermia, drowsiness, decreased or absent reflexes, weakness, irritability and miosis. The frequency of CNS depression may be higher in children than adults. Large overdoses may result in reversible cardiac conduction defects or dysrhythmias, apnea, coma and seizures. Signs and symptoms of overdose generally occur within 30 minutes to two hours after exposure. As little as 0.1 mg of clonidine has produced signs of toxicity in children.

There is no specific antidote for clonidine overdose. Clonidine overdose may result in the rapid development of CNS depression; therefore, induction of vomiting with ipecac syrup is not recommended. Gastric lavage may be indicated following recent and/or large ingestions. Administration of activated charcoal and/or a cathartic may be beneficial. Supportive care may include atropine sulfate

for bradycardia, intravenous fluids and/or vasopressor agents for hypotension and vasodilators for hypertension. Naloxone may be a useful adjunct for the management of clonidine-induced respiratory depression, hypotension and/or coma; blood pressure should be monitored since the administration of naloxone has occasionally resulted in paradoxical hypertension. Tolazoline administration has yielded inconsistent results and is not recommended as first-line therapy. Dialysis is not likely to significantly enhance the elimination of clonidine.

The largest overdose reported to date involved a 28-year old male who ingested 100 mg of clonidine hydrochloride powder. This patient developed hypertension followed by hypotension, bradycardia, apnea, hallucinations, semicoma, and premature ventricular contractions. The patient fully recovered after intensive treatment. Plasma clonidine levels were 60 ng/mL after 1 hour, 190 ng/mL after 1.5 hours, 370 ng/mL after 2 hours, and 120 ng/mL after 5.5 and 6.5 hours. In mice and rats, the oral LD₅₀ of clonidine is 206 and 465 mg/kg, respectively.

DOSAGE AND ADMINISTRATION

Adults

The dose of clonidine hydrochloride tablets must be adjusted according to the patient's individual blood pressure response. The following is a general guide to its administration.

Initial Dose

0.1 mg tablet twice daily (morning and bedtime). Elderly patients may benefit from a lower initial dose.

Maintenance Dose

Further increments of 0.1 mg per day may be made at weekly intervals if necessary until the desired response is achieved. Taking the larger portion of the oral daily dose at bedtime may minimize transient adjustment effects of dry mouth and drowsiness. The therapeutic doses most commonly employed have ranged from 0.2 mg to 0.6 mg per day given in divided doses. Studies have indicated that 2.4 mg is the maximum effective daily dose, but doses as high as this have rarely been employed.

Renal Impairment

Patients with renal impairment may benefit from a lower initial dose. Patients should be carefully monitored. Since only a minimal amount of clonidine is removed during routine hemodialysis, there is no need to give supplemental clonidine following dialysis.

HOW SUPPLIED

Clonidine Hydrochloride Tablets USP are supplied as:

0.1 mg

White rounded off oval shaped, biconvex tablets, with score line having "U" and "135" debossed across the score line on one side and plain on other side.

Blistercards of 30: NDC 0615-7947-39

0.2 mg

White rounded off oval shaped, biconvex tablets, with score line having "U" and "136" debossed across the score line on one side and plain on other side.

Blistercards of 30: NDC 0615-7948-39

Unit-dose Boxes of 30: NDC 0615-7948-30

0.3 mg

White rounded off oval shaped, biconvex tablets with score line having "U" and "137" debossed across the score line on one side and plain on other side.

Blistercards of 30: NDC 0615-8164-39

Store at 20°- 25°C (68°- 77°F) [See USP Controlled Room Temperature].

Dispense in a tight, light-resistant container.

Please address medical inquires to Unichem's toll free # 1-866-562-4616.

Manufactured by:

UNICHEM LABORATORIES LTD.

Pilerne Ind. Estate, Pilerne, Bardez, Goa 403 511, India

Manufactured for:



Hasbrouch Heights, NJ 07604.

05-R-10/2017

13009854

PRINCIPAL DISPLAY PANEL - 0.1MG



VLI NDC 0615-7947-39
CLONIDINE
HYDROCHLORIDE
 TABS, USP 0.1 mg



LOT
 7947
 EXP
 12/31/13

Rx
 Only

QTY
 30

MFG BY UNICHEM
 (NDC 29300-135-10)

PKG BY VANGARD
 GLASGOW, KY 42141

(UNICHEM NDC 29300-135-10)

Clonidine
Hydrochloride
 Tabs, USP 0.1 mg

Received: _____

31	24	16	8
30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
24	17	9	1

START DATE _____ START TIME _____

STORE AT 20°-25°C (68°-77°F)
 (SEE USP CONTROLLED ROOM TEMPERATURE)
 Dispense in a tight, light-resistant container.

See Package Insert Binder for Dosage Information
 FOR INSTITUTIONAL USE ONLY

			16	8
30	23	15	7	
29	22	14	6	
28	21	13	5	
27	20	12	4	
26	19	11	3	
25	18	10	2	
24	17	9	1	

<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>12000 Rev 001 Vanguard Labs Glasgow, KY 42141 </p>
<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>
<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>
<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>
<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>
<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>
<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TABS, USP 0.1 mg LOT 7947 </p>

Omnicare The overall configuration of this package is a trademark of Omnicare, Inc.

PRINCIPAL DISPLAY PANEL - 0.2MG



VLI NDC 0615-7948-39
CLONIDINE
HYDROCHLORIDE
TABS, USP 0.2 mg



LOT
 7948
 EXP
 293001360139
 12394-39

Rx
 Only

QTY
 30

MFG BY UNICHEM
 (NDC 29300-136-01)

PKG BY VANGARD
 GLABOW, KY 42141

(UNICHEM NDC 29300-136-01)

Clonidine
Hydrochloride
Tab, USP 0.2 mg

Received: _____

31	24	16	8
30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
24	17	9	1

START DATE _____ START TIME _____

STORE AT 20°-25°C (68°-77°F)
 (SEE USP CONTROLLED ROOM TEMPERATURE)
 Dispense in a tight, light-resistant container.


See Package Insert Binder for Dosage Information
 FOR INSTITUTIONAL USE ONLY


		16	8
30	23	15	7
29	22	14	6
28	21	13	5
27	20	12	4
26	19	11	3
25	18	10	2
24	17	9	1


<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>32003 Rev 00 Vanguard Labs Glasgow, KY 42141 </p>
<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>
<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>
<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>
<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>
<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>
<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>	<p>Fig by Vanguard, Glasgow, KY 42141 CLONIDINE HYDROCHLORIDE TAB, USP 0.2 mg LOT 7948 </p>





The overall configuration of this package is a trademark of Omnicare, Inc.


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601

Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601

Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948


 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601

Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601

Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601

Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601

Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601

Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601

Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948


 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948


 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948


 2930013601



Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601

Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601


Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601

Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601

Pkg By Vanguard, Glasgow, KY 42141
 CLONIDINE
 HYDROCHLORIDE
 TAB, USP 0.2 mg
 LOT 7948

 2930013601

VLI NDC 0615-7948-30 Clonidine HCl Tablets USP 0.2 mg		VLI NDC 0615-7948-30 Clonidine HCl Tablets USP 0.2 mg	
QTY 30	Rx only STORE AT 20°-25°C (68°-77°F) (SEE USP CONTROLLED ROOM TEMPERATURE) Dispense in a tight, light-resistant container.	FOR INSTITUTIONAL USE ONLY QTY LOT: 30 EXP:	Manufactured By Unichem for Unichem NDC 293001360130
LOT: EXP:		Repackaged by Vanguard Labs Glasgow, KY 42141 Dosage: See my.omicare.com for Insert or Label Information 	7948-3A 2 00001-01 293001360130

PRINCIPAL DISPLAY PANEL - 0.3MG



(Unichem NDC 29300-137-01)
**Clonidine HCl
 Tabs USP 0.3 mg**

Rx Only

Mfg By Unichem
 (NDC 29300-137-01)
 PEG BY VANGUARD
 GLASGOW, KY 42141

QTY
30

			16	8
30	23	15	7	
29	22	14	6	
28	21	13	5	
27	20	12	4	
26	19	11	3	
25	18	10	2	
24	17	9	1	



The overall configuration of this package is a trademark of Omnicare, Inc.

Received: _____

24	16	8
31	23	7
30	22	6
29	21	5
28	20	4
27	19	3
26	18	2
25	17	1

Start Date _____ Start Time _____

STORE AT 20°-25°C (68°-77°F)
 (SEE USP CONTROLLED ROOM TEMPERATURE)
 Dispense in a tight, light-resistant container.

See Package Insert Binder for Dosage Information
 FOR INSTITUTIONAL USE ONLY

Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	12897 Rev 00 Vanguard Label Glasgow, KY 42141
Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164
Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164
Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164
Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164
Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164
Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164 	Pkg by Vanguard, Glasgow, KY 42141 Clonidine HCl Tab USP 0.3 mg LOT B164

CLONIDINE HYDROCHLORIDE

clonidine hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0615-7947(NDC:29300-135)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CLONIDINE HYDROCHLORIDE (UNII: W7616XXF06) (CLONIDINE - UNII:MN3L5RMN02)	CLONIDINE HYDROCHLORIDE	0.1 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
GLYCERIN (UNII: PDC6A3C0OX)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	

MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	OVAL	Size	8mm
Flavor		Imprint Code	U;135
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0615-7947-39	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	05/13/2016	11/30/2020

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078895	09/21/2009	11/30/2020

CLONIDINE HYDROCHLORIDE

clonidine hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0615-7948(NDC:29300-136)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CLONIDINE HYDROCHLORIDE (UNII: W76I6XXF06) (CLONIDINE - UNII:MN3L5RMN02)	CLONIDINE HYDROCHLORIDE	0.2 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
GLYCERIN (UNII: PDC6A3C0OX)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	OVAL	Size	8mm
Flavor		Imprint Code	U;136
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0615-7948-30	6 in 1 BOX, UNIT-DOSE	06/09/2016	11/30/2020
1		5 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0615-7948-39	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	05/17/2016	11/30/2020

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078895	09/21/2009	11/30/2020

CLONIDINE HYDROCHLORIDE

clonidine hydrochloride tablet

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0615-8164(NDC:29300-137)
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CLONIDINE HYDROCHLORIDE (UNII: W76I6XXF06) (CLONIDINE - UNII:MN3L5RMN02)	CLONIDINE HYDROCHLORIDE	0.3 mg

Inactive Ingredients

Ingredient Name	Strength
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
GLYCERIN (UNII: PDC6A3C0OX)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE, UNSPECIFIED (UNII: FZ989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A POTATO (UNII: 5856J3G2A2)	
STARCH, CORN (UNII: O8232NY3SJ)	

Product Characteristics

Color	WHITE	Score	2 pieces
Shape	OVAL	Size	10mm
Flavor		Imprint Code	U;137
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0615-8164-39	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	07/10/2018	05/31/2020

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA078895	07/10/2018	05/31/2020

Labeler - NCS HealthCare of KY, Inc dba Vanguard Labs (050052943)

Establishment

Name	Address	ID/FEI	Business Operations
NCS HealthCare of KY, Inc dba Vanguard Labs		050052943	REPACK(0615-7947, 0615-7948, 0615-8164)

Revised: 12/2018

NCS HealthCare of KY, Inc dba Vanguard Labs