

WART REMOVER- salicylic acid, chlorhexidine ointment
Dongkou County Hecheng Network Technology Co., Ltd.

Wart Remover

SALICYLIC ACID 1%
CHLORHEXIDINE 1%

wart Remove

Keep out of reach of children. If swallowed, get medical help or contact a
Poison control center (1-800-222-1222) right away.

After cleansing the areas to be treated with lukewarm water, apply proper amount of the cream to the surface of the affected area and gently rub in for 5-10 minutes until it is fully absorbed. It is recommended to use this product continuously 2-3 times a day until the warts fall off.

For external use only.

Keep away from fire and flame

wash the affected area

may soak the wart in warm water for 5 minutes

dry area thoroughly

using the applicator (cotton swab), apply a layer of ointment to sufficiently cover each wart
allow it to fully absorb and cover it with a bandage as needed.

repeat this procedure once or twice daily as needed (until the wart is removed) for up to 12 weeks

Store in a dry and cool place

ANGELICA DAHURICA ROOT
CITRUS RETICULATA WHOLE
FORSYTHIA SUSPensa ROOT
PLATYCODON GRANDIFLORUS WHOLE
ASARUM SIEBOLDII
CETOSTEARYL ALCOHOL
SOPHORA FLAVESCENS ROOT
PORTULACA OLERACEA WHOLE
ATRACTYLODES MACROCEPHALA ROOT
ARCTIUM LAPPA WHOLE
BAMBUSA VULGARIS LEAF
BORNEOL
CNIDIUM MONNIERI FRUIT
HONEY
ALCOHOL

Drug Facts

Active ingredient	Purpose
salicylic acid 1%.....	wart Remove!
Chlorhexidine di(acetate) 1%.....	art Remove

Usage:

After cleansing the areas to be treat-edwithlukewarm water, apply proper amou-ntof thec ream to the surface of the affectedareaandgently rub in for 5-10 minutes untillit is fully absorbed.it is recommended tousethis productcontinuously 2-3 times aday un-til the warts falloff.

Warnings

For external use only.
Keep away from fire and flame

When using this product

- e avoid contact with eyes. If the product getsinto the eye, flush with water for 15 minites
- avoid inhaling vapors
- cap the tube tightlv
- out of direct sunlight and store at roomtemperature away from heat

Stop use and ask a doctor

- if discomfort persists.
- if you have diabetes or poor bloodcirculation.

Keep out of reach of children.
If swallowed, get medical help or contact a PoisonControlCenter (1-800-222-1222)rightaway.

Directions


- wash the affected area
- may soak the wart in warm water for 5minutes
- dry area thoroughly
- using the applicator(cotton swab), apply alayer of ointment to sufficiently cover eachwart
- allow it to fully absorb and cover it with dbandage as needed.
- repeat this procedure once or twice dailyas needed (until the wart is removed) for upto 12 weeks

Ingredients:

Angelica Dahurica Root, citrus Reliculata Whole, forsythia Suspensa Root, platycodon Grandiflorus Whole, asarum Sieboldii, cetostearyl Alcohol, sophora Flavescens Root, portulaca Oleracea Whole, atracylodes Macrocephala Root, arctium Lappa Whole, bambusa Vulgaris Leaf, borneol, cnidium Monnieri Fruit, Honey, Alcohol

WoTeReVo

Wart remover OINTMENT CREAM



Net:20g/0.68fl.oz

EXTERNAL USE

NATURAL REMOVES WARTS
FAST PENETRATES



WoTeReVo

Wart remover OINTMENT CREAM



Net:20g/0.68fl.oz

EXTERNAL USE



Wart remover

HEALTHY TREATMENT

CREAM



DERMATOLOGY INSPIRED CARE

*FAST PENETRATES
NATURAL REMOVES WARTS*

Net:20g/0.6fl.OZ

WART REMOVER

When using this product
avoid contact with eyes. If the
product gets into the eye, flush
with water for 15 minutes
avoid inhaling vapors
cap the tube tightly
out of direct sunlight and store
at room temperature away from
heat

Usage:
After cleansing the areas to be
treated with lukewarm water,
apply proper amount of the cream
to the surface of the
affected area and gently rub in
for 5-10 minutes until it is fully
absorbed. It is recommended
to use this product continuously
2-3 times a day until the warts
fall off.



WART REMOVER

salicylic acid, chlorhexidine ointment

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:84288-005
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ)	SALICYLIC ACID	1 g in 100 g
CHLORHEXIDINE (UNII: R4KO0DY52L) (CHLORHEXIDINE - UNII:R4KO0DY52L)	CHLORHEXIDINE	1 g in 100 g

Inactive Ingredients	
Ingredient Name	Strength
ATRACTYLODES MACROCEPHALA ROOT (UNII: 08T3N29QJB)	
ARCTIUM LAPPА WHOLE (UNII: 73070DU1LA)	
CITRUS RETICULATA WHOLE (UNII: 00OX7CMF92)	
ASARUM SIEBOLDII (UNII: T1134STW1K)	
SOPHORA FLAVESCENS ROOT (UNII: IYR6K8KQ5K)	
BAMBUSA VULGARIS LEAF (UNII: EMY54R518C)	
HONEY (UNII: Y9H1V576FH)	
CNIDIUM MONNIERI FRUIT (UNII: V1A3S3CUS)	
PLATYCODON GRANDIFLORUS WHOLE (UNII: WC73QE9274)	
CETOSTEARYL ALCOHOL (UNII: 2DMT128M1S)	
FORSYTHIA SUSPENSА ROOT (UNII: YM9E56VP46)	
BORNEOL (UNII: M89NIB437X)	
PORTULACA OLERACEA WHOLE (UNII: D5J3623SV2)	
ALCOHOL (UNII: 3K9958V90M)	
ANGELICA DAHURICA ROOT (UNII: 1V63N2S972)	

Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84288-005-01	20 g in 1 TUBE; Type 0: Not a Combination Product	08/02/2024	

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M028	08/02/2024	

Labeler - Dongkou County Hecheng Network Technology Co., Ltd. (637818093)

Establishment			
Name	Address	ID/FEI	Business Operations
Dongkou County Hecheng Network Technology Co., Ltd.		637818093	manufacture(84288-005)