

**MAGNESIUM CITRATE SALINE LAXATIVE - CHERRY FLAVOR- magnesium citrate solution**

**Universal Distribution Center LLC**

-----  
**Magnesium Citrate Saline Laxative Oral Solution - Cherry Flavor**

***Drug Facts***

**Active ingredient (in each fl oz)**

Magnesium citrate 1.745 g

**Purpose**

Saline laxative

**Uses:**

- For relief of occasional constipation (irregularity).
- Generally produces bowel movement in ½ to 6 hours

**Warnings:**

**Ask a doctor before use if you have**

- Kidney disease • a magnesium restricted diet
- Abdominal pain, nausea, or vomiting
- Noticed a sudden change in bowel habits that persists over a period of 2 weeks
- Already used a laxative for a period longer than 1 week

**Ask a doctor or pharmacist before use if you are**

- Taking any other drug.
- Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

**Stop use and ask a doctor if you have rectal** bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions:***

- shake well before using • drink a full glass (8 ounces) of liquid each dose • may be taken as a single dose or in divided doses
-

adults and children 12 years of age and over	6.5 to 10 fl oz maximum 10 fl oz in 24 hours
children 6 to under 12 years of age	3 to 7 fl oz maximum 7 fl oz in 24 hours
children 2 to under 6 years of age	2 to 3 fl oz maximum 3 fl oz in 24 hours
children under 2 years of age	ask a doctor

***Other Information:***

- each fl oz contains: magnesium 290 mg
- each fl oz contains: sodium 1 mg

store at temperatures between 46° and 86°F (8° and 30°C)

***Inactive ingredients:***

benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Sugar Free

Alcohol content 0.025%  
(From Flavoring)

**MADE IN USA**

**DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN, OR SEPARATED FROM NECKRING**

Mfd for and Distributed By:  
**Universal Distribution Center LLC**  
330 Applegarth Road,  
Monroe Township, NJ 08831  
**[www.universaldc.com](http://www.universaldc.com)**

**Packaging**

### Drug Facts

store at temperatures between 46° and 86°F (8° and 30°C)

#### Directions:

■ shake well before using ■ drink a full glass (8 ounces) of liquid with each dose ■ may be taken as a single daily dose or in divided doses

adults and children 12 years of age and over 6.5 to 10 fl oz maximum 10 fl oz in 24 hours

children 6 to under 12 years of age 3 to 7 fl oz maximum 7 fl oz in 24 hours

children 2 to under 6 years of age 2 to 3 fl oz maximum 3 fl oz in 24 hours

children under 2 years of age ask a doctor

#### Other Information:

■ each fl oz contains: magnesium 290 mg

■ each fl oz contains: sodium 1 mg

**Inactive ingredients:** benzoic acid, citric acid, disodium EDTA, flavor, sucralose, water

Mfd for and Distributed By:  
Universal Distribution Center LLC  
330 Applegarth Road,  
Monroe Township, NJ 08831  
www.universaldc.com



# Magnesium Citrate

## Saline Laxative Oral Solution

Cherry Flavor

Sugar Free

Alcohol content 0.025%  
(From Flavoring)



10 FL. OZ (296mL)



### Drug Facts (continued)

**Active ingredient (in each fl oz) Purpose**  
Magnesium citrate 1.745 g ....Saline laxative

**Uses:** ■ For relief of occasional constipation (irregularity).

■ Generally produces bowel movement in ½ to 6 hours

#### Warnings:

Ask a doctor before use if you have

■ Kidney disease ■ a magnesium restricted diet

■ Abdominal pain, nausea, or vomiting

■ Noticed a sudden change in bowel habits that persists over a period of 2 weeks

■ Already used a laxative for a period longer than 1 week

Ask a doctor or pharmacist before use if you are

■ Taking any other drug.

■ Take this product 2 or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if you have rectal bleeding or failure to have a bowel movement after use. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

DO NOT USE IF TAMPER EVIDENT TWIST-OFF CAP IS MISSING, BROKEN, OR SEPARATED FROM NECKRING

## MAGNESIUM CITRATE SALINE LAXATIVE - CHERRY FLAVOR

magnesium citrate solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:52000-451
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>MAGNESIUM CITRATE</b> (UNII: RHO26O1T9V) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM CITRATE	0.059 g in 1 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BENZOIC ACID</b> (UNII: 8SKNOB0MIM)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>WATER</b> (UNII: 059QF0KO0R)	

### Product Characteristics

<b>Color</b>		<b>Score</b>	
<b>Shape</b>		<b>Size</b>	

<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:52000-451-10	296 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	03/16/2026	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	03/16/2026	

**Labeler** - Universal Distribution Center LLC (019180459)

Revised: 3/2026

Universal Distribution Center LLC