

**OTIS CLAPP VALIHIST- acetaminophen, chlorpheniramine maleate, phenylephrine hydrochloride tablet, film coated**  
**MEDIQUE MEDICIDIN D- acetaminophen, chlorpheniramine maleate, phenylephrine hydrochloride tablet, film coated**  
**Unifirst First Aid Corporation**

-----

## **Medique Medicidin D/ Otis Clapp Valihist**

### ***Drug Facts***

#### ***Active ingredients (in each tablet)***

Acetaminophen 325 mg

Chlorpheniramine Maleate 2 mg

Phenylephrine Hydrochloride 5 mg

#### ***Purpose***

Pain reliever/fever reducer

Antihistamine

Nasal decongestant

#### ***Uses***

Temporarily relieves

- minor aches and pains
- headache
- nasal congestion
- sinus congestion and pressure
- runny nose
- sneezing

Temporarily

- reduces fever
- relieves runny nose and reduces sneezing, itching of the nose and throat and itchy, watery eyes due to hay fever
- helps clear nasal and/or sinus passages

#### ***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- for more than 10 days for pain unless directed by a doctor
- if you ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

liver disease  
thyroid disease  
diabetes  
high blood pressure  
heart disease  
glaucoma  
chronic bronchitis or emphysema  
difficulty in urination due to enlargement of the prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- may cause drowsiness
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- use caution when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.**

### **Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away (1-

800-222-12220. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- **do not use more than directed**

**Adults and children (12 years and older):** Take 2 tablets every 4 to 6 hours. Do not take more than 10 tablets in 24 hours.

**Children under 12 years:** Ask a doctor

***Other information***

- store at room temperature 59°-86°F (15°-30°C)
- tamper evident sealed packets
- do not use any opened or torn packets
- avoid excessive heat and humidity

***Inactive ingredients***

corn starch, crospovidone, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid

**Questions or comments?** 1-800-634-7680

**Principal Display Panel - Otis Clapp Valihist Label**

Otis Clapp

Quality and Integrity Since 1840

VALIHIST™

Decongestant-Antihistamine

Pain Reliever-Fever Reducer

Antihistamine- Decongestant

For Allergies, Colds, Runny Nose

See Warnings and Directions on Side Panel

This Package is for Households without Young Children.

Acetaminophen 325 mg,

Chlorpheniramine Maleate 2mg

Phenylephrine HCl 5 mg

Tear Out Along Perforation To Dispense

PROFESSIONAL HEALTHCARE

300 TABLETS (150 PACKETS OF 2)



**Medicine**  
**Medicidin-D**  
Gold & Allergy Relief

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

- ✓ Pain Reliever/Fever Reducer • Acetaminophen 325 mg
- ✓ Antihistamine • Chlorpheniramine Maleate 2 mg
- ✓ Nasal Decongestant • Phenylephrine HCl 5 mg

**Medicine**  
**Medicidin-D**  
Gold & Allergy Relief

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.

Pull to Open

- ✓ Pain Reliever/Fever Reducer • Acetaminophen 325 mg
- ✓ Antihistamine • Chlorpheniramine Maleate 2 mg
- ✓ Nasal Decongestant • Phenylephrine HCl 5 mg

100 Tablets (50 x 2)

1203380X  
Rev. 9/021  
Mg. 1

**Drug Facts**

**Active ingredients (in each tablet)**

Active ingredients	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Chlorpheniramine Maleate 2 mg	Antihistamine
Phenylephrine Hydrochloride 5 mg	Nasal decongestant

**Uses**

- Temporarily relieves minor aches and pains
- Relieves sinus congestion and pressure
- Temporarily relieves runny nose and reduces sneezing, itching of the nose and throat and itchy, watery eyes due to hay fever
- Helps clear nasal and/or sinus passages

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product.

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription) if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist
- for more than 10 days unless directed by a doctor
- if you have ever had an allergic reaction to this product or any of its ingredients
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

(continued on opposite panel)

**Drug Facts (continued)**

**Ask a doctor before use if you have**

- liver disease
- heart disease
- difficulty in urination due to enlargement of the prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin

**When using this product**

- do not exceed recommended dosage
- may cause drowsiness
- avoid alcoholic beverages

**Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occur
- fever gets worse or lasts for more than 3 days
- if pregnant or breast-feeding, ask a health professional before use

**Keep out of reach of children.**

**Overdose warning**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- do not use more than directed

Adults and children (10 years and older)	Children under 12 years
Take 2 tablets every 4 to 6 hours. Do not take more than 10 tablets in 24 hours.	Ask a doctor

**Drug Facts (continued)**

**Other information**

- store at room temperature 59°-86°F (15°-30°C)
- tamper evident sealed packets
- do not use any opened or torn packets
- avoid excessive heat and humidity

**Inactive ingredients**

corn starch, croscellose, hypromellose, microcrystalline cellulose, polyethylene glycol, povidone, pregelatinized starch, stearic acid

**Questions or comments?** 1-800-634-7680

Reorder #12033  
Manufactured for:  
Medique Products  
Fort Myers, FL 33967  
1-800-634-7680  
www.medicqueproducts.com

## OTIS CLAPP VALIHIST

acetaminophen, chlorpheniramine maleate, phenylephrine hydrochloride tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-543
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>CROSPVIDONE</b> (UNII: 68401960MK)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	

### Product Characteristics

<b>Color</b>	white (white)	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND)	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	AZ;275
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-543-03	150 in 1 BOX	12/30/2008	
1	NDC:47682-543-99	2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-543-99	2 in 1 PACKET; Type 0: Not a Combination Product	12/30/2008	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/30/2008	

## MEDIQUE MEDICIDIN D

acetaminophen, chlorpheniramine maleate, phenylephrine hydrochloride tablet, film coated

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:47682-120
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>CELLULOSE, MICROCRYSTALLINE</b> (UNII: OP1R32D61U)	
<b>HYPROMELLOSES</b> (UNII: 3NXW29V3WO)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>CROSPVIDONE</b> (UNII: 68401960MK)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	

## Product Characteristics

<b>Color</b>	white (white)	<b>Score</b>	no score
<b>Shape</b>	ROUND (ROUND)	<b>Size</b>	11mm
<b>Flavor</b>		<b>Imprint Code</b>	AZ;275
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:47682-120-64	12 in 1 BOX	12/30/2008	
1		2 in 1 PACKET; Type 0: Not a Combination Product		
2	NDC:47682-120-33	50 in 1 BOX	12/30/2008	
2		2 in 1 PACKET; Type 0: Not a Combination Product		
3	NDC:47682-120-47	100 in 1 BOX	12/30/2008	
3		2 in 1 PACKET; Type 0: Not a Combination Product		
4	NDC:47682-120-13	250 in 1 BOX	12/30/2008	
4	NDC:47682-120-99	2 in 1 PACKET; Type 0: Not a Combination Product		
5	NDC:47682-120-99	2 in 1 PACKET; Type 0: Not a Combination Product	12/30/2008	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/30/2008	

**Labeler** - Unifirst First Aid Corporation (832947092)

Revised: 9/2025

Unifirst First Aid Corporation