

VETRAPURE- ivermectin paste
Casa K'Love, LLC

Vetrapure (IVERMECTIN) Paste 1.87%

For Oral Use in Horses Only

Vetrapure (IVERMECTIN) Paste 1.87%

Net Wt. 0.21 oz (6.08)

Contents will treat up to 1250 lb body weight

Apple Flavored

INDICATIONS: Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. IVERMECTIN (ivermectin paste) provides effective treatment and control of the following parasites in horses. **Large Strongyles (adults)**- *Strongylus vulgaris* (also early forms in blood vessels), *S. edentatus* (also tissue stages), *S. equinus*, *Triodontophorus* spp. including *T. brevicauda* and *T. serratus* and *Craterostomum acuticaudatum*; **Small Strongyles** (adults, including those resistant to some benzimidazole class compounds)-*Coronocylus* spp. including *C. coronatus*, *C. labiatus* and *C. labratus*, *Cyathostomum* spp. including *C. catinatum* and *C. pateratum*, *Cylicocylus* spp. including *C. insigne*, *C. leptostomum*, *C. nassatus*, and *C. brevicapsulatus*, *Cylicodontophorus* spp., *Cylicostephanus* spp. including *C. calicatus*, *C. goldi*, *C. longibursatus* and *C. minutus*, and *Petrovinema poculatum*; **Small Strongyles** - Fourth-stage larvae; **Pinworms** (adults and fourth-stage larvae)-*Oxyuris equi*; **Ascarids** (adults and third- and fourth-stage larvae)- *Parascaris equorum*; **Hairworms** (adults)- *Trichostrongylus axei*; **Large-mouth Stomach Worms** (adults)- *Habronema muscae*; **Bots** (oral and gastric stages)-*Gasterophilus* spp. including *G. intestinalis* and *G. nasalis*; **Lungworms** (adults and fourth-stage larvae)- *Dictyocaulus arnfieldi*; **Intestinal Threadworms** (adults)- *Strongyloides westeri*; **Summer Sores** caused by *Habronema* and *Draschia* spp. cutaneous third-stage larvae; Dermatitis caused by neck threadworm *microfilariae*, *Onchocerca* sp.

DOSAGE: This syringe contains sufficient paste to treat one 1250 lb horse at the recommended dose rate of 91 mcg ivermectin per lb (200 mcg/kg) body weight. Each weight marking on the syringe plunger delivers enough paste to treat 250 lb body weight.

Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of parasite resistance.

CONTROL PROGRAM: All horses should be included in a regular parasite control program with particular attention being paid to mares, foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. Consult your veterinarian for a control program to meet your specific needs. IVERMECTIN (ivermectin paste) effectively controls gastrointestinal nematodes and bots of horses. Regular treatment will reduce the chances of verminous arteritis caused by *Strongylus vulgaris*.

PRODUCT ADVANTAGES: Broad-spectrum Control - IVERMECTIN kills important internal parasites, including bots and the arterial stages of *S. vulgaris*, with a single dose.

IVERMECTIN is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

ANIMAL SAFETY: IVERMECTIN (ivermectin paste) may be used in horses of all ages, including mares at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

STORAGE INFORMATION: Store at 68°F - 77°F (20°C - 25°C). Excursions between 59°F - 86°F (15°C - 30°C) are permitted.

ADMINISTRATION:

- (1) While holding plunger, turn the knurled ring on the plunger 1/4 turn to the left and slide it so the side nearest the barrel is at the prescribed weight marking, aligning the arrow on the ring to the line between the weight and lbs, as shown in the pictogram.
- (2) Lock the ring in place by making a 1/4 turn to the right. Ensure it is locked (it should no longer slide).
- (3) Make sure that the horse's mouth contains no feed.
- (4) Remove the cover from the tip of the syringe.
- (5) Insert the syringe tip into the horse's mouth at the space between the teeth.
- (6) Depress the plunger as far as it will go, depositing paste on the back of the tongue.
- (7) Immediately raise the horse's head for a few seconds after dosing.



WARNING: Do not use in horses intended for human consumption.

Not for use in humans. Keep this and all drugs out of reach of children.

Refrain from smoking and eating when handling. Wash hands after use. Avoid contact with eyes. The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report suspected adverse drug events, for technical assistance, or to obtain a copy of the SDS contact Vetrapure, at +1-909-470-6352. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or www.fda.gov/reportanimalae.

PRECAUTIONS: IVERMECTIN (ivermectin paste) has been formulated specifically for use in horses only. This product should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

OTHER WARNINGS: Parasite resistance may develop to any dewormer, and has been reported for most classes of dewormers.

Treatment with a dewormer used in conjunction with parasite management practices appropriate to the geographic area and the animal(s) to be treated may slow the development of parasite resistance.

Fecal examinations or other diagnostic tests and parasite management history should be used to determine if the product is appropriate for the herd prior to the use of any dewormer. Following the use of any dewormer, effectiveness of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method).

A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your parasite management plan should be adjusted accordingly based on regular monitoring.

Environmental Safety: Ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in an approved landfill or by incineration.

INFORMATION FOR HORSE OWNERS: Swelling and itching reactions after treatment with IVERMECTIN (ivermectin paste) have occurred in horses carrying heavy infections of neck threadworm (*Onchocerca* sp.) microfilariae. These reactions were most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advisable. Consult your veterinarian should any such reactions occur. Healing of summer sores involving extensive tissue changes may require other appropriate therapy in conjunction with treatment with IVERMECTIN. Reinfection, and measures for its prevention, should also be considered. Consult your veterinarian if the condition does not improve.

Distributed by:

Vetrapure

1150s Meadow Ln Colton CA, 92324

support@vetrapure.shop

PRINCIPAL DISPLAY PANEL - 0.21 oz (6.08g)

Keep out of reach of children

For Oral Use in Horses Only

Vetrapure

IVERMECTIN
(ivermectin paste) 1.87%

Net Wt. 0.21 oz (6.08)

NDC: 86246-100-01

Contents will treat up to 1250 lb body weight

Apple Flavored

Net Wt 0.21 oz (6.08 g)
NDC: 86246-100-01

For Oral Use in Horses Only

IVERMECTIN
(ivermectin) PASTE 1.87%

Apple Flavored!

Keep out of reach of children

Contents will treat up to 1250 lb body weight

Manufactured and Distributed by for:
Vetrapure, 1150s Meadow Ln Colton
CA, 92324
support@vetrapure.shop

LOT: VET026
EXP: Oct 2028

Apple Flavored!

IVERMECTIN
(ivermectin) PASTE 1.87%

INDICATIONS: Consult your veterinarian for assistance in the diagnosis, treatment, and control of parasitism. IVERMECTIN (ivermectin paste) provides effective treatment and control of the following protozoan in horses. **Large Strongyles** (adults): *Strongylus vulgaris* (also early forms in blood vessels), *Sedentaria* (also tissue stages), *S. equinus*, *Triodontophorus* spp. including *T. brevicauda* and *T. serratus* and *Chalarostomum aculeicaudatum*, **Small Strongyles** (adults, including those resistant to some benzimidazole class compounds) - *Coronocylus* spp. including *C. coronatus*, *C. labiatus* and *C. labialis*, *Cyathostomum* spp. including *C. colicatum* and *C. potarium*, *C. ylicocylus* spp. including *C. inligne*, *C. leptostomum*, *C. nasutus* and *C. brevicaudatus*, *Cyathostomophorus* spp., *C. ylostephanus* spp. including *C. calceatus*, *C. galei*, *C. longibursatus* and *C. minutus*, and *Petrovirenia pocolatum*; **Small Strongyles** - Fourth-stage larvae; (adults and fourth-stage larvae) - *Oxyuris equi* - **Ascarids** (adults and third- and fourth-stage larvae) - *Parascaris equorum*; **Hookworms** (adults) - *Trichostrongylus axei* **Large-mouth Stomach Worms** (adults) - *Habronema muscae*, *Gasterophilus* spp. including *G. intestinalis* and *G. nasalis*; **Lungworms** (adults and fourth-stage larvae) - *Dictyocaulus immitis*; **Intestinal Threadworms** (adults) - *Strongylides westeri*; **Summer Sores** caused by *habronema* and *dracina* spp. **cutaneous third-stage larvae**; **Demodex** caused by neck threadworm microfilariae, *Onchocerca* sp.

WARNING: Do not use in horses intended for human consumption. Not for use in humans. Keep this and all drugs out of reach of children. Refrain from smoking and eating when handling. Wash hands after use. Avoid contact with eyes. The Safety Data Sheet (SDS) contains more detailed occupational safety information. To report suspected adverse drug events, for technical assistance or to obtain a copy of the SDS, contact Vetrapure, at +1-(800)-470-0352. For additional information about adverse drug experience reporting for animal drugs contact FDA at 1-888-FDA-VETS or www.fda.gov/reportandwatch.

DOSAGE: This syringe contains sufficient paste to treat one 1250 lb horse at the recommended dose rate of 0.1 mg ivermectin per lb (200 mcg/kg) body weight. Each weight marking on the syringe plunger delivers enough paste to treat 250 lb body weight. Do not underdose. Ensure each animal receives a complete dose based on a current body weight. Underdosing may result in ineffective treatment, and encourage the development of resistance.

CONTROL PROGRAM: All horses should be included in a regular control program with particular attention being paid to mares, foals and yearlings. Foals should be treated initially at 6 to 8 weeks of age, and routine treatment repeated as appropriate. Consult your veterinarian for a control program to meet your specific needs. IVERMECTIN (ivermectin paste) effectively controls gastrointestinal nematodes. Regular treatment will reduce the chances of verminous arteritis caused by *Strongylus vulgaris*.

PRODUCT ADVANTAGES: Broad-spectrum Control - IVERMECTIN kills important internal protozoans, including the arterial stages of *S. vulgaris*, with a single dose. IVERMECTIN is a potent antiparasitic agent that is neither a benzimidazole nor an organophosphate.

ANIMAL SAFETY - IVERMECTIN (ivermectin paste) may be used in horses of all ages, including mares at any stage of pregnancy. Stallions may be treated without adversely affecting their fertility.

should not be used in other animal species as severe adverse reactions, including fatalities in dogs, may result.

OTHER WARNINGS: Treatment used in conjunction with practices appropriate to the geographic area and the animal(s) to be treated may slow the development of protozoans resistance. Fecal examinations or other diagnostic tests and management history should be used to determine if the product is appropriate for the herd, prior to the of treatment should be monitored (for example, with the use of a fecal egg count reduction test or another appropriate method). A decrease in a drug's effectiveness over time as calculated by fecal egg count reduction tests may indicate the development of resistance to the dewormer administered. Your management plan should be adjusted accordingly based on regular monitoring.

Environmental Safety: ivermectin and excreted ivermectin residues may adversely affect aquatic organisms. Do not contaminate ground or surface water. Dispose of the syringe in treatment with IVERMECTIN (ivermectin paste) have occurred in horses carrying heavy infections of neck threadworm (*Onchocerca* sp.) microfilariae. These reactions were most likely the result of microfilariae dying in large numbers. Symptomatic treatment may be advised. Consult your veterinarian should any such reactions occur. Healing of summer sores involving extensive tissue changes may require other appropriate therapeutic conjunction with treatment with IVERMECTIN. Reinfection, and measures for its

ADMINISTRATION:
(1) While holding plunger, turn the knurled ring on the plunger 1/4 turn to the left and slide it so the side nearest the barrel is at the prescribed weight marking, aligning the arrow on the ring to the line between the weight and lbs, as shown in the pictogram.
(2) Lock the ring in place by making a 1/4 turn to the right. Ensure it is locked (it should no longer slide).
(3) Make sure that the horse's mouth contains no feed.
(4) Remove the cover from the tip of the syringe.
(5) Insert the syringe tip into the horse's mouth at the space between the teeth.
(6) Depress the plunger as far as it will go, depositing paste on the back of the tongue.
(7) Immediately raise the horse's head for a few seconds after dosing.

STORAGE INFORMATION: Store at 68-77°F (20-25°C). Excursions between 59°F-86°F (15°C-30°C) are permitted.

IVERMECTIN
(ivermectin) PASTE 1.87%

VETRAPURE

ivermectin paste

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:86262-100
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IVERMECTIN (UNII: 8883YP2R6D) (IVERMECTIN - UNII:8883YP2R6D)	IVERMECTIN	18.7 mg in 1 g

Product Characteristics

Color		Score	
Shape		Size	
Flavor	APPLE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:86262-100-01	1 in 1 PACKAGE		
1		6.08 g in 1 SYRINGE, PLASTIC		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200326	10/10/2025	

Labeler - Casa K'Love, LLC (141854924)

Registrant - Casa K'Love, LLC (141854924)

Establishment

Name	Address	ID/FEI	Business Operations
Zhejiang Hisun Pharmaceutical Co., Ltd.		654211754	api manufacture

Revised: 5/2026

Casa K'Love, LLC