

ACID REDUCER- ranitidine tablet
Chain Drug Marketing Association Inc.

Drug Facts

ACTIVE INGREDIENT (IN EACH TABLET)

Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg)

PURPOSE

Acid reducer

USES

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages

WARNINGS

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

Do not use

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor

Ask a doctor before use if you have

- frequent **chest pain**
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition.
- heartburn with **lightheadedness, sweating or dizziness**
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if

- your heartburn continues or worsens
- you need to take this product for more than 14 days

If pregnant or breast-feeding,

Ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

DIRECTIONS

- adults and children 12 years and over:
 - to **relieve** symptoms, swallow 1 tablet with a glass of water
 - to **prevent** symptoms, swallow 1 tablet with a glass of water **30 to 60 minutes before** eating food or drinking beverages that cause heartburn
 - can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

OTHER INFORMATION

- **TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.**
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

INACTIVE INGREDIENTS

Colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

QUESTIONS?

Call **1-800-406-7984**

PRINCIPAL DISPLAY PANEL

QC QUALITY CHOICE®

NDC 63868-0487-50

†Compare to the active ingredient in Zantac 150®

MAXIMUM STRENGTH

HEARTBURN

150 RANITIDINE TABLETS, USP 150 mg

ACID REDUCER

Prevents & Relieves Heartburn Associated with Acid Indigestion & Sour Stomach

50 Tablets

©***DISTRIBUTED BY QUALITY CHOICE***

5093011/0112

Drug Facts

Active ingredient (in each tablet) Ranitidine 150 mg (as ranitidine hydrochloride USP, 168 mg)..... Acid reducer

Uses

- relieves heartburn associated with acid indigestion and sour stomach
- prevents heartburn associated with acid indigestion and sour stomach brought on by certain foods and beverages

Warnings

Allergy alert: Do not use if you are allergic to ranitidine or other acid reducers

- if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
- with other acid reducers
- if you have kidney disease, except under the advice and supervision of a doctor.

Do not use

NDC 63868-0487-50

QC QUALITY CHOICE

Compare to the active ingredient in Zantac 150®

MAXIMUM STRENGTH HEARTBURN 150 RANITIDINE TABLETS, USP 150 mg

ACID REDUCER

Prevents & Relieves Heartburn Associated with Acid Indigestion & Sour Stomach

50 Tablets



QC QUALITY CHOICE

MAXIMUM STRENGTH HEARTBURN 150 RANITIDINE TABLETS, USP 150 mg

50 Tablets

Directions

- adults and children 12 years and over: to relieve symptoms, swallow 1 tablet with a glass of water
- to prevent symptoms, swallow 1 tablet with a glass of water 30 to 60 minutes before eating food or drinking beverages that cause heartburn
- can be used up to twice daily (do not take more than 2 tablets in 24 hours)
- children under 12 years: ask a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF THE CARTON OR PRINTED FOIL UNDER CAP IS OPEN OR TORN.
- store at 20° - 25° C (68° - 77° F)
- avoid excessive heat or humidity
- this product is sodium and sugar free

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, iron oxide red, magnesium stearate, microcrystalline cellulose, polyethylene glycol, talc, titanium dioxide

Questions? call 1-800-406-7984

If pregnant or breast-feeding, ask a health professional before use. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222).

Read the directions, consumer information leaflet and warnings before use. Keep the carton. It contains important information.

Expiration Date

Batch No.

NON VARNISH

6 355 151 96840 0

BUY WITH CONFIDENCE

QC QUALITY CHOICE

MADE IN INDIA

0112

© DISTRIBUTED BY QUALITY CHOICE
PO Box 995
NOVI, MI 48376-0995
QC 96840
www.qualitychoice.com

This product is not manufactured or distributed by Boehringer Ingelheim Pharmaceuticals Inc., owner of the registered trademark Zantac 150®.

5093011

5093011

Drug Facts (continued)

Ask a doctor before use if you have

- frequent chest pain
- frequent wheezing, particularly with heartburn
- unexplained weight loss
- nausea or vomiting
- stomach pain
- had heartburn over 3 months. This may be a sign of a more serious condition
- heartburn with lightheadedness, sweating or dizziness
- chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Stop use and ask a doctor if ■ your heartburn continues or worsens

ACID REDUCER

ranitidine tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:63868-487
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
RANITIDINE HYDROCHLORIDE (UNII: BK76465IHM) (RANITIDINE - UNII:884KT10YB7)	RANITIDINE	150 mg

Inactive Ingredients

Ingredient Name	Strength
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOLS (UNII: 3WJQ0SDW1A)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	pink	Score	no score
Shape	OVAL	Size	12mm
Flavor		Imprint Code	9R
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:63868-487-24	24 in 1 CARTON		
1		1 in 1 BLISTER PACK		
2	NDC:63868-487-50	1 in 1 CARTON		

2	50 in 1 BOTTLE		
Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA200536	03/30/2012	

Labeler - Chain Drug Marketing Association Inc. (011920774)

Establishment			
Name	Address	ID/FEI	Business Operations
Shasun Pharmaceuticals Limited		915786829	manufacture(63868-487)

Revised: 9/2012

Chain Drug Marketing Association Inc.