

SEXUAL FUNCTION ACTIVATOR- curculigoside a granule
Sanaura Group LIMITED

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Sexual Function Activator 84611-102

Active ingredient: CURCULIGOSIDE A 50mg/ 10g

Purpose: Antioxidant

Use to:

Regulate the human immune system;

Enhance human sexual ability and desire

Warnings:

Do not use if you are allergic to this product and its ingredients.

Ask a doctor before use if you have:

Other diseases;

Or take some medicines

Ask a doctor or pharmacist before use if you are:

Children or pregnant women.

When using this product:

Take before meals;

Take it half an hour before going to bed

Stop use and ask a doctor if There was no sign of relief after taking for 4 weeks.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions:

Take one bottle orally with water, three times daily.

For internal use only.

Other information:

Store at 5-30°C;

Keep out of direct sunlight

Inactive ingredients: GANODERIOL A, SMILAGENIN, Corn Starch, YEAST .BETA.-D-GLUCAN

Sexual Function Activator

Drug Facts
Active ingredient (in each dosage unit) Purpose Curculigoside A 50mg/ 10gAntioxidant
Uses <ul style="list-style-type: none"> ■Regulate the human immune system ■Enhance human sexual ability anddesire
Warnings Do not use if you are allergic to this product and its ingredients.
Ask a doctor before use if you have <ul style="list-style-type: none"> ■Other diseases ■Or take some medicines
Ask a doctor or pharmacist before use if you are Children or pregnant women.
When using this product <ul style="list-style-type: none"> ■Take before meals ■Take it half an hour before going to bed
Stop use and ask a doctor if There was no sign of relief after taking for 4 weeks.
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children . In case of overdose, get medical help or contact a Poison Control Center right away. ►

Drug Facts (continued)
Directions <ul style="list-style-type: none"> ■Take one bottle orally with water, three times daily. ■For internal use only.
Other information <ul style="list-style-type: none"> ■Store at 5-30°C ■Keep out of direct sunlight
Inactive ingredients Ganoderiol A Smilagenin CornStarch Yeast beta-D-glucan
Questions? 123-555-1234

Label

SEXUAL FUNCTION ACTIVATOR		
curculigoside a granule		
Product Information		
Product Type	HUMAN OTC DRUG	Item Code (Source) NDC:84611-102
Route of Administration	ORAL	
Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
CURCULIGOSIDE A (UNII: A6S7X76UM5) (CURCULIGOSIDE A - UNII:A6S7X76UM5)	CURCULIGOSIDE A	50 mg in 10 g
Inactive Ingredients		
Ingredient Name	Strength	
STARCH, CORN (UNII: O8232NY3SJ)		
GANODERIOIOL A (UNII: MH91L4SN2J)		

SMILAGENIN (UNII: GQE3Q7YMVZ)

YEAST .BETA.-D-GLUCAN (UNII: 44FQ49X6UN)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:84611-102-01	10 g in 1 BAG; Type 0: Not a Combination Product	08/15/2024	
2	NDC:84611-102-02	10 g in 1 BOTTLE; Type 0: Not a Combination Product	08/15/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		08/15/2024	

Labeler - Sanaura Group LIMITED (989280112)

Establishment

Name	Address	ID/FEI	Business Operations
Sanaura Group LIMITED		989280112	manufacture(84611-102) , pack(84611-102)

Revised: 8/2024

Sanaura Group LIMITED