DERMFREE TATTOO NUMBING- lidocaine 5% tattoo numbing cream Jiangxi Hemei Pharmaceutical Co., Ltd

84010-020

Active Ingredient

Lidocaine 5%

Purpose

Local Anesthetic

Use

For temporary relief of local discomfort, itching, pain, soreness, or burning related to the tattoo process

Warnings

For external use only.

Do not use

In children less than 2 months of age on open skin wounds Do not exceed the recommended daily dosage unless directed by a doctor

When Using

Avoid contact with eyes, If this happens, rinse thoroughly with water.

Stop Use

Do not exceed the recommended daily dosage unless directed by a doctor.

Avoid contact with eyes. If this happens, rinse thoroughly with water.

If symptoms persist for more than 7 days or subside but occur again within 3 days, stop use and consult a physician.

Stop use and seek medical attention if signs of an allergy occur, such as redness, iritation, sweling, pain, or other severe symptoms. fsymptoms do not subside completely within 7 days, seek medica help.

Ask Doctor

Stop use and sek meditcal attention if signs of an alergy occur, such as

rednes, irtation, sweling, pain, or other severe symptoms. If symptoms do not subside completely within 7 days, seek medical help.

Keep Oot Of Reach Of Children

Keep out of reach of children

Directions

Before applying the numbing cream, thoroughly wash your hands and wear disposable tattoo gloves.

Clean the area of the body receiving the tattoo, Dry completely.

Apply a generous amount of numbing cream to the cleaned area and rub it inthoroughly in a circlar motion for 30 to 60 seconds, Then. cover with plastic wrap.

Leave the cream and wrap in place for 30 to 45 minutes, Wipe away the cream and thoroughly clean the area to prepare for a pain-free tattoo.

Other information

Store between 15-30°C(59-86 \square F). Avoid freezing and excessive heat above 40 °C(104°F).


Inactive ingredients

Water, Carbomer 934, propylene Glycol, soybean Lecithin, . Alpha. - Tocopherol Acetate, Benzyl Alcohol.

PRINCIPAL DISPLAY PANEL







DERMFREE[™]

TATTOO NUMBING CREAM

instructions

Purpose Local Anesthetic

For temporary relief of local discomfort, itching, pain, soreness, or burning related to the tattoo process.

related to the tattoo process.
Warnings
For external use only.
Do not use
In children less than 2 months of age on open skin wounds
In children less than 2 months of age on open skin wounds
Do not exceed the recommended daily dosage unless directed by a doctor.
Avoid contact with eyes. If this happens, rinse thoroughly with water,
If symptoms persist for more than 7 days or subside but occur again within
3 days, stop use and consult a physician.
Stop use and seek medical attention if signs of an allergy occur, such as
redness, irritation, swelling, pain, or other severe symptoms. If symptoms do
not subside completely
within 7 days, seek medical help.
Keep out of reach of children.
Divections

Directions

Neep out of reach of chindren.

Directions

Before applying the numbing cream, thoroughly wash your hands and wear disposable tattoo gloves.

Clean the area of the body receiving the tattoo. Dry completely.

Clean the area of the body receiving the tattoo. Dry completely.

Apply a generous amount of numbing cream to the cleaned area and rub it in thoroughly in a circular motion for 30 to 60 seconds. Then, cover with plastic wrap.

Leave the cream and wrap in place for 30 to 45 minutes. Wipe away the cream and thoroughly clean the area to prepare for a pain-free tattoo.

Storage and Handling

Store between 15-30 C (59-86°F). Avoid freezing and excessive heat above 40 C (104°F).

Nactive Ingredints

Water, Carbonner 334, Propylene Glycol. Soybean Lecithin, Alpha. Tocopherol Acetate, Benzyl Alcohol.

DERMFREE TATTOO NUMBING

lidocaine5% tattoo numbing cream

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:84010-020

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name		Basis of Strength	Strength
ı	LIDOCAINE (LINII: 98PI200987) (LIDOCAINE - LINII: 98PI200987)	LIDOCAINE	5 a in 100 a

Inactive Ingredients

Ingredient Name	Strength
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
SOYBEAN LECITHIN (UNII: 1DI56QDM62)	
.ALPHATOCOPHEROL ACETATE (UNII: 9E8X80D2L0)	
WATER (UNII: 059QF0KO0R)	
CARBOMER 934 (UNII: Z135WT9208)	
BENZYL ALCOHOL (UNII: LKG8494WBH)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
		20 g in 1 BOTTLE; Type 0: Not a Combination Product	08/15/2024	
M	larketing I	nformation		
M	larketing I Marketing Category	nformation Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
	Marketing	Application Number or Monograph Citation		

Labeler - Jiangxi Hemei Pharmaceutical Co., Ltd (724892056)

Establishment							
Name	Address	ID/FEI	Business Operations				
Jiangxi Hemei Pharmaceutical Co., Ltd		724892056	manufacture(84010-020)				

Revised: 8/2024 Jiangxi Hemei Pharmaceutical Co., Ltd