

DIMETAPP COLD AND COUGH AND NIGHTTIME COLD AND CONGESTION-
diphenhydramine hydrochloride, phenylephrine hydrochloride,
brompheniramine maleate, and dextromethorphan hydrobromide
Foundation Consumer Brands

Dimetapp Cold and Cough and Nighttime Cold and Congestion

Dimetapp[®]
Nighttime Cold and Congestion

Drug Facts

<i>Active ingredients (in each 10 mL)</i>	<i>Purposes</i>
Diphenhydramine HCl, 12.5 mg	Antihistamine/cough suppressant
Phenylephrine HCl, 5 mg	Nasal decongestant

Uses

- temporarily relieves:
 - nasal congestion
 - runny nose
 - cough
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily restores freer breathing through the nose

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)

- a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occurs
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: **sodium 8 mg**
- store at 20-25°C (68-77°F)
- do not refrigerate

Inactive ingredients

anhydrous citric acid, artificial flavor, FD&C blue no.1, FD&C red no. 40, glycerin, propyl

gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

Call **1-888-594-0828** weekdays 9 AM to 5 PM EST

Dimetapp® Cold and Cough

Drug Facts

Active ingredients (in each 10 mL)	Purposes
Brompheniramine maleate, 2 mg	Antihistamine
Dextromethorphan HBr, 10 mg	Cough suppressant
Phenylephrine HCl, 5 mg	Nasal decongestant

Uses

- temporarily relieves:
 - nasal congestion
 - runny nose
 - cough
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily restores freer breathing through the nose

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts such as occurs with smoking,

asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occurs
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children.In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: **sodium 4 mg**
- store at 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, artificial flavor, FD&C blue no. 1, FD&C red no. 40, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments?

Call **1-888-594-0828** weekdays 9 AM to 5 PM EST

Distributed by:

Foundation Consumer Brands, LLC, Pittsburgh, PA 15212

PRINCIPAL DISPLAY PANEL - Kit Carton

DAYTIME

CHILDREN'S

Dimetapp[®]

Cold &
Cough

BROMPHENIRAMINE MALEATE (Antihistamine)
DEXTROMETHORPHAN HBr (Cough Suppressant)
PHENYLEPHRINE HCl (Nasal Decongestant)

Relieves + comforts:

- Stuffy, runny nose
- Itchy, watery eyes
- Sneezing
- Cough

6+
YRS

Grape Flavor
Alcohol Free

1 BOTTLE

4 FL OZ (118 mL)

8 FL OZ (236 mL) TOTAL

NIGHTTIME

Nighttime
Cold &
Congestion

DIPHENHYDRAMINE HCl (Antihistamine/
Cough Suppressant)
PHENYLEPHRINE HCl (Nasal Decongestant)

Relieves + comforts:

- Stuffy, runny nose
- Itchy, watery eyes
- Sneezing
- Cough

PHARMACIST
RECOMMENDED

6+
YRS

Grape Flavor
Alcohol Free

1 BOTTLE

4 FL OZ (118 mL)

DAYTIME

NIGHTTIME

CHILDREN'S Dimetapp®

DAYTIME

NIGHTTIME

CHILDREN'S Dimetapp®

Cold & Cough

Nighttime Cold & Congestion

BROMPHENIRAMINE MALEATE (Antihistamine)
DEXTROMETHORPHAN HBr (Cough Suppressant)
PHENYLEPHRINE HCl (Nasal Decongestant)

DIPHENHYDRAMINE HCl (Antihistamine/Cough Suppressant)
PHENYLEPHRINE HCl (Nasal Decongestant)

Relieves + comforts:

- ✓ **Stuffy, runny nose**
- ✓ **Itchy, watery eyes**
- ✓ **Sneezing**
- ✓ **Cough**

Relieves + comforts:

- ✓ **Stuffy, runny nose**
- ✓ **Itchy, watery eyes**
- ✓ **Sneezing**
- ✓ **Cough**

6+
YRS



Grape Flavor
Alcohol Free

1 BOTTLE

4 FL OZ (118 mL)



8 FL OZ (236 mL) TOTAL



6+
YRS



Grape Flavor
Alcohol Free

1 BOTTLE

4 FL OZ (118 mL)

Cold & Cough (continued)

Drug Facts (continued)

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occurs
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 4 mg
- store at 20-25°C (68-77°F)

Inactive ingredients anhydrous citric acid, artificial flavor, FD&C blue no. 1, FD&C red no. 40, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments? Call 1-888-594-0828 weekdays 9 AM to 5 PM EST

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

Distributed by:
Foundation Consumer Brands, LLC, Pittsburgh, PA 15212
© 2023 dimetapp.com

B-0503-040/083-36

L60067C1

080223



No print/No varnish
Lot & Exp date

LEBG3007K3
side seam

Tamper Evident: do not use if neckband printed with "sealed for your protection" is torn or missing.

Nighttime Cold & Congestion

Drug Facts

Active ingredients (in each 10 mL)

Diphenhydramine HCl, 12.5 mg Antihistamine/
cough suppressant
Phenylephrine HCl, 5 mg Nasal decongestant

Purposes

Uses

- temporarily relieves:
 - nasal congestion
 - runny nose
 - cough
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily restores freer breathing through the nose

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other product containing diphenhydramine, even one used on skin

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

Do not take Dimetapp® Cold & Cough and Dimetapp® Nighttime Cold & Congestion at the same time. Always wait at least 4 hours before taking another dose of Dimetapp Liquid.

Nighttime Cold & Congestion (continued)

Drug Facts (continued)

When using this product

- do not use more than directed
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occurs
- symptoms do not get better within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- measure only with dosage cup provided
- keep dosage cup with product
- mL = milliliter

age	dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 8 mg
- store at 20-25°C (68-77°F) ■ do not refrigerate

Inactive ingredients anhydrous citric acid, artificial flavor, FD&C blue no. 1, FD&C red no. 40, glycerin, propyl gallate, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol solution, sucralose

Questions or comments? Call 1-888-594-0828 weekdays 9 AM to 5 PM EST

Cold & Cough

Drug Facts

Active ingredients (in each 10 mL)

Brompheniramine maleate, 2 mg Antihistamine
Dextromethorphan HBr, 10 mg Cough suppressant
Phenylephrine HCl, 5 mg Nasal decongestant

Purposes

Uses

- temporarily relieves:
 - nasal congestion
 - runny nose
 - cough
 - sneezing
 - itching of the nose or throat
 - itchy, watery eyes due to hay fever
- temporarily restores freer breathing through the nose

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product

- do not use more than directed
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

DIMETAPP COLD AND COUGH AND NIGHTTIME COLD AND CONGESTION

diphenhydramine hydrochloride, phenylephrine hydrochloride, brompheniramine maleate, and dextromethorphan hydrobromide kit

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:80070-350
---------------------	----------------	---------------------------	---------------

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:80070-350-08	1 in 1 CARTON	09/15/2021	12/31/2026

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE	118 mL
Part 2	1 BOTTLE	118 mL

Part 1 of 2

DIMETAPP NIGHTTIME COLD AND CONGESTION

diphenhydramine hydrochloride and phenylephrine hydrochloride solution

Product Information

Item Code (Source)	NDC:80070-340
---------------------------	---------------

Route of Administration	ORAL
--------------------------------	------

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	6.25 mg in 5 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	purple	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:80070-340-04	1 in 1 CARTON		
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/15/2021	12/31/2026

Part 2 of 2

DIMETAPP COLD AND COUGH

brompheniramine maleate, dextromethorphan hydrobromide, and phenylephrine hydrochloride elixir

Product Information

Item Code (Source)	NDC:80070-310
Route of Administration	ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BROMPHENIRAMINE MALEATE (UNII: IXA7C9ZN03) (BROMPHENIRAMINE - UNII:H57G17P2FN)	BROMPHENIRAMINE MALEATE	1 mg in 5 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 5 mL

PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	2.5 mg in 5 mL
---	-----------------------------	-------------------

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	purple	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:80070-310-04	1 in 1 CARTON		
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/15/2021	12/31/2026

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	09/15/2021	12/31/2026

Labeler - Foundation Consumer Brands (117603632)

