

LAXATIVE- sennosides tablet, film coated
ARMY AND AIR FORCE EXCHANGE SERVICE

Exchange Select 44-773

Active ingredient (in each tablet)

Sennosides USP, 25 mg

Purpose

Stimulant laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in 6 to 12 hours

Warnings

Do not use

laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Ask a doctor before use if you have

noticed a sudden change in bowel habits that persists over a period of 2 weeks.

Ask a doctor or pharmacist before use if you are

taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.

When using this product,

do not use for a period longer than 1 week.

Stop use and ask a doctor if

rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- **do not take more than directed**
- swallow tablet(s) with a glass of water
- swallow tablet(s) whole; do not crush, break, or chew

adults and children 12 years and over	2 tablets once or twice daily
children 6 to under 12 years	1 tablet once or twice daily
children under 6 years	ask a doctor

Other information

- **each tablet contains:** calcium 35 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, dibasic calcium phosphate dihydrate, FD&C blue #1 aluminum lake, hypromellose, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, sodium lauryl sulfate, sucrose, talc, titanium dioxide

Questions or comments?

1-800-426-9391

Principal display panel

exchange/**select**[™]

Compare To The Active Ingredient
of ex•lax® Maximum Strength*

MAXIMUM STRENGTH

laxative

sennosides USP, 25 mg
stimulant laxative

Actual Size

24 Tablets

GENTLE, DEPENDABLE

OVERNIGHT RELIEF

quality

value

*This product is not manufactured or distributed by GSK Consumer Healthcare S.A.R.L., owner of the registered trademark ex•lax® Maximum Strength. 50844 ORG082177308

“SATISFACTION GUARANTEED OR YOUR MONEY BACK”

Manufactured For Your Military Exchanges

Distributed by: LNK International, Inc.

Hauppauge, NY 11788 1-800-426-9391

**TAMPER EVIDENT: DO NOT
USE IF PACKAGE IS OPENED
OR IF BLISTER UNIT IS
TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING**

*This product is not manufactured or distributed by GSK Consumer Healthcare SARL, owner of the registered trademark ex-lax®. Maximum Strength. 50844 ORG082177308
 Manufactured For Your Military Exchanges
 Distributed by: LNK International, Inc.
 Happaugue, NY 11788 1-800-426-9391

exchange **select**

Compare To The Active Ingredient
 of ex-lax® Maximum Strength*

MAXIMUM STRENGTH

laxative

sennosides USP, 25 mg
 stimulant laxative



Actual Size

GENTLE, DEPENDABLE
 OVERNIGHT RELIEF

24 Tablets

quality
 value

TAMPER EVIDENT: DO NOT
 USE IF PACKAGE IS OPENED
 OR IF BLISTER UNIT IS
 TORN, BROKEN OR SHOWS
 ANY SIGNS OF TAMPERING



No print/No varnish Area
 Lot & Exp. Date

Drug Facts (continued)
 ■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
 ■ see end flap for expiration date and lot number
Inactive ingredients corn starch, dibasic calcium phosphate dihydrate, FD & C blue #1 aluminum lake,
 sucrose, talc, titanium dioxide
Drug Facts (continued)
 hypromellose, magnesium stearate, medium chain triglycerides, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, silicon dioxide, sodium lauryl sulfate,

Drug Facts (continued)
 ■ do not take more than directed
 ■ swallow tablet(s) with a glass of water
 ■ swallow tablet(s) whole; do not crush, break, or chew
 adults and children 2 tablets once or 12 years and over twice daily
 children 6 to under 12 years 1 tablet once or twice daily
 children under 6 years ask a doctor
Other information
 ■ each tablet contains: calcium 35 mg
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
Drug Facts (continued)
 Stop use and ask a doctor if rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
 ■ do not take more than directed
 ■ swallow tablet(s) with a glass of water
 ■ swallow tablet(s) whole; do not crush, break, or chew
 adults and children 2 tablets once or 12 years and over twice daily
 children 6 to under 12 years 1 tablet once or twice daily
 children under 6 years ask a doctor
Other information
 ■ each tablet contains: calcium 35 mg
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
Drug Facts (continued)
 Sennosides USP, 25 mg. Stimulant laxative
Active ingredient (in each tablet)
Purpose
 ■ relieves occasional constipation (irregularity)
 ■ generally produces bowel movement in 6 to 12 hours
Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.
 Ask a doctor before use if you have noticed a sudden change in bowel habits that persists over a period of 2 weeks.
 Ask a doctor or pharmacist before use if you are taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.
 When using this product, do not use for a period longer than 1 week.
Drug Facts (continued)
 ■ do not take more than directed
 ■ swallow tablet(s) with a glass of water
 ■ swallow tablet(s) whole; do not crush, break, or chew
 adults and children 2 tablets once or 12 years and over twice daily
 children 6 to under 12 years 1 tablet once or twice daily
 children under 6 years ask a doctor
Other information
 ■ each tablet contains: calcium 35 mg
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
Drug Facts (continued)
 Stop use and ask a doctor if rectal bleeding or failure to have a bowel movement occur after use of a laxative. These may be signs of a serious condition.
 If pregnant or breast-feeding, ask a health professional before use.
 Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.
Directions
 ■ do not take more than directed
 ■ swallow tablet(s) with a glass of water
 ■ swallow tablet(s) whole; do not crush, break, or chew
 adults and children 2 tablets once or 12 years and over twice daily
 children 6 to under 12 years 1 tablet once or twice daily
 children under 6 years ask a doctor
Other information
 ■ each tablet contains: calcium 35 mg
 ■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
Drug Facts (continued)
 Sennosides USP, 25 mg. Stimulant laxative
Active ingredient (in each tablet)
Purpose
 ■ relieves occasional constipation (irregularity)
 ■ generally produces bowel movement in 6 to 12 hours
Warnings
 Do not use laxative products when abdominal pain, nausea, or vomiting are present unless directed by a doctor.
 Ask a doctor before use if you have noticed a sudden change in bowel habits that persists over a period of 2 weeks.
 Ask a doctor or pharmacist before use if you are taking any other drug. Laxatives may affect how other drugs work. Take this product 2 or more hours before or after other drugs.
 When using this product, do not use for a period longer than 1 week.

B-0086E-773-08-R
 ORG082177308

Exchange Select 44-773

LAXATIVE

sennosides tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:55301-773
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
SENNOSIDES (UNII: 3FYP5M0IJX) (SENNOSIDES - UNII:3FYP5M0IJX)	SENNOSIDES	25 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
DIBASIC CALCIUM PHOSPHATE DIHYDRATE (UNII: O7TSZ97GEP)	
FD&C BLUE NO. 1 ALUMINUM LAKE (UNII: J9EQA3S2JM)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	

Product Characteristics

Color	blue	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	L7
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:55301-773-08	2 in 1 CARTON	12/29/2022	
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	12/29/2022	

Labeler - ARMY AND AIR FORCE EXCHANGE SERVICE (001695568)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	pack(55301-773)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	manufacture(55301-773) , pack(55301-773)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(55301-773)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	pack(55301-773)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		117025878	manufacture(55301-773)

Revised: 12/2023

ARMY AND AIR FORCE EXCHANGE SERVICE