

**NASAL DECONGESTANT MAXIMUM STRENGTH, NON-DROWSY- pseudoephedrine
hcl tablet, film coated
Amerisource Bergen**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Good Neighbor Pharmacy 44-112

Active ingredient (in each tablet)

Pseudoephedrine HCl 30 mg

Purpose

Nasal decongestant

Uses

- temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves sinus congestion and pressure

Warnings

Do not use

if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- thyroid disease
- diabetes
- heart disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland

When using this product

do not exceed recommended dose.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- symptoms do not improve within 7 days or occur with fever

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

adults and children 12 years and over	take 2 tablets every 4 to 6 hours; do not take more than 8 tablets in 24 hours
children ages 6 to 12 years	take 1 tablet every 4 to 6 hours; do not take more than 4 tablets in 24 hours
children under 6 years	do not use

Other information

- **each tablet contains:** calcium 15 mg
- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, silicon dioxide, titanium dioxide, triacetin

Questions or comments?

1-800-426-9391

Principal display panel

**Good
Neighbor
Pharmacy®**

**Compare to Sudafed® Sinus Congestion
active ingredient***

NDC 46122-428-62

MAXIMUM STRENGTH

**Nasal Decongestant
pseudoephedrine hydrochloride
(nasal decongestant)**

Non-Drowsy/Pseudoephedrine HCl

Relief of:

- **Nasal & Sinus Congestion**
- **Sinus Pressure**

24 Tablets | 30 mg Each

Actual Size

**TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING**

*This product is not manufactured or distributed by
Johnson & Johnson Corporation, owner of the registered
trademark Sudafed® Sinus Congestion.

50844 REV0619A11208

Distributed By
AmerisourceBergen
1300 Morris Drive
Chesterbrook, PA 19087
Questions or Concerns?
www.mygnp.com

Drug Facts
KEEP OUTER PACKAGE FOR COMPLETE PRODUCT INFORMATION

Active ingredient (in each tablet)
Pseudoephedrine HCl 30 mg..... Nasal decongestant

Uses
■ temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
■ temporarily relieves sinus congestion and pressure

Warnings
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have
■ heart disease ■ diabetes
■ thyroid disease ■ high blood pressure
■ difficulty in urination due to enlargement of the prostate gland
When using this product do not exceed recommended dosage.
■ symptoms do not improve within 7 days or occur with fever
■ nervousness, dizziness, or sleeplessness occur
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Questions or comments? 1-800-426-9391

Inactive ingredients croscarmellose sodium, dibasic calcium phosphate dihydrate, FD&C red #40 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, magnesium stearate, microcrystalline cellulose, polydextrose, polyethylene glycol, silicon dioxide, titanium dioxide, triacetin

Other information
■ each tablet contains: calcium 15 mg
■ TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
■ see end flap for expiration date and lot number

Directions
adults and children 12 years and over
take 2 tablets every 4 to 6 hours; do not take more than 8 tablets in 24 hours
children ages 6 to 11 years
take 1 tablet every 4 to 6 hours; do not take more than 4 tablets in 24 hours
children under 6 years
do not use



MAXIMUM STRENGTH
Nasal Decongestant
pseudoephedrine hydrochloride

B-0045-112-08
REV0619A11208



Compare to Sudafed® Sinus
Congestion active ingredient*

NDC 46122-428-62

MAXIMUM STRENGTH
Nasal Decongestant
pseudoephedrine hydrochloride
(nasal decongestant)

Non-Drowsy/Pseudoephedrine HCl

Relief of:
• Nasal & Sinus Congestion
• Sinus Pressure

24 Tablets | 30 mg Each

Actual Size



ABC#: 10177377



0 87701 42967 5

TAMPER EVIDENT: DO NOT USE IF
PACKAGE IS OPENED OR IF BLISTER
UNIT IS TORN, BROKEN OR SHOWS
ANY SIGNS OF TAMPERING

Distributed By
AmerisourceBergen
1300 Morris Drive
Chesterbrook, PA 19087
Questions or Concerns?
www.mygnp.com



*This product is not manufactured or distributed by
Johnson & Johnson Corporation, owner of the
registered trademark Sudafed® Sinus Congestion.
50844 REV0619A11208

GNP 44-112

NASAL DECONGESTANT MAXIMUM STRENGTH, NON-DROWSY

pseudoephedrine hcl tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:46 122-428	
Route of Administration	ORAL			
Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)		PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg	
Inactive Ingredients				
Ingredient Name			Strength	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)				
ANHYDROUS DIBASIC CALCIUM PHOSPHATE (UNII: L11K75P92J)				
FD&C RED NO. 40 (UNII: WZB9127XOA)				
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)				
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)				
MAGNESIUM STEARATE (UNII: 70097M6I30)				
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POLYDEXTROSE (UNII: VH2XOU12IE)				
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
TRIACETIN (UNII: XHX3C3X673)				
Product Characteristics				
Color	RED	Score	no score	
Shape	ROUND	Size	7mm	
Flavor		Imprint Code	44;112	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:46 122-428-62	1 in 1 CARTON	08/25/1981	
1		24 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH FINAL		part341	08/25/1981	

Labeler - Amerisource Bergen (007914906)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	MANUFACTURE(46122-428)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867837	PACK(46122-428)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	PACK(46122-428)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		038154464	PACK(46122-428)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		868734088	PACK(46122-428)