

**DAYTIME AND NIGHTTIME CHERRY COLD AND FLU- daytime - acetaminophen, dextromethorphan hbr, phenylephrine hcl nighttime - acetaminophen, dextromethorphan hbr, doxylamine succinate  
BETTER LIVING BRANDS, LLC**

-----  
**773L-Albertsons- Daytime and Nighttime Cold & Flu Liquid**

***Nighttime Cold & Flu***

***Drug Facts***

***Active ingredients (in each 30 mL)***

Acetaminophen 650 mg  
Dextromethorphan HBr 30 mg  
Doxylamine succinate 12.5 mg

***Purpose***

Pain reliever/ fever reducer  
Cough suppressant  
Antihistamine

***Uses***

temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

***Warnings***

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

■ more than 4 doses in 24 hours, which is the maximum daily amount for this product.

■ with other drugs containing acetaminophen

■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include:

■ skin reddening

■ blisters

■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more

than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

### **Allergy alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

### **Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

■ cough comes back or occurs with rash or headache that lasts.  
These could be signs of a serious condition.

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

---

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

---

### **Other information**

- **each 30 mL tablespoon contains:**sodium 52 mg
- store at no greater than 25°C and do not refrigerate

### **Inactive ingredients**

alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

### **Questions or comments?**

Call **1-877-290-4008**

### **Daytime Cold & Flu Drug Facts**

#### **Active ingredients (in each 15 mL)**

Acetaminophen 325 mg  
Dextromethorphan HBr 10 mg  
Phenylephrine HCl 5 mg

#### **Purpose**

Pain reliever/Fever reducer  
Cough suppressant

## Nasal decongestant

### **Uses**

temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

### **Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

■ adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product

■ child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product

■ taken with other drugs containing acetaminophen

■ adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include:

■ skin reddening

■ blisters

■ rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Liver warning:**

This product contains acetaminophen. Severe liver damage may occur if

- adults take more than 4 doses (30 mL each) in 24 hours, which is the maximum daily amount for this product
- child take more than 4 doses (15 mL each) in 24 hours, which is the maximum daily amount for this product
- with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

### **Allergy alert:**

Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Sore throat warning:**

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin.

**When using this product,**

**do not use more than directed.**

**Stop use and ask a doctor if**

- You get nervous, dizzy, or sleepless.
- Pain, nasal congestion or cough get worse or last more than 5 days (children) or 7 days (adults)
- Fever gets worse or last more than 3 days
- Redness or swelling is present
- New symptoms occur
- Cough comes back, or occurs with rash or headache that lasts.

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

## Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

---

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

---

### Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

**Ask a doctor or pharmacist before use if you are** taking the blood thinning drug warfarin.

### When using this product, do not use more than directed.

#### Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough get worse or last more than 5 days (children) and 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Other information***

- **each 15 mL contains:** sodium 56 mg
- store at no greater than 25°C and do not refrigerate.

***Inactive ingredients***

anhydrous citric acid, FD&C yellow no. 6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

***Questions or comments?***

Call **1-877-290-4008**

**PRINCIPAL DISPLAY PANEL - Convenience Pack**

**COMPARE TO**  
Vicks® DayQuil™ Cold &  
Flu Liquid Active Ingredients\*

NDC 21130-947-01



**DAYTIME  
COLD & FLU  
RELIEF**

**FOR AGES 6  
& OVER**

**ORIGINAL  
FLAVOR**

**ACETAMINOPHEN** 325 mg

- Pain Reliever/Fever Reducer

**DEXTROMETHORPHAN HBr** 10 mg

- Cough Suppressant

**PHENYLEPHRINE HCl** 5 mg

- Nasal Decongestant

• Relieves headache, fever, sore throat,  
minor aches and pains

• Cough

• Nasal congestion, sinus pressure

ALCOHOL FREE

12 FL OZ (354 mL)

**FRONT**

TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND SEAL AROUND THE NECK IS BROKEN OR MISSING

← PEEL BACK FOR DRUG FACTS

**Drug Facts**

Active Ingredients (In each 15 mL)	Purpose
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant

**Uses** temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- headache
- minor aches & pains
- fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4 doses (30 mL each) in 24 hrs, which is

DISTRIBUTED BY:  
BETTER LIVING BRANDS LLC  
P.O. BOX 99  
PLEASANTON, CA 94566-0099  
T: 985-723-3329  
772L 0425 Made in India  
RD 25099 S6850



smartlabel™  
Scan here for more information



**PARENTS:**  
Learn about teen medicine abuse  
www.StopMedicineAbuse.org

**NOT FOR RESALE**

LOT:  
EXP.:

Print/Varnish Omit Area

BACK

**Drug Facts (continued)**

the maximum daily amount for this product

- child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away. **Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- sodium-restricted diet

**Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.**

BACK PANEL 1

**Drug Facts (continued)**

**When using this product, do not use more than directed.**

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough get worse or last more than 5 days (children) and 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back, or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
  - only use the dose cup provided
  - do not exceed 4 doses per 24 hrs
- |                                 |                   |
|---------------------------------|-------------------|
| adults & children 12 yrs & over | 30 mL every 4 hrs |
| children 6 to under 12 yrs      | 15 mL every 4 hrs |
| children 4 to under 6 yrs       | ask a doctor      |
| children under 4 yrs            | do not use        |

**Other information**

- each 15 mL contains: sodium 56 mg
- store at no greater than 25°C and do not refrigerate

**Inactive Ingredients** anhydrous citric acid, FD&C yellow no. 6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium chloride, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

**Questions or comments?** Call 1-877-290-4008

\*This product is not manufactured or distributed by Procter & Gamble Company, owner of the registered trademarks Vicks® DayQuil™ Cold & Flu Liquid.

BACK PANEL 2

GLUE AREA

**COMPARE TO**

Vicks® NyQuil™ Cold & Flu  
Cherry Liquid Active Ingredients\*

NDC 21130-907-01

*Signature*  
SELECT

**NIGHTTIME  
COLD & FLU  
RELIEF**

**FOR AGES 12  
& OVER**

**CHERRY  
FLAVOR**

**ACETAMINOPHEN 650 mg**

- Pain Reliever/Fever Reducer

**DEXTROMETHORPHAN HBr 30 mg**

- Cough Suppressant

**DOXYLAMINE SUCCINATE 12.5 mg**

- Antihistamine

• Relieves headaches, fever, sore  
throat, minor aches & pains

• Sneezing, runny nose, cough

• Nighttime relief



**ALCOHOL 10%**

**12 FL OZ (354 mL)**

**FRONT**

**TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND SEAL AROUND THE NECK IS BROKEN OR MISSING**

**← PEEL BACK FOR DRUG FACTS**

**Drug Facts**

**Active Ingredients (in each 30 mL)** **Purpose**

Acetaminophen 650 mg.....Pain reliever/fever reducer

Dextromethorphan HBr 30 mg.....Cough suppressant

Doxylamine succinate 12.5 mg.....Antihistamine

**Uses** temporarily relieves common cold/flu symptoms:

- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever
- runny nose and sneezing

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4 doses in 24 hours, which is the maximum daily amount

← for this product

DISTRIBUTED BY:  
BETTER LIVING BRANDS LLC  
P.O. BOX 99  
PLEASANTON, CA 94566-0099  
1-888-723-3929

750L 0425 Made in India  
RD 25099 S6850



smartlabel™  
Scan here for more information

**PARENTS:**  
Learn about teen medicine abuse  
www.StopMedicineAbuse.org

**NOT FOR RESALE**

LOT: \_\_\_\_\_  
EXP: \_\_\_\_\_ Print/Varnish Omit Area

**Drug Facts (continued)**

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin redness
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or is caused with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

**Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

**When using this product**

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

**Drug Facts (continued)**

- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- each 30 mL contains: sodium 52 mg
- store at no greater than 25°C and do not refrigerate

**Inactive ingredients** alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum

**Questions or comments?** Call 1-877-290-4008

\*This product is not manufactured or distributed by Procter & Gamble Company, owner of the registered trademarks Vicks® NyQuil™ Cold & Flu Cherry Liquid.

**TWIN PACK**

**COMPARE TO Vicks® DayQuil™ Cold & Flu Liquid Active Ingredients\*** **COMPARE TO Vicks® NyQuil™ Cold & Flu Liquid Active Ingredients\***

NDC 21130-967-02

**DAYTIME**

**COLD & FLU RELIEF**

ALCOHOL FREE

Relieves headache, fever, sore throat, minor aches and pains, cough, nasal congestion, sinus pressure

**FOR AGES 6 & OVER** **ORIGINAL FLAVOR**

ACETAMINOPHEN 325 mg  
- Pain Reliever/Fever Reducer

DEXTROMETHORPHAN HBr 10 mg  
- Cough Suppressant

PHENYLEPHRINE HCl 5 mg  
- Nasal Decongestant

**NIGHTTIME**

**COLD & FLU RELIEF**

ALCOHOL 10% | NIGHTTIME RELIEF

Relieves headache, fever, sore throat, minor aches & pains, sneezing, runny nose, cough

**FOR AGES 12 & OVER** **CHERRY FLAVOR**

ACETAMINOPHEN 650 mg  
- Pain Reliever/Fever Reducer

DEXTROMETHORPHAN HBr 30 mg  
- Cough Suppressant

DOXYLAMINE SUCCINATE 12.5 mg  
- Antihistamine

When using Daytime and Nighttime products, carefully read the labeling to ensure correct dosing.

**2 BOTTLES - 12 FL OZ (354 mL) EACH 24 FL OZ (708 mL) TOTAL**

NIGHTTIME COLD & FLU	DAYTIME COLD & FLU
<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each 30 mL)</b></p> <p>Acetaminophen 650 mg.....Pain reliever/fever reducer</p> <p>Dextromethorphan HBr 30 mg.....Cough suppressant</p> <p>Phenylephrine HCl 5 mg.....Nasal decongestant</p> <p><b>Uses</b> temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ minor aches and pains ■ fever ■ sore throat ■ headache ■ minor aches &amp; pains ■ cough</p> <p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses in 24 hours, which is the maximum daily amount for this product. ■ child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product. ■ takes with other drugs containing acetaminophen. ■ adult has 3 or more alcoholic drinks every day while using this product. ■ allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash ■ if a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p><b>Do not use</b> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p><b>Ask a doctor before use if you have</b> liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or is caused with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to enlarged prostate gland</p> <p><b>Ask a doctor or pharmacist before use if you are</b> taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin</p> <p><b>When using this product</b> ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ do not use if you are taking the blood thinning drug warfarin</p> <p><b>Directions</b> ■ take only as directed ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs</p> <p>adults &amp; children 12 yrs &amp; over 30 mL every 6 hrs</p> <p>children 4 to under 12 yrs ask a doctor</p> <p>children under 4 yrs do not use</p> <p><b>Other information</b> ■ each 30 mL contains: sodium 52 mg ■ store at no greater than 25°C and do not refrigerate</p> <p><b>Inactive ingredients</b> alcohol, anhydrous citric acid, FD&amp;C blue no. 1, FD&amp;C red no. 40, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum</p> <p><b>Questions or comments?</b> Call 1-877-290-4008</p>	<p><b>Drug Facts</b></p> <p><b>Active ingredients (in each 15 mL)</b></p> <p>Acetaminophen 325 mg.....Pain reliever/fever reducer</p> <p>Dextromethorphan HBr 10 mg.....Cough suppressant</p> <p>Phenylephrine HCl 5 mg.....Nasal decongestant</p> <p><b>Uses</b> temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ minor aches and pains ■ fever ■ sore throat ■ headache ■ minor aches &amp; pains ■ cough</p> <p><b>Warnings</b></p> <p><b>Liver warning:</b> This product contains acetaminophen. Severe liver damage may occur if you take more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product. ■ child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product. ■ takes with other drugs containing acetaminophen. ■ adult has 3 or more alcoholic drinks every day while using this product. ■ allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin redness ■ blisters ■ rash ■ if a skin reaction occurs, stop use and seek medical help right away.</p> <p><b>Sore throat warning:</b> If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p><b>Do not use</b> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p> <p><b>Ask a doctor before use if you have</b> liver disease ■ glaucoma ■ cough that occurs with too much phlegm (mucus) ■ a breathing problem or chronic cough that lasts or is caused with smoking, asthma, chronic bronchitis, or emphysema ■ trouble urinating due to enlarged prostate gland</p> <p><b>Ask a doctor or pharmacist before use if you are</b> taking sedatives or tranquilizers ■ taking the blood thinning drug warfarin</p> <p><b>When using this product</b> ■ do not use more than directed</p> <p><b>Directions</b> ■ take only as directed ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs</p> <p>adults &amp; children 12 yrs &amp; over 15 mL every 6 hrs</p> <p>children 4 to under 12 yrs ask a doctor</p> <p>children under 4 yrs do not use</p> <p><b>Other information</b> ■ each 15 mL contains: sodium 50 mg ■ store at no greater than 25°C and do not refrigerate</p> <p><b>Inactive ingredients</b> alcohol, anhydrous citric acid, FD&amp;C yellow no. 6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate, sodium citrate, sodium saccharin, sorbitol solution, sucralose, xanthan gum</p> <p><b>Questions or comments?</b> Call 1-877-290-4008</p>

**TAMPER EVIDENT: DO NOT USE IF PRINTED SHRINKBAND SEAL AROUND THE NECK IS BROKEN OR MISSING**

DISTRIBUTED BY:  
BETTER LIVING BRANDS LLC  
P.O. BOX 99  
PLEASANTON, CA 94566-0099  
1-888-723-3929

774 0425

**PARENTS:**  
Learn about teen medicine abuse  
www.StopMedicineAbuse.org

LOT: \_\_\_\_\_  
EXP: \_\_\_\_\_ Print/Varnish Omit Area

# DAYTIME AND NIGHTTIME CHERRY COLD AND FLU

daytime - acetaminophen, dextromethorphan hbr, phenylephrine hcl nighttime - acetaminophen, dextromethorphan hbr, doxylamine succinate kit

Product Information				
Product Type	HUMAN OTC DRUG			
Item Code (Source)	NDC:21130-967			
<b>Packaging</b>				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:21130-967-	1 in 1 BOTTLE; Type 0: Not a Combination	03/01/2025	

**Quantity of Parts**

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE, PLASTIC	354 mL
Part 2	1 BOTTLE, PLASTIC	354 mL

**Part 1 of 2****DAYTIME COLD AND FLU**

acetaminophen, dextromethorphan hydrobromide, phenylephrine hcl liquid

**Product Information**

Item Code (Source)	NDC:21130-947
Route of Administration	ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

**Inactive Ingredients**

Ingredient Name	Strength
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

**Product Characteristics**

<b>Color</b>	orange	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	APRICOT	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/01/2025	

### Part 2 of 2

### NIGHTTIME CHERRY COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, and doxylamine succinate liquid

### Product Information

<b>Item Code (Source)</b>	NDC:21130-907
<b>Route of Administration</b>	ORAL

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	

<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>SORBITOL SOLUTION</b> (UNII: 8KW3E207O2)	

### Product Characteristics

<b>Color</b>	red	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		354 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/01/2025	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/01/2025	

**Labeler** - BETTER LIVING BRANDS, LLC (009137209)

**Registrant** - TIME CAP LABORATORIES, INC. (037052099)

### Establishment

Name	Address	ID/FEI	Business Operations
MARKSANS PHARMA LIMITED		677604129	manufacture(21130-967)