

OXALIPLATIN - oxaliplatin injection, solution, concentrate
Sun Pharmaceutical Industries, Inc.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use OXALIPLATIN INJECTION safely and effectively. See full prescribing information for OXALIPLATIN INJECTION.

OXALIPLATIN injection,
for intravenous use

Initial U.S. Approval: 2002

WARNING: ANAPHYLACTIC REACTIONS
See full prescribing information for complete boxed warning.

Anaphylactic reactions to oxaliplatin injection have been reported, and may occur within minutes of oxaliplatin injection administration. Epinephrine, corticosteroids, and antihistamines have been employed to alleviate symptoms. (5.1)

RECENT MAJOR CHANGES

Dosage and Administration (2.2) 10/2015

Warnings and Precautions (5.3, 5.4, 5.7, 5.8, 5.9) 10/2015

INDICATIONS AND USAGE

Oxaliplatin injection is a platinum-based drug used in combination with infusional 5-fluorouracil/leucovorin, which is indicated for:

- adjuvant treatment of stage III colon cancer in patients who have undergone complete resection of the primary tumor. (1)
- treatment of advanced colorectal cancer. (1)

DOSAGE AND ADMINISTRATION

- Administer oxaliplatin injection in combination with 5-fluorouracil/leucovorin every 2 weeks. (2.1):
 - **Day 1:** Oxaliplatin injection 85 mg/m² intravenous infusion in 250 to 500 mL 5% Dextrose Injection, USP and leucovorin 200 mg/m² intravenous infusion in 5% Dextrose Injection, USP both given over 120 minutes at the same time in separate bags using a Y-line, followed by 5-fluorouracil 400 mg/m² intravenous bolus given over 2 to 4 minutes, followed by 5-fluorouracil 600 mg/m² intravenous infusion in 500 mL 5% Dextrose Injection, USP (recommended) as a 22-hour continuous infusion.
 - **Day 2:** leucovorin 200 mg/m² intravenous infusion over 120 minutes, followed by 5-fluorouracil 400 mg/m² intravenous bolus given over 2 to 4 minutes, followed by 5-fluorouracil 600 mg/m² intravenous infusion in 500 mL 5% Dextrose Injection, USP (recommended) as a 22-hour continuous infusion.
- Reduce the dose of oxaliplatin injection to 75 mg/m² (adjuvant setting) or 65 mg/m² (advanced colorectal cancer) (2.2):
 - if there are persistent grade 2 neurosensory events that do not resolve.
 - after recovery from grade 3/4 gastrointestinal toxicities (despite prophylactic treatment) or grade 4 neutropenia or febrile neutropenia or grade 3/4 thrombocytopenia. Delay next dose until neutrophils $\geq 1.5 \times 10^9/L$ and platelets $\geq 75 \times 10^9/L$.
- For patients with severe renal impairment (creatinine clearance < 30 mL/min), the initial recommended dose is 65 mg/m². (2.2)
- Discontinue oxaliplatin injection if there are persistent Grade 3 neurosensory events. (2.2)
- Never prepare a final dilution with a sodium chloride solution or other chloride-containing solutions. (2.3)

DOSAGE FORMS AND STRENGTHS

Single-use vials of 50 mg or 100 mg oxaliplatin as a sterile, preservative-free, aqueous solution at a concentration of 5 mg/mL. (3)

CONTRAINDICATIONS

- Known allergy to oxaliplatin injection or other platinum compounds. (4, 5.1)

WARNINGS AND PRECAUTIONS

- Allergic Reactions: Monitor for development of rash, urticaria, erythema, pruritis, bronchospasm, and hypotension. (5.1)
- Neuropathy: Reduce the dose or discontinue oxaliplatin injection if necessary. (5.2)
- Severe Neutropenia: Delay oxaliplatin injection until neutrophils are $\geq 1.5 \times 10^9/L$. Withhold oxaliplatin injection for sepsis. (5.3)
- Pulmonary Toxicity: May need to discontinue oxaliplatin injection until interstitial lung disease or pulmonary fibrosis are excluded. (5.4)
- Hepatotoxicity: Monitor liver function tests. (5.5)
- Cardiovascular Toxicity: Correct hypokalemia or hypomagnesemia prior to initiating oxaliplatin injection. (5.6)
- Rhabdomyolysis: Discontinue oxaliplatin injection if rhabdomyolysis occurs. (5.7)
- Pregnancy. Fetal harm can occur when administered to a pregnant woman. Women should be apprised of the potential harm to the fetus. (5.8, 8.1)

ADVERSE REACTIONS

Most common adverse reactions (incidence $\geq 40\%$) were peripheral sensory neuropathy, neutropenia, thrombocytopenia, anemia, nausea, increase in transaminases and alkaline phosphatase, diarrhea, emesis, fatigue and stomatitis. Other adverse reactions, including serious adverse reactions, have been reported. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact Sun Pharmaceutical Industries, Inc. 1-800-818-4555 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

See 17 for PATIENT COUNSELING INFORMATION.

Revised: 10/2018

FULL PRESCRIBING INFORMATION: CONTENTS*

WARNING: ANAPHYLACTIC REACTIONS

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Dosage
- 2.2 Dose Modification Recommendations
- 2.3 Preparation of Infusion Solution

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Allergic Reactions
- 5.2 Neurologic Toxicity
- 5.3 Severe Neutropenia
- 5.4 Pulmonary Toxicity
- 5.5 Hepatotoxicity
- 5.6 Cardiovascular Toxicity
- 5.7 Rhabdomyolysis
- 5.8 Use in Pregnancy
- 5.9 Recommended Laboratory Tests

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

7 DRUG INTERACTIONS

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.3 Nursing Mothers
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Patients with Renal Impairment

10 OVERDOSAGE

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

- 14.1 Combination Adjuvant Therapy with Oxaliplatin Injection and Infusional 5-fluorouracil/leucovorin in Patients with Colon Cancer
- 14.2 Combination Therapy with Oxaliplatin Injection and 5-fluorouracil/leucovorin in Patients Previously Untreated for Advanced Colorectal Cancer
- 14.3 Combination Therapy with Oxaliplatin Injection and 5-fluorouracil/leucovorin in Previously Treated Patients with Advanced Colorectal Cancer

15 REFERENCES

16 HOW SUPPLIED/STORAGE AND HANDLING

- 16.1 How Supplied
- 16.2 Storage
- 16.3 Handling and Disposal

17 PATIENT COUNSELING INFORMATION

- FDA-Approved Patient Labeling

* Sections or subsections omitted from the full prescribing information are not listed.

FULL PRESCRIBING INFORMATION

WARNING: ANAPHYLACTIC REACTIONS

Anaphylactic reactions to oxaliplatin injection have been reported, and may occur within minutes of oxaliplatin injection administration. Epinephrine, corticosteroids, and antihistamines have been employed to alleviate symptoms of anaphylaxis [see Warnings and Precautions (5.1)].

1 INDICATIONS AND USAGE

Oxaliplatin injection, USP, used in combination with infusional 5-fluorouracil/leucovorin, is indicated for:

- adjuvant treatment of stage III colon cancer in patients who have undergone complete resection of the primary tumor.
- treatment of advanced colorectal cancer.

2 DOSAGE AND ADMINISTRATION

Oxaliplatin injection should be administered under the supervision of a qualified physician experienced in the use of cancer chemotherapeutic agents. Appropriate management of therapy and complications is possible only when adequate diagnostic and treatment facilities are readily available.


2.1 Dosage

Administer oxaliplatin injection in combination with 5-fluorouracil/leucovorin every 2 weeks. For advanced disease, treatment is recommended until disease progression or unacceptable toxicity. For adjuvant use, treatment is recommended for a total of 6 months (12 cycles):

Day 1: Oxaliplatin injection 85 mg/m² intravenous infusion in 250 to 500 mL 5% Dextrose injection, USP and leucovorin 200 mg/m² intravenous infusion in 5% Dextrose Injection, USP both given over 120 minutes at the same time in separate bags using a Y-line, followed by 5-fluorouracil 400 mg/m² intravenous bolus given over 2 to 4 minutes, followed by 5-fluorouracil 600 mg/m² intravenous infusion in 500 mL 5% Dextrose Injection, USP (recommended) as a 22-hour continuous infusion.

Day 2: Leucovorin 200 mg/m² intravenous infusion over 120 minutes, followed by 5-fluorouracil 400 mg/m² intravenous bolus given over 2 to 4 minutes, followed by 5-fluorouracil 600 mg/m² intravenous infusion in 500 mL 5% Dextrose Injection, USP (recommended) as a 22-hour continuous infusion.

Figure 1



The administration of oxaliplatin injection does not require prehydration. Premedication with antiemetics, including 5-HT₃ blockers with or without dexamethasone, is recommended.

For information on 5-fluorouracil and leucovorin, see the respective package inserts.

2.2 Dose Modification Recommendations

Prior to subsequent therapy cycles, patients should be evaluated for clinical toxicities and recommended laboratory tests [see Warnings and Precautions (5.11)]. Prolongation of infusion time for oxaliplatin injection from 2 hours to 6 hours may mitigate acute toxicities. The infusion times for 5-fluorouracil and leucovorin do not need to be changed.

Adjuvant Therapy in Patients with Stage III Colon Cancer

Neuropathy and other toxicities were graded using the NCI CTC scale version 1 [see Warnings and Precautions (5.2)].

For patients who experience persistent Grade 2 neurosensory events that do not resolve, a dose reduction of oxaliplatin injection to 75 mg/m² should be considered. For patients with persistent Grade 3 neurosensory events, discontinuing therapy should be considered. The infusional 5-fluorouracil/leucovorin regimen need not be altered.

A dose reduction of oxaliplatin injection to 75 mg/m² and infusional 5-fluorouracil to 300 mg/m² bolus and 500 mg/m² 22 hour infusion is recommended for patients after recovery from grade 3/4 gastrointestinal (despite prophylactic treatment) or grade 4 neutropenia, or febrile neutropenia, or grade 3/4 thrombocytopenia. The next dose should be delayed until: neutrophils $\geq 1.5 \times 10^9/L$ and platelets $\geq 75 \times 10^9/L$.

Dose Modifications in Therapy in Previously Untreated and Previously Treated Patients with Advanced Colorectal Cancer

Neuropathy was graded using a study-specific neurotoxicity scale [see Warnings and Precautions (5.2)]. Other toxicities were graded by the NCI CTC, Version 2.

For patients who experience persistent Grade 2 neurosensory events that do not resolve, a dose reduction of oxaliplatin injection to 65 mg/m² should be considered. For patients with persistent Grade 3 neurosensory events, discontinuing therapy should be considered. The 5-fluorouracil/leucovorin regimen need not be altered.

A dose reduction of oxaliplatin injection to 65 mg/m² and 5-fluorouracil by 20% (300 mg/m² bolus and 500 mg/m² 22-hour infusion) is recommended for patients after recovery from grade 3/4 gastrointestinal (despite prophylactic treatment) or grade 4 neutropenia, or febrile neutropenia, or grade 3/4 thrombocytopenia. The next dose should be delayed until: neutrophils $\geq 1.5 \times 10^9/L$ and platelets $\geq 75 \times 10^9/L$.

Dose Modifications in Therapy for Patients with Renal Impairment

In patients with normal renal function or mild to moderate renal impairment, the recommended dose of oxaliplatin injection is 85 mg/m². In patients with severe renal impairment, the initial recommended oxaliplatin injection dose should be reduced to 65 mg/m² [see *Use in Specific Populations (8.6)* and *Clinical Pharmacology (12.3)*].

2.3 Preparation of Infusion Solution

Do not freeze and protect from light the concentrated solution.

A final dilution must never be performed with a sodium chloride solution or other chloride-containing solutions.

The solution must be further diluted in an infusion solution of 250 to 500 mL of 5% Dextrose Injection, USP. After dilution with 250 to 500 mL of 5% Dextrose Injection, USP, the shelf life is **6 hours at room temperature [20° to 25°C (68° to 77°F)] or up to 24 hours under refrigeration [2° to 8°C (36° to 46°F)]**.

After final dilution, protection from light is not required.

Oxaliplatin injection is incompatible in solution with alkaline medications or media (such as basic solutions of 5-fluorouracil) and must not be mixed with these or administered simultaneously through the same infusion line. **The infusion line should be flushed with 5% Dextrose Injection, USP prior to administration of any concomitant medication.**

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration and discarded if present.

Needles or intravenous administration sets containing aluminum parts that may come in contact with oxaliplatin injection should not be used for the preparation or mixing of the drug. Aluminum has been reported to cause degradation of platinum compounds.

3 DOSAGE FORMS AND STRENGTHS

Oxaliplatin injection is supplied in single-use vials containing 50 mg or 100 mg of oxaliplatin, USP as a sterile, preservative-free, aqueous solution at a concentration of 5 mg/mL.

4 CONTRAINDICATIONS

Oxaliplatin injection should not be administered to patients with a history of known allergy to oxaliplatin injection or other platinum compounds [see *Warnings and Precautions (5.1)*].

5 WARNINGS AND PRECAUTIONS

5.1 Allergic Reactions

See boxed warning

Grade 3/4 hypersensitivity, including anaphylactic/anaphylactoid reactions, to oxaliplatin injection has been observed in 2 to 3% of colon cancer patients. These allergic reactions which can be fatal, can occur within minutes of administration and at any cycle, and were similar in nature and severity to those reported with other platinum-containing compounds, such as rash, urticaria, erythema, pruritus, and, rarely, bronchospasm and hypotension. The symptoms associated with hypersensitivity reactions reported in the previously untreated patients were urticaria, pruritus, flushing of the face, diarrhea associated with oxaliplatin infusion, shortness of breath, bronchospasm, diaphoresis, chest pains, hypotension, disorientation and syncope. These reactions are usually managed with standard epinephrine, corticosteroid, antihistamine therapy, and require discontinuation of therapy. Rechallenge is contraindicated in these patients [see *Contraindications (4)*]. Drug-related deaths associated with platinum compounds from anaphylaxis have been reported.

5.2 Neurologic Toxicity

Neuropathy

Oxaliplatin injection is associated with two types of neuropathy:

An acute, reversible, primarily peripheral, sensory neuropathy that is of early onset, occurring within hours or one to two days of dosing, that resolves within 14 days, and that frequently recurs with further dosing. The symptoms may be precipitated or exacerbated by exposure to cold temperature or cold objects and they usually present as transient paresthesia, dysesthesia and hypoesthesia in the hands, feet, perioral area, or throat. Jaw spasm, abnormal tongue sensation, dysarthria, eye pain, and a feeling of chest pressure have also been observed. The acute, reversible pattern of sensory neuropathy was observed in about 56% of study patients who received oxaliplatin injection with 5-fluorouracil/leucovorin. In any individual cycle acute neurotoxicity was observed in approximately 30% of patients. In adjuvant patients the median cycle of onset for grade 3 peripheral sensory neuropathy was 9 in the previously treated patients the median number of cycles administered on the oxaliplatin injection with 5-fluorouracil/leucovorin combination arm was 6.

An acute syndrome of pharyngolaryngeal dysesthesia seen in 1 to 2% (grade 3/4) of patients previously untreated for advanced colorectal cancer, and the previously treated patients, is characterized by subjective sensations of dysphagia or dyspnea, without any laryngospasm or bronchospasm (no stridor or wheezing). Ice (mucositis prophylaxis) should be avoided during the infusion of oxaliplatin injection because cold temperature can exacerbate acute neurological symptoms.

A persistent (>14 days), primarily peripheral, sensory neuropathy that is usually characterized by paresthesias, dysesthesias, hypoesthesias, but may also include deficits in proprioception that can interfere with daily activities (e.g., writing, buttoning, swallowing, and difficulty walking from impaired proprioception). These forms of neuropathy occurred in 48% of the study patients receiving oxaliplatin injection with 5-fluorouracil/leucovorin. Persistent neuropathy can occur without any prior acute neuropathy event. The majority of the patients (80%) who developed grade 3 persistent neuropathy progressed from prior Grade 1 or 2 events. These symptoms may improve in some patients upon discontinuation of oxaliplatin injection.

In the adjuvant colon cancer trial, neuropathy was graded using a prelisted module derived from the Neuro-Sensory section of the National Cancer Institute Common Toxicity Criteria (NCI CTC) scale, Version 1, as follows:

Table 1 - NCI CTC Grading for Neuropathy in Adjuvant Patients

| Grade | Definition |
|---------|--|
| Grade 0 | No change or none |
| Grade 1 | Mild paresthesias, loss of deep tendon reflexes |
| Grade 2 | Mild or moderate objective sensory loss, moderate paresthesias |
| Grade 3 | Severe objective sensory loss or paresthesias that interfere with function |
| Grade 4 | Not applicable |

Peripheral sensory neuropathy was reported in adjuvant patients treated with the oxaliplatin injection combination with a frequency of 92% (all grades) and 13% (grade 3). At the 28-day follow-up after the last treatment cycle, 60% of all patients had any grade (Grade 1=40%, Grade 2=16%, Grade 3=5%) peripheral sensory neuropathy decreasing to 39% at 6 months follow-up (Grade 1=31%, Grade 2=7%, Grade 3=1%) and 21% at 18 months of follow-up (Grade 1=17%, Grade 2=3%, Grade 3=1%). In the advanced colorectal cancer studies, neuropathy was graded using a study-specific neurotoxicity scale, which was different from the NCI CTC scale, Version 2 (see below).

Table 2 - Grading Scale for Paresthesias/Dysesthesias in Advanced Colorectal Cancer Patients

| Grade | Definition |
|---------|---|
| Grade 1 | Resolved and did not interfere with functioning |
| Grade 2 | Interfered with function but not daily activities |
| Grade 3 | Pain or functional impairment that interfered with daily activities |
| Grade 4 | Persistent impairment that is disabling or life-threatening |

Overall, neuropathy was reported in patients previously untreated for advanced colorectal cancer in 82% (all grades) and 19% (grade 3/4), and in the previously treated patients in 74% (all grades) and 7% (grade 3/4) events. Information regarding reversibility of neuropathy was not available from the trial for patients who had not been previously treated for colorectal cancer.

Reversible Posterior Leukoencephalopathy Syndrome

Reversible Posterior Leukoencephalopathy Syndrome (RPLS, also known as PRES, Posterior Reversible Encephalopathy Syndrome) has been observed in clinical trials (< 0.1%) and postmarketing experience. Signs and symptoms of RPLS could be headache, altered mental functioning, seizures, abnormal vision from blurriness to blindness, associated or not with hypertension [see *Adverse Reactions* (6.2)]. Diagnosis of RPLS is based upon confirmation by brain imaging.

5.3 Severe Neutropenia

Grade 3 or 4 neutropenia occurred in 41-44% of patients with colorectal cancer treated with oxaliplatin injection in combination with 5-fluorouracil (5-FU) and leucovorin compared to 5% with 5-FU plus leucovorin alone. Sepsis, neutropenic sepsis and septic shock have been reported in patients treated with oxaliplatin injection, including fatal outcomes [see *Adverse Reactions* (6.1)].

Delay oxaliplatin injection until neutrophils are $\geq 1.5 \times 10^9/L$. Withhold oxaliplatin injection for sepsis or septic shock. Dose reduce oxaliplatin injection after recovery from Grade 4 neutropenia or febrile neutropenia [see *Dosage and Administration* (2.2)].

5.4 Pulmonary Toxicity

Oxaliplatin injection has been associated with pulmonary fibrosis (<1% of study patients), which may be fatal. The combined incidence of cough and dyspnea was 7.4% (any grade) and <1% (grade 3) with no grade 4 events in the oxaliplatin injection plus infusional 5-fluorouracil/leucovorin arm compared to 4.5% (any grade) and no grade 3 and 0.1% grade 4 events in the infusional 5-fluorouracil/leucovorin alone arm in adjuvant colon cancer patients. In this study, one patient died from eosinophilic pneumonia in the oxaliplatin injection combination arm. The combined incidence of cough, dyspnea and hypoxia was 43% (any grade) and 7% (grade 3 and 4) in the oxaliplatin injection plus 5-fluorouracil/leucovorin arm compared to 32% (any grade) and 5% (grade 3 and 4) in the irinotecan plus 5-fluorouracil/leucovorin arm of unknown duration for patients with previously untreated colorectal

cancer. In case of unexplained respiratory symptoms such as non-productive cough, dyspnea, crackles, or radiological pulmonary infiltrates, oxaliplatin injection should be discontinued until further pulmonary investigation excludes interstitial lung disease or pulmonary fibrosis.

5.5 Hepatotoxicity

Hepatotoxicity as evidenced in the adjuvant study, by increase in transaminases (57% vs. 34%) and alkaline phosphatase (42% vs. 20%) was observed more commonly in the oxaliplatin injection combination arm than in the control arm. The incidence of increased bilirubin was similar on both arms. Changes noted on liver biopsies include: peliosis, nodular regenerative hyperplasia or sinusoidal alterations, perisinusoidal fibrosis, and veno-occlusive lesions. Hepatic vascular disorders should be considered, and if appropriate, should be investigated in case of abnormal liver function test results or portal hypertension, which cannot be explained by liver metastases [see *Clinical Trials Experience (6.1)*].

5.6 Cardiovascular Toxicity

QT prolongation and ventricular arrhythmias including fatal Torsade de Pointes have been reported in postmarketing experiences following oxaliplatin injection administration. ECG monitoring is recommended if therapy is initiated in patients with congestive heart failure, bradyarrhythmias, drugs known to prolong the QT interval, including Class Ia and III antiarrhythmics, and electrolyte abnormalities. Correct hypokalemia or hypomagnesemia prior to initiating oxaliplatin injection and monitor these electrolytes periodically during therapy. Avoid oxaliplatin injection in patients with congenital long QT syndrome [see *Adverse Reactions (6.2)*].

5.7 Rhabdomyolysis

Rhabdomyolysis, including fatal cases, has been reported in patients treated with oxaliplatin injection. Discontinue oxaliplatin injection if any signs or symptoms of rhabdomyolysis occur. [see *Adverse Reactions (6.2)*].

5.8 Use in Pregnancy

Pregnancy Category D

Oxaliplatin injection may cause fetal harm when administered to a pregnant woman. There are no adequate and well-controlled studies of oxaliplatin injection in pregnant women. Women of childbearing potential should be advised to avoid becoming pregnant while receiving treatment with oxaliplatin injection. [see *Use in Specific Populations (8.1)*].

5.9 Recommended Laboratory Tests

Standard monitoring of the white blood cell count with differential, hemoglobin, platelet count, and blood chemistries (including ALT, AST, bilirubin and creatinine) is recommended before each oxaliplatin injection cycle [see *Dosage and Administration (2)*].

There have been reports while on study and from postmarketing surveillance of prolonged prothrombin time and INR occasionally associated with hemorrhage in patients who received oxaliplatin injection plus 5-fluorouracil/leucovorin while on anticoagulants. Patients receiving oxaliplatin injection plus 5-fluorouracil/leucovorin and requiring oral anticoagulants may require closer monitoring.

6 ADVERSE REACTIONS

The following serious adverse reactions are discussed in greater detail in other sections of the label:

- Anaphylaxis and Allergic reactions [see *Boxed Warning, Warnings and Precautions (5.1)*.]
- Neuropathy [see *Warnings and Precautions (5.2)*.]
- Severe Neutropenia [see *Warnings and Precautions (5.3)*.]
- Pulmonary Toxicities [see *Warnings and Precautions (5.4)*.]
- Hepatotoxicity [see *Warnings and Precautions (5.5)*.]
- Cardiovascular Toxicities [see *Warnings and Precautions (5.6)*.]
- Rhabdomyolysis [see *Warnings and Precautions (5.7)*.]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

More than 1100 patients with stage II or III colon cancer and more than 4,000 patients with advanced colorectal cancer have been treated in clinical studies with oxaliplatin injection. The most common adverse reactions in patients with stage II or III colon cancer receiving adjuvant therapy were peripheral sensory neuropathy, neutropenia, thrombocytopenia, anemia, nausea, increase in transaminases and alkaline phosphatase, diarrhea, emesis, fatigue and stomatitis. The most common adverse reactions in previously untreated and treated patients were peripheral sensory neuropathies, fatigue, neutropenia, nausea, emesis, and diarrhea [see *Warnings and Precautions (5)*].

Combination Adjuvant Therapy with Oxaliplatin Injection and Infusional 5-fluorouracil/leucovorin in Patients with Colon Cancer

One thousand one hundred and eight patients with stage II or III colon cancer, who had undergone complete resection of the primary tumor, have been treated in a clinical study with oxaliplatin injection in combination with infusional 5-fluorouracil/leucovorin [see *Clinical Studies (14)*]. The incidence of grade 3 or 4 adverse reactions was 70% on the oxaliplatin injection combination arm, and 31% on the infusional 5-fluorouracil/leucovorin arm. The adverse reactions in this trial are shown in the tables below. Discontinuation of treatment due to adverse reactions occurred in 15% of the patients receiving oxaliplatin injection and infusional 5-fluorouracil/leucovorin. Both 5-fluorouracil/leucovorin and oxaliplatin injection are associated with gastrointestinal or hematologic adverse reactions. When oxaliplatin injection is administered in combination with infusional 5-fluorouracil/leucovorin, the incidence of these events is increased.

The incidence of death within 28 days of last treatment, regardless of causality, was 0.5% (n=6) in both the oxaliplatin injection combination and infusional 5-fluorouracil/leucovorin arms, respectively. Deaths within 60 days from initiation of therapy were 0.3% (n=3) in both the oxaliplatin injection combination and infusional 5-fluorouracil/leucovorin arms, respectively. On the oxaliplatin injection combination arm, 3 deaths were due to sepsis/neutropenic sepsis, 2 from intracerebral bleeding and one from eosinophilic pneumonia. On the 5-fluorouracil/leucovorin arm, one death was due to suicide, 2 from Steven-Johnson Syndrome (1 patient also had sepsis), 1 unknown cause, 1 anoxic cerebral infarction and 1 probable abdominal aorta rupture.

The following table provides adverse reactions reported in the adjuvant therapy colon cancer clinical trial [see *Clinical Studies (14)*] by body system and decreasing order of frequency in the oxaliplatin injection and infusional 5-fluorouracil/leucovorin arm for events with overall incidences $\geq 5\%$ and for NCI grade 3/4 events with incidences $\geq 1\%$.

Table 3 - Adverse Reactions Reported in Patients with Colon Cancer receiving Adjuvant Treatment ($\geq 5\%$ of all patients and with $\geq 1\%$ NCI Grade 3/4 events)

| Adverse reaction (WHO/Pref) | Oxaliplatin Injection + 5-FU/LV N=1108 | | 5-FU/LV N=1111 | |
|---------------------------------------|---|---------------|-------------------|---------------|
| | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) |
| Any Event | 100 | 70 | 99 | 31 |
| Allergy/Immunology | | | | |
| Allergic Reaction | 10 | 3 | 2 | <1 |
| Constitutional Symptoms/Pain | | | | |
| Fatigue | 44 | 4 | 38 | 1 |
| Abdominal Pain | 18 | 1 | 17 | 2 |
| Dermatology/Skin | | | | |
| Skin Disorder | 32 | 2 | 36 | 2 |
| Injection Site Reaction* | 11 | 3 | 10 | 3 |
| Gastrointestinal | | | | |
| Nausea | 74 | 5 | 61 | 2 |
| Diarrhea | 56 | 11 | 48 | 7 |
| Vomiting | 47 | 6 | 24 | 1 |
| Stomatitis | 42 | 3 | 40 | 2 |
| Anorexia | 13 | 1 | 8 | <1 |
| Fever/Infection | | | | |
| Fever | 27 | 1 | 12 | 1 |
| Infection | 25 | 4 | 25 | 3 |
| Neurology | | | | |
| Overall Peripheral Sensory Neuropathy | 92 | 12 | 16 | <1 |

* Includes thrombosis related to the catheter

The following table provides adverse reactions reported in the adjuvant therapy colon cancer clinical trial [see *Clinical Studies (14)*] by body system and decreasing order of frequency in the oxaliplatin injection and infusional 5-fluorouracil/leucovorin arm for events with overall incidences $\geq 5\%$ but with incidences $<1\%$ NCI grade 3/4 events.

Table 4 - Adverse Reactions Reported in Patients with Colon Cancer receiving Adjuvant Treatment ($\geq 5\%$ of all patients, but with $<1\%$ NCI Grade 3/4 events)

| Adverse reaction (WHO/Pref) | Oxaliplatin Injection + 5-FU/LV N=1108 | 5-FU/LV N=1111 |
|---|---|-------------------|
| | All Grades (%) | All Grades (%) |
| Allergy/Immunology | | |
| Rhinitis | 6 | 8 |
| Constitutional Symptoms/Pain/Ocular/Visual | | |
| Epistaxis | 16 | 12 |
| Weight Increase | 10 | 10 |
| Conjunctivitis | 9 | 15 |
| Headache | 7 | 5 |
| Dyspnea | 5 | 3 |
| Pain | 5 | 5 |
| Lacrimation Abnormal | 4 | 12 |
| Dermatology/Skin | | |
| Alopecia | 30 | 28 |

| Gas trointes tinal | | |
|------------------------------|----|----|
| Constipation | 22 | 19 |
| Taste Perversion | 12 | 8 |
| Dyspepsia | 8 | 5 |
| Metabolic | | |
| Phosphate Alkaline increased | 42 | 20 |
| Neurology | | |
| Sensory Disturbance | 8 | 1 |

Although specific events can vary, the overall frequency of adverse reactions was similar in men and women and in patients <65 and ≥65 years. However, the following grade 3/4 events were more common in females: diarrhea, fatigue, granulocytopenia, nausea and vomiting. In patients ≥65 years old, the incidence of grade 3/4 diarrhea and granulocytopenia was higher than in younger patients. Insufficient subgroup sizes prevented analysis of safety by race. The following additional adverse reactions, were reported in ≥2% and <5% of the patients in the oxaliplatin injection and infusional 5-fluorouracil/leucovorin combination arm (listed in decreasing order of frequency): pain, leukopenia, weight decrease, coughing.

The number of patients who developed secondary malignancies was similar; 62 in the oxaliplatin injection combination arm and 68 in the infusional 5-fluorouracil/leucovorin arm. An exploratory analysis showed that the number of deaths due to secondary malignancies was 1.96% in the oxaliplatin injection combination arm and 0.98% in infusional 5-fluorouracil/leucovorin arm. In addition, the number of cardiovascular deaths was 1.4% in the oxaliplatin injection combination arm as compared to 0.7% in the infusional 5-fluorouracil/leucovorin arm. Clinical significance of these findings is unknown.

Patients Previously Untreated for Advanced Colorectal Cancer

Two hundred and fifty-nine patients were treated in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm of the randomized trial in patients previously untreated for advanced colorectal cancer [see *Clinical Studies (14)*]. The adverse reaction profile in this study was similar to that seen in other studies and the adverse reactions in this trial are shown in the tables below.

Both 5-fluorouracil and oxaliplatin injection are associated with gastrointestinal and hematologic adverse reactions. When oxaliplatin injection is administered in combination with 5-fluorouracil, the incidence of these events is increased.

The incidence of death within 30 days of treatment in the previously untreated for advanced colorectal cancer study, regardless of causality, was 3% with the oxaliplatin injection and 5-fluorouracil/leucovorin combination, 5% with irinotecan plus 5-fluorouracil/leucovorin, and 3% with oxaliplatin injection plus irinotecan. Deaths within 60 days from initiation of therapy were 2.3% with the oxaliplatin injection and 5-fluorouracil/leucovorin combination, 5.1% with irinotecan plus 5-fluorouracil/leucovorin, and 3.1% with oxaliplatin injection plus irinotecan. The following table provides adverse reactions reported in the previously untreated for advanced colorectal cancer study [see *Clinical Studies (14)*] by body system and decreasing order of frequency in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm for events with overall incidences ≥5% and for grade 3/4 events with incidences ≥1%.

Table 5 – Adverse Reactions Reported in Patients Previously Untreated for Advanced Colorectal Cancer Clinical Trial (≥5% of all patients and with ≥1% NCI Grade 3/4 events)

| Adverse reaction (WHO/Pref) | Oxaliplatin Injection + 5-FU/LV N=259 | | Irinotecan + 5-FU/LV N=256 | | Oxaliplatin Injection + Irinotecan N=258 | |
|---|--|---------------------|----------------------------------|------------------|---|---------------------|
| | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) |
| Any Event | 99 | 82 | 98 | 70 | 99 | 76 |
| Allergy/Immunology | | | | | | |
| Hypersensitivity | 12 | 2 | 5 | 0 | 6 | 1 |
| Cardiovascular | | | | | | |
| Thrombosis | 6 | 5 | 6 | 6 | 3 | 3 |
| Hypotension | 5 | 3 | 6 | 3 | 4 | 3 |
| Constitutional Symptoms/Pain/Ocular/Visual | | | | | | |
| Fatigue | 70 | 7 | 58 | 11 | 66 | 16 |
| Abdominal Pain | 29 | 8 | 31 | 7 | 39 | 10 |
| Myalgia | 14 | 2 | 6 | 0 | 9 | 2 |
| Pain | 7 | 1 | 5 | 1 | 6 | 1 |
| Vision abnormal | 5 | 0 | 2 | 1 | 6 | 1 |
| Neuralgia | 5 | 0 | 0 | 0 | 2 | 1 |
| Dermatology/Skin | | | | | | |
| Skin reaction – hand/foot | 7 | 1 | 2 | 1 | 1 | 0 |
| Injection site reaction | 6 | 0 | 1 | 0 | 4 | 1 |
| Gastrointestinal | | | | | | |
| Nausea | 71 | 6 | 67 | 15 | 83 | 19 |
| Diarrhea | 56 | 12 | 65 | 29 | 76 | 25 |
| Vomiting | 41 | 4 | 43 | 13 | 64 | 23 |
| Stomatitis | 38 | 0 | 25 | 1 | 19 | 1 |
| Anorexia | 35 | 2 | 25 | 4 | 27 | 5 |

| | | | | | | |
|---|----|----|----|----|----|----|
| Constipation | 32 | 4 | 27 | 2 | 21 | 2 |
| Diarrhea-colostomy | 13 | 2 | 16 | 7 | 16 | 3 |
| Gastrointestinal NOS* | 5 | 2 | 4 | 2 | 3 | 2 |
| Hematology/Infection | | | | | | |
| Infection normal ANC [†] | 10 | 4 | 5 | 1 | 7 | 2 |
| Infection low ANC [†] | 8 | 8 | 12 | 11 | 9 | 8 |
| Lymphopenia | 6 | 2 | 4 | 1 | 5 | 2 |
| Febrile neutropenia | 4 | 4 | 15 | 14 | 12 | 11 |
| Hepatic/Metabolic/Laboratory/Renal | | | | | | |
| Hyperglycemia | 14 | 2 | 11 | 3 | 12 | 3 |
| Hypokalemia | 11 | 3 | 7 | 4 | 6 | 2 |
| Dehydration | 9 | 5 | 16 | 11 | 14 | 7 |
| Hypoalbuminemia | 8 | 0 | 5 | 2 | 9 | 1 |
| Hyponatremia | 8 | 2 | 7 | 4 | 4 | 1 |
| Urinary frequency | 5 | 1 | 2 | 1 | 3 | 1 |
| Neurology | | | | | | |
| Overall Neuropathy | 82 | 19 | 18 | 2 | 69 | 7 |
| Paresthesias | 77 | 18 | 16 | 2 | 62 | 6 |
| Pharyngo-laryngeal dysesthesias | 38 | 2 | 1 | 0 | 28 | 1 |
| Neuro-sensory | 12 | 1 | 2 | 0 | 9 | 1 |
| Neuro NOS* | 1 | 0 | 1 | 0 | 1 | 0 |
| Pulmonary | | | | | | |
| Cough | 35 | 1 | 25 | 2 | 17 | 1 |
| Dyspnea | 18 | 7 | 14 | 3 | 11 | 2 |
| Hiccups | 5 | 1 | 2 | 0 | 3 | 2 |

* Not otherwise specified

† Absolute neutrophil count

The following table provides adverse reactions reported in the previously untreated for advanced colorectal cancer study [see *Clinical Studies (14)*] by body system and decreasing order of frequency in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm for events with overall incidences ≥5% but with incidences <1% NCI Grade 3/4 events.

Table 6 - Adverse Reactions Reported in Patients Previously Untreated for Advanced Colorectal Cancer Clinical Trial (≥5% of all patients but with < 1% NCI Grade 3/4 events)

| Adverse reaction (WHO/Pref) | Oxaliplatin Injection + 5-FU/LV N=259 | Irinotecan + 5-FU/LV N=256 | Oxaliplatin Injection + Irinotecan N=258 |
|---|--|-------------------------------|---|
| | All Grades (%) | All Grades (%) | All Grades (%) |
| Allergy/Immunology | | | |
| Rash | 11 | 4 | 7 |
| Rhinitis allergic | 10 | 6 | 6 |
| Cardiovascular | | | |
| Edema | 15 | 13 | 10 |
| Constitutional Symptoms/Pain/Ocular/Visual | | | |
| Headache | 13 | 6 | 9 |
| Weight loss | 11 | 9 | 11 |
| Epistaxis | 10 | 2 | 2 |
| Tearing | 9 | 1 | 2 |
| Rigors | 8 | 2 | 7 |
| Dysphasia | 5 | 3 | 3 |
| Sweating | 5 | 6 | 12 |
| Arthralgia | 5 | 5 | 8 |
| Dermatology/Skin | | | |
| Alopecia | 38 | 44 | 67 |
| Flushing | 7 | 2 | 5 |
| Pruritis | 6 | 4 | 2 |
| Dry Skin | 6 | 2 | 5 |
| Gastrointestinal | | | |
| Taste perversion | 14 | 6 | 8 |
| Dyspepsia | 12 | 7 | 5 |
| Flatulence | 9 | 6 | 5 |
| Mouth Dryness | 5 | 2 | 3 |
| Hematology/Infection | | | |
| Fever normal ANC* | 16 | 9 | 9 |
| Hepatic/Metabolic/Laboratory/Renal | | | |
| Hypocalcemia | 7 | 5 | 4 |
| Elevated Creatinine | 4 | 4 | 5 |
| Neurology | | | |
| Insomnia | 13 | 9 | 11 |
| Depression | 9 | 5 | 7 |
| Dizziness | 8 | 6 | 10 |
| Anxiety | 5 | 2 | 6 |

* Absolute neutrophil count

Adverse reactions were similar in men and women and in patients <65 and ≥65 years, but older patients may have been more susceptible to diarrhea, dehydration, hypokalemia, leukopenia, fatigue and syncope.

The following additional adverse reactions, at least possibly related to treatment and potentially important, were reported in $\geq 2\%$ and $< 5\%$ of the patients in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm (listed in decreasing order of frequency): metabolic, pneumonitis, catheter infection, vertigo, prothrombin time, pulmonary, rectal bleeding, dysuria, nail changes, chest pain, rectal pain, syncope, hypertension, hypoxia, unknown infection, bone pain, pigmentation changes, and urticaria.

Previously Treated Patients with Advanced Colorectal Cancer

Four hundred and fifty patients (about 150 receiving the combination of oxaliplatin injection and 5-fluorouracil/leucovorin) were studied in a randomized trial in patients with refractory and relapsed colorectal cancer [see Clinical Studies (14)]. The adverse reaction profile in this study was similar to that seen in other studies and the adverse reactions in this trial are shown in the tables below. Thirteen percent of patients in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm and 18% in the 5-fluorouracil/leucovorin arm of the previously treated study had to discontinue treatment because of adverse effects related to gastrointestinal, or hematologic adverse reactions, or neuropathies. Both 5-fluorouracil and oxaliplatin injection are associated with gastrointestinal and hematologic adverse reactions. When oxaliplatin injection is administered in combination with 5-fluorouracil, the incidence of these events is increased.

The incidence of death within 30 days of treatment in the previously treated study, regardless of causality, was 5% with the oxaliplatin injection and 5-fluorouracil/leucovorin combination, 8% with oxaliplatin injection alone, and 7% with 5-fluorouracil/leucovorin. Of the 7 deaths that occurred on the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm within 30 days of stopping treatment, 3 may have been treatment related, associated with gastrointestinal bleeding or dehydration.

The following table provides adverse reactions reported in the previously treated study [see Clinical Studies (14)] by body system and in decreasing order of frequency in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm for events with overall incidences $\geq 5\%$ and for grade 3/4 events with incidences $\geq 1\%$. This table does not include hematologic and blood chemistry abnormalities; these are shown separately below.

Table 7 – Adverse Reactions Reported In Previously Treated Colorectal Cancer Clinical Trial ($\geq 5\%$ of all patients and with $\geq 1\%$ NCI Grade 3/4 events)

| Adverse reaction (WHO/Pref) | 5-FU/LV (N = 142) | | Oxaliplatin Injection (N = 153) | | Oxaliplatin Injection + 5-FU/LV (N = 150) | |
|---|----------------------|---------------|------------------------------------|---------------|--|---------------|
| | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) |
| Any Event | 98 | 41 | 100 | 46 | 99 | 73 |
| Cardiovascular | | | | | | |
| Dyspnea | 11 | 2 | 13 | 7 | 20 | 4 |
| Coughing | 9 | 0 | 11 | 0 | 19 | 1 |
| Edema | 13 | 1 | 10 | 1 | 15 | 1 |
| Thromboembolism | 4 | 2 | 2 | 1 | 9 | 8 |
| Chest Pain | 4 | 1 | 5 | 1 | 8 | 1 |
| Constitutional Symptoms/Pain | | | | | | |
| Fatigue | 52 | 6 | 61 | 9 | 68 | 7 |
| Back Pain | 16 | 4 | 11 | 0 | 19 | 3 |
| Pain | 9 | 3 | 14 | 3 | 15 | 2 |
| Dermatology/Skin | | | | | | |
| Injection Site Reaction | 5 | 1 | 9 | 0 | 10 | 3 |
| Gastrointestinal | | | | | | |
| Diarrhea | 44 | 3 | 46 | 4 | 67 | 11 |
| Nausea | 59 | 4 | 64 | 4 | 65 | 11 |
| Vomiting | 27 | 4 | 37 | 4 | 40 | 9 |
| Stomatitis | 32 | 3 | 14 | 0 | 37 | 3 |
| Abdominal Pain | 31 | 5 | 31 | 7 | 33 | 4 |
| Anorexia | 20 | 1 | 20 | 2 | 29 | 3 |
| Gastroesophageal Reflux | 3 | 0 | 1 | 0 | 5 | 2 |
| Hematology/Infection | | | | | | |
| Fever | 23 | 1 | 25 | 1 | 29 | 1 |
| Febrile Neutropenia | 1 | 1 | 0 | 0 | 6 | 6 |
| Hepatic/Metabolic/Laboratory/Renal | | | | | | |
| Hypokalemia | 3 | 1 | 3 | 2 | 9 | 4 |
| Dehydration | 6 | 4 | 5 | 3 | 8 | 3 |
| Neurology | | | | | | |
| Neuropathy | 17 | 0 | 76 | 7 | 74 | 7 |
| Acute | 10 | 0 | 65 | 5 | 56 | 2 |
| Persistent | 9 | 0 | 43 | 3 | 48 | 6 |

The following table provides adverse reactions reported in the previously treated study [see Clinical Studies (14)] by body system and in decreasing order of frequency in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm for events with overall incidences $\geq 5\%$ but with incidences $< 1\%$ NCI Grade 3/4 events.

Table 8 - Adverse Reactions Reported In Previously Treated Colorectal Cancer Clinical Trial ($\geq 5\%$ of all patients but with $< 1\%$ NCI Grade 3/4 events)

| Adverse reaction (WHO/Pref) | 5-FU/LV (N = 142) | Oxaliplatin Injection (N = 153) | Oxaliplatin Injection + 5-FU/LV (N = 150) |
|---|----------------------|------------------------------------|--|
| | All Grades (%) | All Grades (%) | All Grades (%) |
| Allergy/Immunology | | | |
| Rhinitis | 4 | 6 | 15 |
| Allergic Reaction | 1 | 3 | 10 |
| Rash | 5 | 5 | 9 |
| Cardiovascular | | | |
| Peripheral Edema | 11 | 5 | 10 |
| Constitutional Symptoms/Pain/Ocular/Visual | | | |
| Headache | 8 | 13 | 17 |
| Arthralgia | 10 | 7 | 10 |
| Epistaxis | 1 | 2 | 9 |
| Abnormal Lacrimation | 6 | 1 | 7 |
| Rigors | 6 | 9 | 7 |
| Dermatology/Skin | | | |
| Hand-Foot Syndrome | 13 | 1 | 11 |
| Flushing | 2 | 3 | 10 |
| Alopecia | 3 | 3 | 7 |
| Gastrointestinal | | | |
| Constipation | 23 | 31 | 32 |
| Dyspepsia | 10 | 7 | 14 |
| Taste Perversion | 1 | 5 | 13 |
| Mucositis | 10 | 2 | 7 |
| Flatulence | 6 | 3 | 5 |
| Hepatic/Metabolic/Laboratory/Renal | | | |
| Hematuria | 4 | 0 | 6 |
| Dysuria | 1 | 1 | 6 |
| Neurology | | | |
| Dizziness | 8 | 7 | 13 |
| Insomnia | 4 | 11 | 9 |
| Pulmonary | | | |
| Upper Resp Tract Infection | 4 | 7 | 10 |
| Pharyngitis | 10 | 2 | 9 |
| Hiccup | 0 | 2 | 5 |

Adverse reactions were similar in men and women and in patients <65 and ≥65 years, but older patients may have been more susceptible to dehydration, diarrhea, hypokalemia and fatigue. The following additional adverse reactions, at least possibly related to treatment and potentially important, were reported in ≥2% and <5% of the patients in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm (listed in decreasing order of frequency): anxiety, myalgia, erythematous rash, increased sweating, conjunctivitis, weight decrease, dry mouth, rectal hemorrhage, depression, ataxia, ascites, hemorrhoids, muscle weakness, nervousness, tachycardia, abnormal micturition frequency, dry skin, pruritus, hemoptysis, purpura, vaginal hemorrhage, melena, somnolence, pneumonia, proctitis, involuntary muscle contractions, intestinal obstruction, gingivitis, tenesmus, hot flashes, enlarged abdomen, urinary incontinence.

Hematologic Changes

The following tables list the hematologic changes occurring in ≥5% of patients, based on laboratory values and NCI grade, with the exception of those events occurring in adjuvant patients and anemia in the patients previously untreated for advanced colorectal cancer, respectively, which are based on AE reporting and NCI grade alone.

Table 9 - Adverse Hematologic Reactions in Patients with Colon Cancer Receiving Adjuvant Therapy (≥5% of patients)

| Hematology Parameter | Oxaliplatin Injection + 5-FU/LV (N=1108) | | 5-FU/LV (N=1111) | |
|----------------------|---|---------------|---------------------|---------------|
| | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) |
| Anemia | 76 | 1 | 67 | <1 |
| Neutropenia | 79 | 41 | 40 | 5 |
| Thrombocytopenia | 77 | 2 | 19 | <1 |

Table 10 – Adverse Hematologic Reactions in Patients Previously Untreated for Advanced Colorectal Cancer (≥5% of patients)

| Hematology Parameter | Oxaliplatin Injection + 5-FU/LV N=259 | | Irinotecan+ 5-FU/LV N=256 | | Oxaliplatin Injection + Irinotecan N=258 | |
|----------------------|--|---------------|---------------------------------|---------------|---|---------------|
| | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) |
| Anemia | 27 | 3 | 28 | 4 | 25 | 3 |
| Leukopenia | 85 | 20 | 84 | 23 | 76 | 24 |
| Neutropenia | 81 | 53 | 77 | 44 | 71 | 36 |
| Thrombocytopenia | 71 | 5 | 26 | 2 | 44 | 4 |

Table 11 – Adverse Hematologic Reactions in Previously Treated Patients (≥5% of patients)

| Hematology Parameter | 5-FU/LV (N=142) | | Oxaliplatin Injection (N=153) | | Oxaliplatin Injection + 5-FU/LV (N=150) | |
|----------------------|-----------------|---------------|-------------------------------|---------------|---|---------------|
| | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) |
| Anemia | 68 | 2 | 64 | 1 | 81 | 2 |
| Leukopenia | 34 | 1 | 13 | 0 | 76 | 19 |
| Neutropenia | 25 | 5 | 7 | 0 | 73 | 44 |
| Thrombocytopenia | 20 | 0 | 30 | 3 | 64 | 4 |

Thrombocytopenia and Bleeding

Thrombocytopenia was frequently reported with the combination of oxaliplatin injection and infusional 5-fluorouracil/leucovorin. The incidence of all hemorrhagic events in the adjuvant and previously treated patients was higher on the oxaliplatin injection combination arm compared to the infusional 5-fluorouracil/leucovorin arm. These events included gastrointestinal bleeding, hematuria, and epistaxis. In the adjuvant trial, two patients died from intracerebral hemorrhages.

The incidence of Grade 3/4 thrombocytopenia was 2% in adjuvant patients with colon cancer. In patients treated for advanced colorectal cancer the incidence of Grade 3/4 thrombocytopenia was 3 to 5%, and the incidence of these events was greater for the combination of oxaliplatin injection and 5-fluorouracil/leucovorin over the irinotecan plus 5-fluorouracil/leucovorin or 5-fluorouracil/leucovorin control groups. Grade 3/4 gastrointestinal bleeding was reported in 0.2% of adjuvant patients receiving oxaliplatin injection and 5-fluorouracil/leucovorin. In the previously untreated patients, the incidence of epistaxis was 10% in the oxaliplatin injection and 5-fluorouracil/leucovorin arm, and 2% and 1%, respectively, in the irinotecan plus 5-fluorouracil/leucovorin or irinotecan plus oxaliplatin injection arms.

Neutropenia

Neutropenia was frequently observed with the combination of oxaliplatin injection and 5-fluorouracil/leucovorin, with Grade 3 and 4 events reported in 29% and 12% of adjuvant patients with colon cancer, respectively. In the adjuvant trial, 3 patients died from sepsis/neutropenic sepsis. Grade 3 and 4 events were reported in 35% and 18% of the patients previously untreated for advanced colorectal cancer, respectively. Grade 3 and 4 events were reported in 27% and 17% of previously treated patients, respectively. In adjuvant patients the incidence of either febrile neutropenia (0.7%) or documented infection with concomitant grade 3/4 neutropenia (1.1%) was 1.8% in the oxaliplatin injection and 5-fluorouracil/leucovorin arm. The incidence of febrile neutropenia in the patients previously untreated for advanced colorectal cancer was 15% (3% of cycles) in the irinotecan plus 5-fluorouracil/leucovorin arm and 4% (less than 1% of cycles) in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm. Additionally, in this same population, infection with grade 3 or 4 neutropenia was 12% in the irinotecan plus 5-fluorouracil/leucovorin, and 8% in the oxaliplatin injection and 5-fluorouracil/leucovorin combination. The incidence of febrile neutropenia in the previously treated patients was 1% in the 5-fluorouracil/leucovorin arm and 6% (less than 1% of cycles) in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm.

Gastrointestinal

In patients receiving the combination of oxaliplatin injection plus infusional 5-fluorouracil/leucovorin for adjuvant treatment for colon cancer the incidence of Grade 3/4 nausea and vomiting was greater than those receiving infusional 5-fluorouracil/leucovorin alone (see table). In patients previously untreated for advanced colorectal cancer receiving the combination of oxaliplatin injection and 5-fluorouracil/leucovorin, the incidence of Grade 3 and 4 vomiting and diarrhea was less compared to irinotecan plus 5-fluorouracil/leucovorin controls (see table). In previously treated patients receiving the combination of oxaliplatin injection and 5-fluorouracil/leucovorin, the incidence of Grade 3 and 4 nausea, vomiting, diarrhea, and mucositis/stomatitis increased compared to 5-fluorouracil/leucovorin controls (see table).

The incidence of gastrointestinal adverse reactions in the previously untreated and previously treated patients appears to be similar across cycles. Premedication with antiemetics, including 5-HT₃ blockers, is recommended. Diarrhea and mucositis may be exacerbated by the addition of oxaliplatin injection to 5-fluorouracil/leucovorin, and should be managed with appropriate supportive care. Since cold temperature can exacerbate acute neurological symptoms, ice (mucositis prophylaxis) should be avoided during the infusion of oxaliplatin injection.

Dermatologic

Oxaliplatin injection did not increase the incidence of alopecia compared to 5-fluorouracil/leucovorin alone. No complete alopecia was reported. The incidence of Grade 3/4 skin disorders was 2% in both the oxaliplatin injection plus infusional 5-fluorouracil/leucovorin and the infusional 5-fluorouracil/leucovorin alone arms in the adjuvant colon cancer patients. The incidence of hand-foot syndrome in patients previously untreated for advanced colorectal cancer was 2% in the irinotecan plus 5-fluorouracil/leucovorin arm and 7% in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm. The incidence of hand-foot syndrome in previously treated patients was 13% in the 5-fluorouracil/leucovorin arm and 11% in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm.

Intravenous Site Reactions

Extravasation, in some cases including necrosis, has been reported. Injection site reaction, including redness, swelling, and pain, has been reported.

Anticoagulation and Hemorrhage

There have been reports while on study and from postmarketing surveillance of prolonged prothrombin time and INR occasionally associated with hemorrhage in patients who received oxaliplatin injection plus 5-fluorouracil/leucovorin while on anticoagulants. Patients receiving oxaliplatin injection plus 5-fluorouracil/leucovorin and requiring oral anticoagulants may require closer monitoring.

Renal

About 5 to 10% of patients in all groups had some degree of elevation of serum creatinine. The incidence of Grade 3/4 elevations in serum creatinine in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm was 1% in the previously treated patients. Serum creatinine measurements were not reported in the adjuvant trial.

Hepatic

Hepatotoxicity (defined as elevation of liver enzymes) appears to be related to oxaliplatin injection combination therapy [see *Warnings and Precautions (5.5)*]. The following tables list the clinical chemistry changes associated with hepatic toxicity occurring in ≥5% of patients, based on adverse reactions reported and NCI CTC grade for adjuvant patients and patients previously untreated for advanced colorectal cancer, laboratory values and NCI CTC grade for previously treated patients.

Table 12 - Adverse Hepatic Reactions in Patients with Stage II or III Colon Cancer Receiving Adjuvant Therapy (≥5% of patients)

| Hepatic Parameter | Oxaliplatin Injection + 5-FU/LV (N=1108) | | 5-FU/LV (N=1111) | |
|---------------------------|--|---------------|------------------|---------------|
| | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) |
| Increase in transaminases | 57 | 2 | 34 | 1 |
| ALP increased | 42 | <1 | 20 | <1 |
| Bilirubinaemia | 20 | 4 | 20 | 5 |

Table 13 – Adverse Hepatic – Clinical Chemistry Abnormalities in Patients Previously Untreated for Advanced Colorectal Cancer (≥5% of patients)

| Clinical Chemistry | Oxaliplatin Injection + 5-FU/LV N=259 | | Irinotecan + 5-FU/LV N=256 | | Oxaliplatin Injection + Irinotecan N=258 | |
|----------------------|---------------------------------------|---------------|----------------------------|---------------|--|---------------|
| | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) |
| ALT (SGPT-ALAT) | 6 | 1 | 2 | 0 | 5 | 2 |
| AST (SGOT-ASAT) | 17 | 1 | 2 | 1 | 11 | 1 |
| Alkaline Phosphatase | 16 | 0 | 8 | 0 | 14 | 2 |
| Total Bilirubin | 6 | 1 | 3 | 1 | 3 | 2 |

Table 14 – Adverse Hepatic – Clinical Chemistry Abnormalities in Previously Treated Patients (≥5% of patients)

| Clinical Chemistry | 5-FU/LV (N=142) | | Oxaliplatin Injection (N=153) | | Oxaliplatin Injection + 5-FU/LV (N=150) | |
|--------------------|-----------------|---------------|-------------------------------|---------------|---|---------------|
| | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) | All Grades (%) | Grade 3/4 (%) |
| ALT (SGPT-ALAT) | 28 | 3 | 36 | 1 | 31 | 0 |
| AST (SGOT-ASAT) | 39 | 2 | 54 | 4 | 47 | 0 |
| Total Bilirubin | 22 | 6 | 13 | 5 | 13 | 1 |

Thromboembolism

The incidence of thromboembolic events in adjuvant patients with colon cancer was 6% (1.8% grade 3/4) in the infusional 5-fluorouracil/leucovorin arm and 6% (1.2% grade 3/4) in the oxaliplatin injection and infusional 5-fluorouracil/leucovorin combined arm, respectively. The incidence was 6 and 9% of the patients previously untreated for advanced colorectal cancer and previously treated patients in the oxaliplatin injection and 5-fluorouracil/leucovorin combination arm, respectively.

6.2 Postmarketing Experience

The following adverse reactions have been identified during post-approval use of oxaliplatin injection. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

Body as a whole:

angioedema, anaphylactic shock

Cardiovascular disorders:

QT prolongation leading to ventricular arrhythmias including fatal Torsade de Pointes

Central and peripheral nervous system disorders:

loss of deep tendon reflexes, dysarthria, Lhermitte's sign, cranial nerve palsies, fasciculations, convulsion, Reversible Posterior Leukoencephalopathy Syndrome (RPLS, also known as PRES).

Hearing and vestibular system disorders:

deafness

Infections:

septic shock, including fatal outcomes

Infusion reactions/hypersensitivity:

laryngospasm

Liver and Gastrointestinal system disorders:

severe diarrhea/vomiting resulting in hypokalemia, colitis (including *Clostridium difficile* diarrhea), metabolic acidosis; ileus; intestinal obstruction, pancreatitis; veno-occlusive disease of liver also known as sinusoidal obstruction syndrome, and perisinusoidal fibrosis which rarely may progress.

Musculoskeletal and connective tissue disorders

rhabdomyolysis, including fatal outcomes.

Platelet, bleeding, and clotting disorders:

immuno-allergic thrombocytopenia

prolongation of prothrombin time and of INR in patients receiving anticoagulants

Red Blood Cell disorders:

hemolytic uremic syndrome, immuno-allergic hemolytic anemia

Renal disorders:

Acute tubular necrosis, acute interstitial nephritis and acute renal failure.

Respiratory system disorders:

pulmonary fibrosis, and other interstitial lung diseases (sometimes fatal)

Vision disorders:

decrease of visual acuity, visual field disturbance, optic neuritis and transient vision loss (reversible following therapy discontinuation)

7 DRUG INTERACTIONS

No specific cytochrome P-450-based drug interaction studies have been conducted. No pharmacokinetic interaction between 85 mg/m² oxaliplatin injection and 5-fluorouracil/leucovorin has been observed in patients treated every 2 weeks. Increases of 5-fluorouracil plasma concentrations by approximately 20% have been observed with doses of 130 mg/m² oxaliplatin injection dosed every 3 weeks. Because platinum-containing species are eliminated primarily through the kidney, clearance of these products may be decreased by coadministration of potentially nephrotoxic compounds; although, this has not been specifically studied [see *Clinical Pharmacology* (12.3)].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category D

Based on direct interaction with DNA, oxaliplatin injection may cause fetal harm when administered to a pregnant woman. There are no adequate and well-controlled studies of oxaliplatin injection in pregnant women. Reproductive toxicity studies in rats demonstrated adverse effects on fertility and embryo-fetal development at maternal doses that were below the recommended human dose based on body surface area. If this drug is used during pregnancy or if the patient becomes pregnant while taking this drug, the patient should be apprised of the potential hazard to the fetus. Women of childbearing potential should be advised to avoid becoming pregnant and use effective contraception while receiving treatment with oxaliplatin injection.

Pregnant rats were administered oxaliplatin at less than one-tenth the recommended human dose based on body surface area during gestation days 1 to 5 (pre-implantation), 6 to 10, or 11 to 16 (during organogenesis). Oxaliplatin caused developmental mortality (increased early resorptions) when administered on days 6 to 10 and 11 to 16 and adversely affected fetal growth (decreased fetal weight, delayed ossification) when administered on days 6 to 10. Administration of oxaliplatin to male and female rats prior to mating resulted in 97% post-implantation loss in animals that received approximately one-seventh the recommended human dose based on the body surface area.

8.3 Nursing Mothers

It is not known whether oxaliplatin injection or its derivatives are excreted in human milk. Because many

drugs are excreted in human milk and because of the potential for serious adverse reactions in nursing infants from oxaliplatin injection, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

8.4 Pediatric Use

The effectiveness of oxaliplatin in children has not been established. Oxaliplatin has been tested in 2 Phase 1 and 2 Phase 2 trials in 235 patients ages 7 months to 22 years with solid tumors (see below) and no significant activity observed.

In a Phase 1/2 study, oxaliplatin was administered as a 2-hour intravenous infusion on Days 1, 8 and 15 every 4 weeks (1 cycle), for a maximum of 6 cycles, to 43 patients with refractory or relapsed malignant solid tumors, mainly neuroblastoma and osteosarcoma. Twenty eight pediatric patients in the Phase 1 study received oxaliplatin at 6 dose levels starting at 40 mg/m² with escalation to 110 mg/m². The dose limiting toxicity (DLT) was sensory neuropathy at the 110 mg/m² dose. Fifteen patients received oxaliplatin at a dose of 90 mg/m² intravenous in the Phase 2 portion of the study. At this dose, paresthesia (60%, G3/4: 7%), fever (40%, G3/4: 7%) and thrombocytopenia (40%, G3/4: 27%) were the main adverse reactions. No responses were observed.

In a second Phase 1 study, oxaliplatin was administered to 26 pediatric patients as a 2-hour intravenous infusion on day 1 every 3 weeks (1 cycle) at 5 dose levels starting at 100 mg/m² with escalation to 160 mg/m², for a maximum of 6 cycles. In a separate cohort, oxaliplatin 85 mg/m² was administered on day 1 every 2 weeks, for a maximum of 9 doses. Patients had metastatic or unresectable solid tumors mainly neuroblastoma and ganglioneuroblastoma. No responses were observed. The DLT was sensory neuropathy at the 160 mg/m² dose. Based on these studies, oxaliplatin 130 mg/m² as a 2-hour intravenous infusion on day 1 every 3 weeks (1 cycle) was used in subsequent Phase II studies. A dose of 85 mg/m² on day 1 every 2 weeks was also found to be tolerable.

In one Phase 2 study, 43 pediatric patients with recurrent or refractory embryonal CNS tumors received oxaliplatin 130 mg/m² every 3 weeks for a maximum of 12 months in absence of progressive disease or unacceptable toxicity. In patients < 10 kg the oxaliplatin dose used was 4.3 mg/kg. The most common adverse reactions reported were leukopenia (67%, G3/4: 12%), anemia (65%, G3/4: 5%), thrombocytopenia (65%, G3/4: 26%), vomiting (65%, G3/4: 7%), neutropenia (58%, G3/4: 16%) and sensory neuropathy (40%, G3/4: 5%). One partial response was observed.

In a second Phase 2 study, 123 pediatric patients with recurrent solid tumors, including neuroblastoma, osteosarcoma, Ewing sarcoma or peripheral PNET, ependymoma, rhabdomyosarcoma, hepatoblastoma, high grade astrocytoma, Brain stem glioma, low grade astrocytoma, malignant germ cell tumor and other tumors of interest received oxaliplatin 130 mg/m² every 3 weeks for a maximum of 12 months or 17 cycles. In patients < 12 months old the oxaliplatin dose used was 4.3 mg/kg. The most common adverse reactions reported were sensory neuropathy (52%, G3/4: 12%), thrombocytopenia (37%, G3/4: 17%), anemia (37%, G3/4: 9%), vomiting (26%, G3/4: 4%), ALT increased (24%, G3/4: 6%), AST increased (24%, G3/4: 2%), and nausea (23%, G3/4: 3%). Two partial responses were observed.

The pharmacokinetic parameters of ultrafiltrable platinum have been evaluated in 105 pediatric patients during the first cycle. The mean clearance in pediatric patients estimated by the population pharmacokinetic analysis was 4.7 L/h. The inter-patient variability of platinum clearance in pediatric cancer patients was 41%. Mean platinum pharmacokinetic parameters in ultrafiltrate were C_{max} of 0.75 ± 0.24 mcg/mL, AUC₀₋₄₈ of 7.52 ± 5.07 mcg·h/mL and AUC_{inf} of 8.83 ± 1.57 mcg·h/mL at 85 mg/m² of oxaliplatin and C_{max} of 1.1 ± 0.43 mcg/mL, AUC₀₋₄₈ of 9.74 ± 2.52 mcg·h/mL and AUC_{inf} of 17.3 ± 5.34 mcg·h/mL at 130 mg/m² of oxaliplatin.

8.5 Geriatric Use

No significant effect of age on the clearance of ultrafilterable platinum has been observed. In the adjuvant therapy colon cancer randomized clinical trial, [see *Clinical Studies (14)*] 723 patients treated with oxaliplatin injection and infusional 5-fluorouracil/leucovorin were <65 years and 400 patients were ≥65 years. A descriptive subgroup analysis demonstrated that the improvement in DFS for the oxaliplatin injection combination arm compared to the infusional 5-fluorouracil/leucovorin alone arm appeared to be maintained across genders. The effect of oxaliplatin injection in patients ≥65 years of age was not conclusive. Insufficient subgroup sizes prevented analysis by race. Patients ≥ 65 years of age receiving the oxaliplatin injection combination therapy experienced more grade 3 and 4 granulocytopenia than patients < 65 years of age (45% versus 39%).

In the previously untreated for advanced colorectal cancer randomized clinical trial [see *Clinical Studies (14)*] of oxaliplatin injection, 160 patients treated with oxaliplatin injection and 5-fluorouracil/leucovorin were < 65 years and 99 patients were ≥65 years. The same efficacy improvements in response rate, time to tumor progression, and overall survival were observed in the ≥65 year old patients as in the overall study population. In the previously treated for advanced colorectal cancer randomized clinical trial [see *Clinical Studies (14)*] of oxaliplatin injection, 95 patients treated with oxaliplatin injection and 5-fluorouracil/leucovorin were <65 years and 55 patients were ≥65 years. The rates of overall adverse reactions, including grade 3 and 4 events, were similar across and within arms in the different age groups in all studies. The incidence of diarrhea, dehydration, hypokalemia, leukopenia, fatigue and syncope were higher in patients ≥65 years old. No adjustment to starting dose was required in patients ≥65 years old.

8.6 Patients with Renal Impairment

The exposure (AUC) of unbound platinum in plasma ultrafiltrate tends to increase in renally impaired patients [see *Pharmacokinetics* (12.3)]. Caution and close monitoring should be exercised when oxaliplatin injection is administered to patients with renal impairment. The starting oxaliplatin injection dose does not need to be reduced in patients with mild (creatinine clearance = 50 to 80 mL/min) or moderate (creatinine clearance = 30 to 49 mL/min) renal impairment. However, the starting dose of oxaliplatin injection should be reduced in patients with severe renal impairment (creatinine clearance < 30 mL/min) [see *Dosage and Administration* (2.2)].

10 OVERDOSAGE


There is no known antidote for oxaliplatin injection overdose. In addition to thrombocytopenia, the anticipated complications of an oxaliplatin injection overdose include hypersensitivity reaction, myelosuppression, nausea, vomiting, diarrhea and neurotoxicity.

Several cases of overdoses have been reported with oxaliplatin injection. Adverse reactions observed were Grade 4 thrombocytopenia (<25,000/mm³) without any bleeding, anemia, sensory neuropathy such as paresthesia, dysesthesia, laryngospasm and facial muscle spasms, gastrointestinal disorders such as nausea, vomiting, stomatitis, flatulence, abdomen enlarged and Grade 4 intestinal obstruction, Grade 4 dehydration, dyspnea, wheezing, chest pain, respiratory failure, severe bradycardia and death.

Patients suspected of receiving an overdose should be monitored, and supportive treatment should be administered. The maximum dose of oxaliplatin that has been administered in a single infusion is 825 mg.

11 DESCRIPTION

Oxaliplatin injection is an antineoplastic agent with the molecular formula C₈H₁₄N₂O₄Pt and the chemical name of *cis*-[(1*R*,2*R*)-1,2-cyclohexanediamine-*N,N'*] [oxalato(2-)-*O,O'*] platinum. Oxaliplatin is an organoplatinum complex in which the platinum atom is complexed with 1,2-diaminocyclohexane (DACH) and with an oxalate ligand as a leaving group.



The molecular weight is 397.3. Oxaliplatin is slightly soluble in water at 6 mg/mL, very slightly soluble in methanol, and practically insoluble in ethanol and acetone. Oxaliplatin injection, USP is for intravenous use. Each mL of oxaliplatin injection contains 5 mg of oxaliplatin, USP. It is supplied as a sterile, preservative-free, aqueous solution in 50 mg/10 mL and 100 mg/20 mL vials. Water for injection is present as an inactive ingredient.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Oxaliplatin undergoes nonenzymatic conversion in physiologic solutions to active derivatives via displacement of the labile oxalate ligand. Several transient reactive species are formed, including monoquo and diaquo DACH platinum, which covalently bind with macromolecules. Both inter- and intrastrand Pt-DNA crosslinks are formed. Crosslinks are formed between the *N7* positions of two adjacent guanines (GG), adjacent adenine-guanines (AG), and guanines separated by an intervening nucleotide (GNG). These crosslinks inhibit DNA replication and transcription. Cytotoxicity is cell-cycle nonspecific. *In vivo* studies have shown antitumor activity of oxaliplatin against colon carcinoma. In combination with 5-fluorouracil, oxaliplatin exhibits *in vitro* and *in vivo* antiproliferative activity greater than either compound alone in several tumor models [HT29 (colon), GR (mammary), and L1210 (leukemia)].

12.3 Pharmacokinetics

The reactive oxaliplatin derivatives are present as a fraction of the unbound platinum in plasma ultrafiltrate. The decline of ultrafilterable platinum levels following oxaliplatin administration is triphasic, characterized by two relatively short distribution phases ($t_{1/2\alpha}$; 0.43 hours and $t_{1/2\beta}$; 16.8 hours) and a long terminal elimination phase ($t_{1/2\gamma}$; 391 hours). Pharmacokinetic parameters obtained after a single 2-hour intravenous infusion of oxaliplatin injection at a dose of 85 mg/m² expressed as ultrafilterable platinum were C_{\max} of 0.814 mcg/mL and volume of distribution of 440 L. Interpatient and inpatient variability in ultrafilterable platinum exposure (AUC_{0-48hr}) assessed over 3 cycles was moderate to low (23% and 6%, respectively). A pharmacodynamic relationship between platinum

ultrafiltrate levels and clinical safety and effectiveness has not been established.

Distribution

At the end of a 2-hour infusion of oxaliplatin injection, approximately 15% of the administered platinum is present in the systemic circulation. The remaining 85% is rapidly distributed into tissues or eliminated in the urine. In patients, plasma protein binding of platinum is irreversible and is greater than 90%. The main binding proteins are albumin and gamma-globulins. Platinum also binds irreversibly and accumulates (approximately 2-fold) in erythrocytes, where it appears to have no relevant activity. No platinum accumulation was observed in plasma ultrafiltrate following 85 mg/m² every two weeks.

Metabolism

Oxaliplatin undergoes rapid and extensive nonenzymatic biotransformation. There is no evidence of cytochrome P450-mediated metabolism *in vitro*.

Up to 17 platinum-containing derivatives have been observed in plasma ultrafiltrate samples from patients, including several cytotoxic species (monochloro DACH platinum, dichloro DACH platinum, and monoquo and diaquo DACH platinum) and a number of noncytotoxic, conjugated species.

Elimination

The major route of platinum elimination is renal excretion. At five days after a single 2-hour infusion of oxaliplatin injection, urinary elimination accounted for about 54% of the platinum eliminated, with fecal excretion accounting for only about 2%. Platinum was cleared from plasma at a rate (10 to 17 L/h) that was similar to or exceeded the average human glomerular filtration rate (GFR; 7.5 L/h). There was no significant effect of gender on the clearance of ultrafilterable platinum. The renal clearance of ultrafilterable platinum is significantly correlated with GFR.

Pharmacokinetics in Special Populations

Pediatric

[See Use In Specific Patient Populations (8.4)].

Renal Impairment

A study was conducted in 38 patients with advanced GI cancer and varying degrees of renal impairment. Patients in the normal (creatinine clearance (CrCL) > 80 mL/min, N = 11), mild (CrCL = 50 to 80 mL/min, N = 13), and moderate (CrCL = 30 to 49 mL/min, N = 10) groups were treated with 85 mg/m² oxaliplatin injection and those in the severe (CrCL < 30 mL/min, N = 4) group were treated with 65 mg/m² oxaliplatin injection. The mean AUC of unbound platinum was 40%, 95%, and 342% higher in the mild, moderate, and severe groups, respectively, than in the normal group. Mean C_{max} of unbound platinum appeared to be similar among the normal, mild and moderate renal function groups, but was 38% higher in the severe group than in the normal group. Caution should be exercised in renally impaired patients [see Use in Specific Populations (8.6)]. The starting dose of oxaliplatin injection should be reduced in patients with severe renal impairment [see Dosage and Administration (2.2)].

Drug - Drug Interactions

No pharmacokinetic interaction between 85 mg/m² of oxaliplatin injection and infusional 5-fluorouracil has been observed in patients treated every 2 weeks, but increases of 5-fluorouracil plasma concentrations by approximately 20% have been observed with doses of 130 mg/m² of oxaliplatin injection administered every 3 weeks. *In vitro*, platinum was not displaced from plasma proteins by the following medications: erythromycin, salicylate, sodium valproate, granisetron, and paclitaxel. *In vitro*, oxaliplatin is not metabolized by, nor does it inhibit, human cytochrome P450 isoenzymes. No P450-mediated drug-drug interactions are therefore anticipated in patients. Since platinum-containing species are eliminated primarily through the kidney, clearance of these products may be decreased by coadministration of potentially nephrotoxic compounds, although this has not been specifically studied.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Long-term animal studies have not been performed to evaluate the carcinogenic potential of oxaliplatin. Oxaliplatin was not mutagenic to bacteria (Ames test) but was mutagenic to mammalian cells *in vitro* (L5178Y mouse lymphoma assay). Oxaliplatin was clastogenic both *in vitro* (chromosome aberration in human lymphocytes) and *in vivo* (mouse bone marrow micronucleus assay).

In a fertility study, male rats were given oxaliplatin at 0, 0.5, 1, or 2 mg/kg/day for five days every 21 days for a total of three cycles prior to mating with females that received two cycles of oxaliplatin on the same schedule. A dose of 2 mg/kg/day (less than one-seventh the recommended human dose on a body surface area basis) did not affect pregnancy rate, but caused developmental mortality (increased early resorptions, decreased live fetuses, decreased live births) and delayed growth (decreased fetal weight).

Testicular damage, characterized by degeneration, hypoplasia, and atrophy, was observed in dogs administered oxaliplatin at 0.75 mg/kg/day x 5 days every 28 days for three cycles. A no effect level

was not identified. This daily dose is approximately one-sixth of the recommended human dose on a body surface area basis.

14 CLINICAL STUDIES

14.1 Combination Adjuvant Therapy with Oxaliplatin Injection and Infusional 5-fluorouracil/leucovorin in Patients with Colon Cancer

An international, multicenter, randomized study compared the efficacy and evaluated the safety of oxaliplatin injection in combination with an infusional schedule of 5-fluorouracil/leucovorin to infusional 5-fluorouracil/leucovorin

alone, in patients with stage II (Dukes' B2) or III (Dukes' C) colon cancer who had undergone complete resection of the primary tumor. The primary objective of the study was to compare the 3-year disease-free survival (DFS) in patients receiving oxaliplatin injection and infusional 5-fluorouracil/leucovorin to those receiving 5-fluorouracil/leucovorin

alone. Patients were to be treated for a total of 6 months (i.e., 12 cycles). A total of 2246 patients were randomized; 1123 patients per study arm. Patients in the study had to be between 18 and 75 years of age, have histologically proven stage II (T₃-T₄ N₀ M₀; Dukes' B2) or III (any T N₁₋₂ M₀; Dukes' C) colon carcinoma (with the inferior pole of the tumor above the peritoneal reflection, i.e., ≥15 cm from the anal margin) and undergone (within 7 weeks prior to randomization) complete resection of the primary tumor without gross or microscopic evidence of residual disease. Patients had to have had no prior chemotherapy, immunotherapy or radiotherapy, and have an ECOG performance status of 0, 1, or 2 (KPS ≥ 60%), absolute neutrophil count (ANC) > 1.5x10⁹/L, platelets ≥100x10⁹/L, serum creatinine ≤ 1.25 x ULN total bilirubin < 2 x ULN, AST/ALT < 2 x ULN and carcino-embryonic antigen (CEA) < 10 ng/mL. Patients with preexisting peripheral neuropathy (NCI grade ≥ 1) were ineligible for this trial.

The following table shows the dosing regimens for the two arms of the study.

Table 15 - Dosing Regimens in Adjuvant Therapy Study

| Treatment Arm | Dose | Regimen |
|---|--|-------------------------|
| Oxaliplatin injection + 5-FU/LV (FOLFOX4) (N =1123) 5-FU/LV (N=1123) | Day 1: Oxaliplatin Injection : 85 mg/m ² (2-hour infusion) + LV: 200 mg/m ² (2-hour infusion), followed by 5-FU: 400 mg/m ² (bolus), 600 mg/m ² (22-hour infusion) | every 2 weeks 12 cycles |
| | Day 2: LV: 200 mg/m ² (2-hour infusion), followed by 5-FU: 400 mg/m ² (bolus), 600 mg/m ² (22-hour infusion) | |
| | Day 1: LV: 200 mg/m ² (2-hour infusion), followed by 5-FU: 400 mg/m ² (bolus), 600 mg/m ² (22-hour infusion) | every 2 weeks 12 cycles |
| | Day 2: LV: 200 mg/m ² (2-hour infusion), followed by 5-FU: 400 mg/m ² (bolus), 600 mg/m ² (22-hour infusion) | |

The following tables show the baseline characteristics and dosing of the patient population entered into this study. The baseline characteristics were well balanced between arms.

Table 16 - Patient Characteristics in Adjuvant Therapy Study

| | Oxaliplatin Injection + Infusional 5-FU/LV N=1123 | Infusional 5-FU/LV N=1123 |
|--|--|------------------------------|
| Sex: Male (%) | 56.1 | 52.4 |
| Female (%) | 43.9 | 47.6 |
| Median age (years) | 61 | 60 |
| <65 years of age (%) | 64.4 | 66.2 |
| ≥65 years of age (%) | 35.6 | 33.8 |
| Karnofsky Performance Status (KPS) (%) | | |
| 100 | 29.7 | 30.5 |
| 90 | 52.2 | 53.9 |
| 80 | 4.4 | 3.3 |
| 70 | 13.2 | 11.9 |
| ≤60 | 0.6 | 0.4 |
| Primary site (%) | | |
| Colon including cecum | 54.6 | 54.4 |
| Sigmoid | 31.9 | 33.8 |
| Recto sigmoid | 12.9 | 10.9 |
| Other including rectum | 0.6 | 0.9 |
| Bowel obstruction (%) | | |
| Yes | 17.9 | 19.3 |
| Perforation (%) | | |
| Yes | 6.9 | 6.9 |
| Stage at Randomization (%) | | |
| II (T=3,4 N=0, M=0) | 40.1 | 39.9 |
| III (T=any, N=1,2, M=0) | 59.6 | 59.3 |
| IV (T=any, N=any, M=1) | 0.4 | 0.8 |
| Staging – T (%) | | |

| | | |
|-----------------|------|------|
| T1 | 0.5 | 0.7 |
| T2 | 4.5 | 4.8 |
| T3 | 76 | 75.9 |
| T4 | 19 | 18.5 |
| Staging – N (%) | | |
| N0 | 40.2 | 39.9 |
| N1 | 39.4 | 39.4 |
| N2 | 20.4 | 20.7 |
| Staging – M (%) | | |
| M1 | 0.4 | 0.8 |

Table 17 - Dosing in Adjuvant Therapy Study

| | Oxaliplatin Injection + Infusional 5-FU/LV N=1108 | Infusional 5-FU/LV N=1111 |
|--|--|--|
| Median Relative Dose Intensity (%) | | |
| 5-FU | 84.4 | 97.7 |
| Oxaliplatin Injection | 80.5 | N/A |
| Median Number of Cycles | 12 | 12 |
| Median Number of cycles with oxaliplatin injection | 11 | N/A |

The following table and figures summarize the disease-free survival (DFS) results in the overall randomized population and in patients with stage II and III disease based on an ITT analysis. The median duration of follow-up was approximately 77 months.

Table 18 - Summary of DFS analysis - ITT analysis

| Parameter | Oxaliplatin Injection + Infusional 5-FU/LV | Infusional 5-FU/LV |
|---|---|-------------------------------|
| Overall | | |
| N | 1123 | 1123 |
| Number of events – relapse or death (%) | 304 (27.1) | 360 (32.1) |
| Disease-free survival % [95% CI]* | 73.3 [70.7, 76] | 67.4 [64.6, 70.2] |
| Hazard ratio [95% CI]† | 0.8 [0.68, 0.93] | |
| Stratified Logrank test | p=0.003 | |
| Stage III (Dukes' C) | | |
| N | 672 | 675 |
| Number of events –relapse or death (%) | 226 (33.6) | 271 (40.1) |
| Disease-free survival % [95% CI]* | 66.4 [62.7, 70] | 58.9 [55.2, 62.7] |
| Hazard ratio [95% CI]† | 0.78 [0.65, 0.93] | |
| Logrank test | p=0.005 | |
| Stage II (Dukes' B2) | | |
| N | 451 | 448 |
| Number of events –relapse or death (%) | 78 (17.3) | 89 (19.9) |
| Disease-free survival % [95% CI]* | 83.7 [80.2, 87.1] | 79.9 [76.2, 83.7] |
| Hazard ratio [95% CI]† | 0.84 [0.62, 1.14] | |
| Logrank test | p=0.258 | |

Data cut off for disease free survival 1 June 2006

* Disease-free survival at 5 years

† A hazard ratio of less than 1 favors Oxaliplatin Injection + Infusional 5-fluorouracil/leucovorin

In the overall and stage III colon cancer populations DFS was statistically significantly improved in the oxaliplatin injection combination arm compared to infusional 5-fluorouracil/leucovorin alone. However, a statistically significant improvement in DFS was not noted in Stage II patients.

Figure 2 shows the DFS Kaplan-Meier curves for the comparison of oxaliplatin injection and infusional 5-fluorouracil/leucovorin

combination and infusional 5-fluorouracil/leucovorin

alone for the overall population (ITT analysis).

Figure 3 shows the DFS Kaplan-Meier curves for the comparison of oxaliplatin injection and infusional 5-fluorouracil/leucovorin

combination and infusional 5-fluorouracil/leucovorin

alone in Stage III patients.




Figure 2 - DFS Kaplan-Meier curves by treatment arm (cutoff: 1 June 2006) – ITT population




Figure 3 - DFS Kaplan-Meier curves by treatment arm in Stage III patients (cutoff: 1 June 2006) – ITT population

The following table summarizes the overall survival (OS) results in the overall randomized population and in patients with stage II and III disease, based on the ITT analysis.

Table 19 - Summary of OS analysis - ITT analysis

| Parameter | Oxaliplatin Injection +Infusional 5-FU/LV | Infusional 5-FU/LV |
|----------------------------|---|--------------------|
| Overall | | |
| N | 1123 | 1123 |
| Number of death events (%) | 245 (21.8) | 283 (25.2) |
| Hazard ratio* [95% CI] | 0.84 [0.71, 1] | |
| Stage III (Dukes' C) | | |
| N | 672 | 675 |
| Number of death events (%) | 182 (27.1) | 220 (32.6) |
| Hazard ratio* [95% CI] | 0.8 [0.65, 0.97] | |
| Stage II (Dukes' B2) | | |
| N | 451 | 448 |
| Number of death events (%) | 63 (14) | 63 (14.1) |
| Hazard ratio* [95% CI] | 1 [0.7, 1.41] | |

Data cut off for overall survival 16 January 2007

* A hazard ratio of less than 1 favors Oxaliplatin Injection + Infusional 5-fluorouracil/leucovorin

14.2 Combination Therapy with Oxaliplatin Injection and 5-fluorouracil/leucovorin in Patients Previously Untreated for Advanced Colorectal Cancer

A North American, multicenter, open-label, randomized controlled study was sponsored by the National Cancer Institute (NCI) as an intergroup study led by the North Central Cancer Treatment Group (NCCTG). The study had 7 arms at different times during its conduct, four of which were closed due to either changes in the standard of care, toxicity, or simplification. During the study, the control arm was changed to irinotecan plus 5-fluorouracil/leucovorin. The results reported below compared the efficacy and safety of two experimental regimens, oxaliplatin injection in combination with infusional 5-fluorouracil/leucovorin and a combination of oxaliplatin injection plus irinotecan, to an approved control regimen of irinotecan plus 5-fluorouracil/leucovorin in 795 concurrently randomized patients previously untreated for locally advanced or metastatic colorectal cancer. After completion of enrollment, the dose of irinotecan plus 5-fluorouracil/leucovorin was decreased due to toxicity. Patients had to be at least 18 years of age, have known locally advanced, locally recurrent, or metastatic colorectal adenocarcinoma not curable by surgery or amenable to radiation therapy with curative intent, histologically proven colorectal adenocarcinoma, measurable or evaluable disease, with an ECOG performance status 0, 1, or 2. Patients had to have granulocyte count $\geq 1.5 \times 10^9/L$, platelets $\geq 100 \times 10^9/L$, hemoglobin ≥ 9 gm/dL, creatinine $\leq 1.5 \times$ ULN, total bilirubin ≤ 1.5 mg/dL, AST $\leq 5 \times$ ULN, and alkaline phosphatase $\leq 5 \times$ ULN. Patients may have received adjuvant therapy for resected Stage II or III disease without recurrence within 12 months. The patients were stratified for ECOG performance status (0, 1 vs. 2), prior adjuvant chemotherapy (yes vs. no), prior immunotherapy (yes vs. no), and age (<65 vs. ≥ 65 years). Although no post study treatment was specified in the protocol, 65 to 72% of patients received additional post study chemotherapy after study treatment discontinuation on all arms. Fifty-eight percent of patients on the oxaliplatin injection plus 5-fluorouracil/leucovorin arm received an irinotecan-containing regimen and 23% of patients on the irinotecan plus 5-fluorouracil/leucovorin arm received oxaliplatin-containing regimens. Oxaliplatin was not commercially available during the trial.

The following table presents the dosing regimens of the three arms of the study.

Table 20 – Dosing Regimens in Patients Previously Untreated for Advanced Colorectal Cancer Clinical Trial

| Treatment Arm | Dose | Regimen |
|---|--|---------------|
| Oxaliplatin Injection + 5-FU/LV (FOLFOX4) (N=267) | Day 1: Oxaliplatin Injection: 85 mg/m ² (2-hour infusion) + LV 200 mg/m ² (2-hour infusion), followed by 5-FU: 400 mg/m ² (bolus), 600 mg/m ² (22-hour infusion) Day 2: LV 200 mg/m ² (2-hour infusion), followed by 5-FU: 400 mg/m ² (bolus), 600 mg/m ² (22-hour infusion) | every 2 weeks |
| Irinotecan + 5-FU/LV (IFL) (N=264) | Day 1: irinotecan 125 mg/m ² as a 90-min infusion + LV 20 mg/m ² as a 15-min infusion or intravenous push, followed by 5-FU 500 mg/m ² intravenous bolus weekly x 4 | every 6 weeks |
| Oxaliplatin Injection + Irinotecan (IROX) (N=264) | Day 1: oxaliplatin injection: 85 mg/m ² intravenous (2-hour infusion) + irinotecan 200 mg/m ² intravenous over 30 minutes | every 3 weeks |

The following table presents the demographics of the patient population entered into this study.

Table 21 – Patient Demographics in Patients Previously Untreated for Advanced Colorectal Cancer Clinical Trial

| | Oxaliplatin Injection + 5-FU/LV N=267 | Irinotecan + 5-FU/LV N=264 | Oxaliplatin Injection + Irinotecan N=264 |
|-------------------------------|--|-------------------------------|---|
| Sex: Male (%) | 58.8 | 65.2 | 61 |
| Female (%) | 41.2 | 34.8 | 39 |
| Median age (years) | 61 | 61 | 61 |
| <65 years of age (%) | 61 | 62 | 63 |
| ≥ 65 years of age (%) | 39 | 38 | 37 |
| ECOG (%) | | | |
| 0 to 1 | 94.4 | 95.5 | 94.7 |
| 2 | 5.6 | 4.5 | 5.3 |
| Involved organs (%) | | | |
| Colon only | 0.7 | 0.8 | 0.4 |
| Liver only | 39.3 | 44.3 | 39 |
| Liver + other | 41.2 | 38.6 | 40.9 |
| Lung only | 6.4 | 3.8 | 5.3 |
| Other (including lymph nodes) | 11.6 | 11 | 12.9 |
| Not reported | 0.7 | 1.5 | 1.5 |
| Prior radiation (%) | 3 | 1.5 | 3 |
| Prior surgery (%) | 74.5 | 79.2 | 81.8 |

| | | | |
|--------------------|------|------|------|
| Prior adjuvant (%) | 15.7 | 14.8 | 15.2 |
|--------------------|------|------|------|

The length of a treatment cycle was 2 weeks for the oxaliplatin injection and 5-fluorouracil/leucovorin regimen; 6 weeks for the irinotecan plus 5-fluorouracil/leucovorin regimen; and 3 weeks for the oxaliplatin injection plus irinotecan regimen. The median number of cycles administered per patient was 10 (23.9 weeks) for the oxaliplatin injection and 5-fluorouracil/leucovorin regimen, 4 (23.6 weeks) for the irinotecan plus 5-fluorouracil/leucovorin regimen, and 7 (21 weeks) for the oxaliplatin injection plus irinotecan regimen. Patients treated with the oxaliplatin injection and 5-fluorouracil/leucovorin combination had a significantly longer time to tumor progression based on investigator assessment, longer overall survival, and a significantly higher confirmed response rate based on investigator assessment compared to patients given irinotecan plus 5-fluorouracil/leucovorin. The following table summarizes the efficacy results.

Table 22 – Summary of Efficacy


| | Oxaliplatin Injection + 5-FU/LV N=267 | Irinotecan + 5-FU/LV N=264 | Oxaliplatin Injection + Irinotecan N=264 |
|---|--|-------------------------------|---|
| Survival (ITT) | | | |
| Number of deaths N (%) | 155 (58.1) | 192 (72.7) | 175 (66.3) |
| Median survival (months) | 19.4 | 14.6 | 17.6 |
| Hazard Ratio and (95% confidence interval) | 0.65 (0.53 to 0.8)* | | |
| P-value | <0.0001* | - | - |
| TTP (ITT, investigator assessment) | | | |
| Percentage of progressors | 82.8 | 81.8 | 89.4 |
| Median TTP (months) | 8.7 | 6.9 | 6.5 |
| Hazard Ratio and (95% confidence interval) [†] | 0.74 (0.61 to 0.89)* | | |
| P-value | 0.0014* | - | - |
| Response Rate (investigator assessment) [‡] | | | |
| Patients with measurable disease | 210 | 212 | 215 |
| Complete response N (%) | 13 (6.2) | 5 (2.4) | 7 (3.3) |
| Partial response N (%) | 82 (39) | 64 (30.2) | 67 (31.2) |
| Complete and partial response N (%) | 95 (45.2) | 69 (32.5) | 74 (34.4) |
| 95% confidence interval | (38.5 to 52) | (26.2 to 38.9) | (28.1 to 40.8) |
| P-value | 0.008* | - | - |
| The numbers in the response rate and TTP analysis are based on unblinded investigator assessment. | | | |

* Compared to irinotecan plus 5-fluorouracil/leucovorin (IFL) arm

† A hazard ratio of less than 1 favors oxaliplatin injection + Irinotecan plus 5-fluorouracil/leucovorin

‡ Based on all patients with measurable disease at baseline

Figure 4 illustrates the Kaplan-Meier survival curves for the comparison of oxaliplatin injection and 5-fluorouracil/leucovorin combination and oxaliplatin injection plus irinotecan to irinotecan plus 5-fluorouracil/leucovorin.



* Log rank test comparing oxaliplatin injection plus 5-FU/LV to irinotecan plus 5-FU/LV.

Figure 4- Kaplan-Meier Overall Survival by treatment arm

A descriptive subgroup analysis demonstrated that the improvement in survival for oxaliplatin injection plus 5-fluorouracil/leucovorin compared to irinotecan plus 5-fluorouracil/leucovorin appeared to be maintained across age groups, prior adjuvant therapy, and number of organs involved. An estimated survival advantage in oxaliplatin injection plus 5-fluorouracil/leucovorin versus irinotecan plus 5-fluorouracil/leucovorin was seen in both genders; however it was greater among women than men. Insufficient subgroup sizes prevented analysis by race.

14.3 Combination Therapy with Oxaliplatin Injection and 5-fluorouracil/leucovorin in Previously

Treated Patients with Advanced Colorectal Cancer

A multicenter, open-label, randomized, three-arm controlled study was conducted in the US and Canada comparing the efficacy and safety of oxaliplatin injection in combination with an infusional schedule of 5-fluorouracil/leucovorin

to the same dose and schedule of 5-fluorouracil/leucovorin

alone and to single agent oxaliplatin in patients with advanced colorectal cancer who had relapsed/progressed during or within 6 months of first-line therapy with bolus 5-fluorouracil/leucovorin

and irinotecan. The study was intended to be analyzed for response rate after 450 patients were enrolled. Survival will be subsequently assessed in all patients enrolled in the completed study. Accrual to this study is complete, with 821 patients enrolled. Patients in the study had to be at least 18 years of age, have unresectable, measurable, histologically proven colorectal adenocarcinoma, with a Karnofsky performance status >50%. Patients had to have SGOT (AST) and SGPT (ALT) $\leq 2x$ the institution's upper limit of normal (ULN), unless liver metastases were present and documented at baseline by CT or MRI scan, in which case $\leq 5x$ ULN was permitted. Patients had to have alkaline phosphatase $\leq 2x$ the institution's ULN, unless liver metastases were present and documented at baseline by CT or MRI scan, in which cases $\leq 5x$ ULN was permitted. Prior radiotherapy was permitted if it had been completed at least 3 weeks before randomization.

The dosing regimens of the three arms of the study are presented in the table below.

Table 23 – Dosing Regimens in Refractory and Relapsed Colorectal Cancer Clinical Trial

| Treatment Arm | Dose | Regimen |
|--|--|---------------|
| Oxaliplatin Injection + 5-FU/LV (N =152) | Day 1: Oxaliplatin Injection: 85 mg/m ² (2-hour infusion) + LV 200 mg/m ² (2-hour infusion), followed by 5-FU: 400 mg/m ² (bolus), 600 mg/m ² (22-hour infusion) Day 2: LV 200 mg/m ² (2-hour infusion), followed by 5-FU: 400 mg/m ² (bolus), 600 mg/m ² (22-hour infusion) | every 2 weeks |
| 5-FU/LV (N=151) | Day 1: LV 200 mg/m ² (2-hour infusion), followed by 5-FU: 400 mg/m ² (bolus), 600 mg/m ² (22-hour infusion) Day 2: LV 200 mg/m ² (2-hour infusion), followed by 5-FU: 400 mg/m ² (bolus), 600 mg/m ² (22-hour infusion) | every 2 weeks |
| Oxaliplatin Injection (N=156) | Day 1: Oxaliplatin Injection 85 mg/m ² (2-hour infusion) | every 2 weeks |

Patients entered into the study for evaluation of response must have had at least one unidimensional lesion measuring ≥ 20 mm using conventional CT or MRI scans, or ≥ 10 mm using a spiral CT scan. Tumor response and progression were assessed every 3 cycles (6 weeks) using the Response Evaluation Criteria in Solid Tumors (RECIST) until radiological documentation of progression or for 13 months following the first dose of study drug(s), whichever came first. Confirmed responses were based on two tumor assessments separated by at least 4 weeks.

The demographics of the patient population entered into this study are shown in the table below.

Table 24 – Patient Demographics in Refractory and Relapsed Colorectal Cancer Clinical Trial

| | 5-FU/LV (N = 151) | Oxaliplatin Injection (N = 156) | Oxaliplatin Injection + 5-FU/LV (N = 152) |
|--------------------------------|-------------------|---------------------------------|---|
| Sex: Male (%) | 54.3 | 60.9 | 57.2 |
| Female (%) | 45.7 | 39.1 | 42.8 |
| Median age (years) | 60 | 61 | 59 |
| Range | 21 to 80 | 27 to 79 | 22 to 88 |
| Race (%) | | | |
| Caucasian | 87.4 | 84.6 | 88.8 |
| Black | 7.9 | 7.1 | 5.9 |
| Asian | 1.3 | 2.6 | 2.6 |
| Other | 3.3 | 5.8 | 2.6 |
| KPS (%) | | | |
| 70 to 100 | 94.7 | 92.3 | 95.4 |
| 50 to 60 | 2.6 | 4.5 | 2 |
| Not reported | 2.6 | 3.2 | 2.6 |
| Prior radiotherapy (%) | 25.2 | 19.2 | 25 |
| Prior pelvic radiation (%) | 18.5 | 13.5 | 21.1 |
| Number of metastatic sites (%) | | | |
| 1 | 27.2 | 31.4 | 25.7 |
| ≥ 2 | 72.2 | 67.9 | 74.3 |
| Liver involvement (%) | | | |
| Liver only | 22.5 | 25.6 | 18.4 |
| Liver + other | 60.3 | 59 | 53.3 |

The median number of cycles administered per patient was 6 for the oxaliplatin injection and 5-fluorouracil/leucovorin combination and 3 each for 5-fluorouracil/leucovorin alone and oxaliplatin injection alone.

Patients treated with the combination of oxaliplatin injection and 5-fluorouracil/leucovorin had an increased response rate compared to patients given 5-fluorouracil/leucovorin or oxaliplatin alone. The efficacy results are summarized in the tables below.

Table 25 - Response Rates (ITT Analysis)

| Best Response | 5-FU/LV (N=151) | Oxaliplatin Injection (N=156) | Oxaliplatin Injection + 5-FU/LV (N=152) |
|---------------|--|-------------------------------|---|
| CR | 0 | 0 | 0 |
| PR | 0 | 2 (1%) | 13 (9%) |
| p-value | 0.0002 for 5-FU/LV vs. Oxaliplatin Injection + 5-FU/LV | | |
| 95%CI | 0 to 2.4% | 0.2 to 4.6% | 4.6 to 14.2% |

Table 26 - Summary of Radiographic Time to Progression*

| Arm | 5-FU/LV (N=151) | Oxaliplatin Injection (N=156) | Oxaliplatin Injection + 5-FU/LV (N=152) |
|---|-----------------|-------------------------------|---|
| No. of Progressors | 74 | 101 | 50 |
| No. of patients with no radiological evaluation beyond baseline | 22 (15%) | 16 (10%) | 17 (11%) |
| Median TTP (months) | 2.7 | 1.6 | 4.6 |
| 95% CI | 1.8 to 3 | 1.4 to 2.7 | 4.2 to 6.1 |

* This is not an ITT analysis. Events were limited to radiographic disease progression documented by independent review of radiographs. Clinical progression was not included in this analysis, and 18% of patients were excluded from the analysis based on unavailability of the radiographs for independent review.

At the time of the interim analysis 49% of the radiographic progression events had occurred. In this interim analysis an estimated 2-month increase in median time to radiographic progression was observed compared to 5-fluorouracil/leucovorin alone.

Of the 13 patients who had tumor response to the combination of oxaliplatin injection and 5-fluorouracil/leucovorin, 5 were female and 8 were male, and responders included patients <65 years old and ≥65 years old. The small number of non-Caucasian participants made efficacy analyses in these populations uninterpretable.

15 REFERENCES

1. NIOSH Alert: Preventing occupational exposures to antineoplastic and other hazardous drugs in healthcare settings. 2004. U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Institute for Occupational Safety and Health, DHHS (NIOSH) Publication No. 2004-165.
2. OSHA Technical Manual, TED 1-0.15A, Section VI: Chapter 2. Controlling Occupational Exposure to Hazardous Drugs. OSHA, 1999. http://www.osha.gov/dts/osta/otm/otm_vi/otm_vi_2.html
3. American Society of Health-System Pharmacists. (2006) ASHP Guidelines on Handling Hazardous Drugs.
4. Polovich, M., White, J. M., & Kelleher, L.O. (eds.) 2005. Chemotherapy and biotherapy guidelines and recommendations for practice (2nd. ed.) Pittsburgh, PA: Oncology Nursing Society.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Oxaliplatin injection, USP is supplied in clear, glass, single-use vials with gray elastomeric stoppers and aluminum flip-off seals containing 50 mg/10 mL or 100 mg/20 mL of oxaliplatin, USP as a sterile, preservative-free, aqueous solution at a concentration of 5 mg/mL. Water for Injection, USP is present as an inactive ingredient.

NDC 47335-046-40: 50 mg/10 mL single-use vial with parrot green flip-off seal individually packaged in a carton.

NDC 47335-047-40: 100 mg/20 mL single-use vial with golden brown flip-off seal individually packaged in a carton.

16.2 Storage

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° and 30°C (59° and 86°F) [see USP Controlled Room Temperature]. Do not freeze and protect from light (keep in original outer carton).

16.3 Handling and Disposal

As with other potentially toxic anticancer agents, care should be exercised in the handling and preparation of infusion solutions prepared from oxaliplatin injection. The use of gloves is recommended. If a solution of oxaliplatin injection contacts the skin, wash the skin immediately and thoroughly with soap and water. If oxaliplatin injection contacts the mucous membranes, flush thoroughly with water.

Procedures for the handling and disposal of anticancer drugs should be considered. Several guidelines on the subject have been published [see *References (15)*]. There is no general agreement that all of the procedures recommended in the guidelines are necessary or appropriate.

17 PATIENT COUNSELING INFORMATION

Advise patients:

- To expect side effects of oxaliplatin injection, particularly its neurologic effects, both the acute, reversible effects and the persistent neurosensory toxicity. Patients should be informed that the acute neurosensory toxicity may be precipitated or exacerbated by exposure to cold or cold objects.
- To avoid cold drinks, use of ice, and should cover exposed skin prior to exposure to cold temperature or cold objects.
- Of the risk of low blood cell counts and to contact their physician immediately should fever, particularly if associated with persistent diarrhea, or evidence of infection develop.
- To contact their physician if persistent vomiting, diarrhea, signs of dehydration, cough or breathing difficulties occur, or signs of allergic reaction appear.
- To exercise caution when driving and using machines. No studies on the effects of the ability to operate cars and machines have been performed; however, oxaliplatin treatment resulting in an increase risk of dizziness, nausea and vomiting, and other neurologic symptoms that affect gait and balance may lead to a minor or moderate influence on the ability to drive and use machines.
- Of the potential effects of vision abnormalities, in particular transient vision loss (reversible following therapy discontinuation), which may affect patients' ability to drive and use machines.

FDA-Approved Patient Labeling

Patient Information

Oxaliplatin(ox-Al-ah-platin)Injection, USP

for intravenous use

Read this Patient Information leaflet carefully before you start receiving oxaliplatin injection. There may be new information. It will help you learn more about oxaliplatin injection. This leaflet does not take the place of talking to your doctor about your medical condition or your treatment. Ask your doctor about any questions you have.

What is the most important information I should know about oxaliplatin injection?

Oxaliplatin can cause serious allergic reactions, including allergic reactions that can lead to death.

Oxaliplatin injection is a platinum base medicine. Serious allergic reactions including death can happen in people who take oxaliplatin injection and who have had previous allergic reactions to platinum medicines. Serious allergic reactions can happen within a few minutes of your infusion or any time during your treatment with oxaliplatin injection.

Get emergency help right away if you:

- **have trouble breathing.**
- **feel like your throat is closing up.**

Call your doctor right away if you have any of the following signs or symptoms of an allergic reaction:

- rash
- flushed face
- hives
- itching
- swelling of your lips or tongue
- sudden cough
- dizziness or feel faint
- sweating
- chest pain

See “**What are the possible side effects of oxaliplatin injection?**” for information about other serious side effects.

What is oxaliplatin injection?

Oxaliplatin injection is an anti-cancer (chemotherapy) medicine that is used with other anti-cancer medicines called 5-fluorouracil and leucovorin to treat people with:

- stage III colon cancer after surgery to remove the tumor
- advanced colon or rectal cancer (colorectal cancer).

It is not known if oxaliplatin injection is effective in children.

Who should not receive oxaliplatin injection?

Do not receive oxaliplatin injection if you are allergic to any of the ingredients in oxaliplatin injection or other medicines that contain platinum. See the end of this leaflet for a complete list of the ingredients oxaliplatin injection.

Ask your doctor if you are not sure if you take a medicine that contains platinum.

What should I tell my doctor before receiving oxaliplatin injection?

Before receiving oxaliplatin injection, tell your doctor about all of your medical conditions, including if you:

- have an infection
- have lung, liver, or kidney problems
- have or had heart problems such as an abnormal heart test called an electrocardiogram (ECG or EKG), a condition called long QT syndrome, an irregular or slow heartbeat, or a family history of heart problems.
- have had changes in the level of certain blood salt (electrolytes) levels, including potassium, magnesium, and calcium
- are pregnant or plan to become pregnant. Oxaliplatin injection may harm your unborn baby. Females who are able to become pregnant should avoid becoming pregnant and should use effective birth control during treatment with oxaliplatin injection.
- are breastfeeding or plan to breastfeed. It is not known if oxaliplatin passes into your breast milk. You and your doctor should decide if you will receive oxaliplatin injection or breastfeed. You should not do both.

Tell your doctor about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

Know the medicines you take. Keep a list of them and show it to your doctor and pharmacist when you get a new medicine.

How will I receive oxaliplatin injection?

- Oxaliplatin injection is given to you into your vein through an intravenous (IV) tube.
- Your doctor will prescribe oxaliplatin injection in a dose that is right for you.
- Your doctor may change how often you receive oxaliplatin injection, your dose, or how long your infusion will take.
- You and your doctor will decide how many oxaliplatin injection treatments you will receive.
- It is very important that you do exactly what your doctor and nurse tell you to do.
- Some medicines may be given to you before oxaliplatin injection to help prevent nausea and vomiting.
- Each treatment course is given to you over 2 days. You will receive oxaliplatin injection on the first day only.
- There are usually 14 days between each chemotherapy treatment course.
- It is important for you to keep all of your medical appointments. Call your doctor if you miss an appointment. There may be special instructions for you.

Treatment Day 1:

- Oxaliplatin injection and leucovorin will be given through a thin plastic tube into a vein (intravenous infusion or IV) and given for 2 hours. You will be watched by a healthcare provider during this time.
- Right after the oxaliplatin injection and leucovorin are given, 2 doses of 5-fluorouracil will be given. The first dose is given right away into your IV tube. The second dose will be given into your IV tube over the next 22 hours, using a pump device.

Treatment Day 2:

You **will not** get oxaliplatin injection on Day 2. Leucovorin and 5-fluorouracil will be given the same way as on Day 1.

The 5-fluorouracil will be given through your IV with a pump. If you have any problems with the pump or the tube, call your doctor, your nurse, or the person who is responsible for your pump. Do not let anyone other than a healthcare provider touch your infusion pump or tubing.

What should I avoid while receiving oxaliplatin injection?

- Avoid cold temperatures and cold objects. Cover your skin if you go outdoors in cold temperatures.
- Do not drink cold drinks or use ice cubes in drinks.
- Do not put ice or ice packs on your body.
- Oxaliplatin injection can cause dizziness, vision problems, or vision loss that can affect your ability to drive or use machines. You should not drive or operate machinery if you develop these symptoms while receiving oxaliplatin injection.

See “How can I reduce the side effects caused by cold temperatures?” for more information.

Talk with your doctor and nurse about your level of activity during treatment with oxaliplatin injection. Follow their instructions.

What are the possible side effects of oxaliplatin injection?

Oxaliplatin injection can cause serious side effects, including:

- See “What is the most important information I should know about oxaliplatin injection?”

- **Nerve problems.** Oxaliplatin injection can affect how your nerves work and make you feel. Nerve problems may happen with the first treatment or within two days after your treatment of oxaliplatin injection. Nerve problems may last a short time (acute) or may become persistent. Symptoms may improve after stopping treatment with oxaliplatin injection. Exposure to cold or cold objects may cause or worsen nerve problems. Tell your doctor right away if you get any signs of nerve problems, including:

- o very sensitive to cold temperatures and cold objects

- o trouble breathing, swallowing, or saying words, jaw tightness, odd feelings in your tongue, or chest pressure

- o pain, tingling, burning (pins and needles, numb feeling) in your hands, feet, or around your mouth or throat, which may cause problems walking or performing activities of daily living.

For information on ways to lessen or help with the nerve problems, see the end of this leaflet, “How can I reduce the side effects caused by cold temperatures?”

- **Reversible Posterior Leukoencephalopathy (RPLS).** RPLS is a rare condition that affects the brain.

Tell your doctor right away if you have any of the following signs and symptoms of RPLS:

- o headache

- o confusion or a change in the way you think

- o seizures

- o vision problems, such as blurriness or vision loss

- **Low white blood cell counts (neutropenia).** Oxaliplatin injection can cause low white blood cells counts. Low blood cell counts are common with oxaliplatin injection and can lead to serious infection and death. Tell your doctor right away if you have a fever greater than 100.9°F (38.3°C) or a prolonged fever greater than 100.4°F (38°C) for more than one hour (febrile neutropenia). Call your doctor right away if you get any of the following signs of infection:

- o chills or shivering

- o pain on swallowing

- o sore throat

- o cough that brings up mucus

- o burning or pain on urination

- o redness or swelling at intravenous site

- o persistent diarrhea

- **Lung problems (interstitial fibrosis).** Oxaliplatin injection can cause lung problems that may lead to death. Tell your doctor right away if you get a dry cough and have trouble breathing (shortness of breath) before your next treatment. These may be signs of a serious lung disease.

- **Liver problems (hepatotoxicity).** Your doctor will do blood tests to check your liver.

- **Heart problems.** Oxaliplatin injection can cause heart problems that have led to death. Your doctor may do blood and heart tests during treatment with oxaliplatin injection if you have certain heart problem. If you faint (lose consciousness) or have an irregular heartbeat or chest pain during treatment with oxaliplatin injection, tell your doctor right away as this may be a sign of a serious heart condition.

- **Muscle problems.** Oxaliplatin injection can cause muscle damage (rhabdomyolysis) which can lead to death. Tell your doctor right away if you have muscle pain and swelling, along with weakness, fever, or red- brown urine.

- **Harm to an unborn baby.** See “What should I tell my doctor before receiving oxaliplatin injection?”

The most common side effects of oxaliplatin injection include:

- Numbness, pain, tingling, and/or burning along the nerves
- Low white blood cells (neutropenia)
- Low platelet count (important for clotting and to control bleeding)
- Low red blood cells (blood cells that carry oxygen to the tissues)
- Nausea
- Changes in liver function tests
- Diarrhea
- Vomiting
- Tiredness
- Mouth sores

Tell your doctor if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of oxaliplatin injection. For more information, ask your doctor or

pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How can I reduce the side effects caused by cold temperatures?

- Cover yourself with a blanket while you are getting your oxaliplatin injection infusion.
- Do not breathe deeply when exposed to cold air.
- Wear warm clothing in cold weather at all times. Cover your mouth and nose with a scarf or a pull-down cap (ski cap) to warm the air that goes to your lungs.
- Wear gloves when taking things from the freezer or refrigerator.
- Drink fluids warm or at room temperature.
- Always drink through a straw.
- **Do not** use ice chips if you have nausea or mouth sores. Ask your doctor about what you can use.
- Be aware that most metals are cold to touch, especially in the winter. These include your car door and mailbox. Wear gloves to touch cold objects.
- Do not run the air-conditioning at high levels in the house or in the car in hot weather.
- If your body gets cold, warm-up the affected part. If your hands get cold, wash them with warm water.
- Always let your doctor know before your next treatment how well you did since your last visit.

Your doctor may have other useful tips for helping you with side effects.

General information about the safe and effective use of oxaliplatin injection

Medicines are sometimes prescribed for purposes other than those listed in the Patient Information leaflet.

This Patient Information leaflet summarizes the most important information about oxaliplatin injection. If you would like more information, talk with your doctor. You can ask your doctor or pharmacist for information about oxaliplatin injection that is written for health professionals.

What are the ingredients in oxaliplatin injection?

Active ingredient: oxaliplatin

Inactive ingredients: water for injection

Distributed by:

Sun Pharmaceutical Industries, Inc.
Cranbury, NJ 08512

Manufactured by:

Sun Pharmaceutical Industries Ltd.
Halol-Baroda Highway,
Halol-389 350, Gujarat, India.
Under license from Sanofi-Aventis U.S. LLC,
Sanofi-Aventis, and Debiopharm S.A.
of U.S. Pat Nos. 5,420,319; 5,959,133; and 5,716,988

ISS. 10/2015
PJPI0332B

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - LABEL - 50MG

NDC 47335-046-40

Oxaliplatin Injection, USP

50 mg/10 mL (5 mg/mL)

Cytotoxic Agent

FOR INTRAVENOUS USE ONLY

Single Use Vial

Sterile - Preservative Free

See package insert for further required dilution

DO NOT MIX OR ADD TO SODIUM CHLORIDE/CHLORIDE-CONTAINING SOLUTIONS

Rx only

SUN PHARMA

Each mL contains: oxaliplatin, USP 5 mg
Inactive: Water for Injection 1 mL

USUAL DOSAGE: Further dilute in 5% dextrose injection, USP. See insert.

Do not freeze; protect from light.

Retain in carton until time of use.

Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° and 30°C (59° and 86°F) [see USP Controlled Room Temperature].

Discard unused portion (see package insert for storage of further diluted solutions.)

Distributed by:
Sun Pharmaceutical Industries, Inc.
Cranbury, NJ 08512

NDC 47335-046-40

Oxaliplatin Injection, USP


50 mg/10 mL (5 mg/mL)

Cytotoxic Agent
FOR INTRAVENOUS USE ONLY
Single use vial

Sterile – Preservative Free
See package insert for further required dilution

DO NOT MIX OR ADD TO SODIUM CHLORIDE/CHLORIDE-CONTAINING SOLUTIONS

Rx only


SUN PHARMA

Manufactured by:
Sun Pharmaceutical Industries Ltd.
Halol-Baroda Highway,
Halol-389 350, Gujarat, India.
Under license from Sanofi-Aventis U.S. LLC,
Sanofi-Aventis, and Debiopharm S.A.
of U.S. Pat Nos. 5,420,319; 5,959,133;
and 5,716,988

PJLB1800A PJLB1800A PJLB1800A
PJLB1800A ISS. 02/2015

GUJ/DRUGS/28/396

Batch No.:
Exp.:

4 7 3 3 5 0 4 6 4 0 3

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - CARTON - 50MG

NDC 47335-046-40
Oxaliplatin Injection, USP
50 mg/10 mL (5 mg/mL)
Cytotoxic Agent
FOR INTRAVENOUS USE ONLY
Single Use Vial Only
Sterile - Preservative Free
See package insert for further required dilution
DO NOT MIX OR ADD TO SODIUM CHLORIDE/CHLORIDE-CONTAINING SOLUTIONS
Rx only
10 mL Single-Use Vial
SUN PHARMA

NDC 47335-046-40

Oxaliplatin Injection, USP


50 mg/10 mL (5 mg/mL)

Cytotoxic Agent
FOR INTRAVENOUS USE ONLY
Single use vial only

Sterile – Preservative Free
See package insert for further required dilution

DO NOT MIX OR ADD TO SODIUM CHLORIDE/CHLORIDE-CONTAINING SOLUTIONS

Rx only
10 mL Single-Use Vial


SUN PHARMA

Manufactured by:
Sun Pharmaceutical Industries Ltd.
Halol-Baroda Highway,
Halol-389 350, Gujarat, India.
Under license from Sanofi-Aventis U.S. LLC,
Sanofi-Aventis, and Debiopharm S.A.
of U.S. Pat Nos. 5,420,319; 5,959,133; and 5,716,988

PJSB0667A ISS. 02/2015
GUJ/DRUGS/28/396

4 7 3 3 5 0 4 6 4 0 3

Batch No. / Exp. } To be over print

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - LABEL - 100MG

NDC 47335-047-40
Oxaliplatin Injection, USP
100 mg/20 mL (5 mg/mL)
Cytotoxic Agent
FOR INTRAVENOUS USE ONLY
Single Use Vial
Sterile - Preservative Free
See package insert for further required dilution
DO NOT MIX OR ADD TO SODIUM CHLORIDE/CHLORIDE-CONTAINING SOLUTIONS
Rx only
SUN PHARMA

Each mL contains: oxaliplatin, USP 5 mg
 Inactive: Water for Injection 1 mL
USUAL DOSAGE: Further dilute in 5% dextrose injection, USP. See insert.
Do not freeze; protect from light.
Retain in carton until time of use.
 Store at 20° to 25°C (68° to 77°F); excursions permitted between 15° and 30°C (59° and 86°F) [see USP Controlled Room Temperature].
 Discard unused portion (see package insert for storage of further diluted solutions.)

Distributed by:
Sun Pharmaceutical Industries, Inc.
 Cranbury, NJ 08512

NDC 47335-047-40

Oxaliplatin Injection, USP

100 mg/20 mL (5 mg/mL)

Cytotoxic Agent
FOR INTRAVENOUS USE ONLY
Single use vial

Sterile – Preservative Free
 See package insert for further required dilution

DO NOT MIX OR ADD TO SODIUM CHLORIDE/CHLORIDE-CONTAINING SOLUTIONS

Rx only




Manufactured by:
Sun Pharmaceutical Industries Ltd.
 Halol-Baroda Highway,
 Halol-389 350, Gujarat, India.
 Under license from Sanofi-Aventis U.S. LLC,
 Sanofi-Aventis, and Debiopharm S.A.
 of U.S. Pat Nos. 5,420,319; 5,959,133;
 and 5,716,988

PJLB1801A PJLB1801A PJLB1801a
 PJLB1801A ISS. 02/2015
 GUJ/DRUGS/28/396

Batch No.:
 Exp.:

PACKAGE LABEL.PRINCIPAL DISPLAY PANEL - CARTON - 100MG

NDC 47335-047-40
Oxaliplatin Injection, USP
100 mg/20 mL (5 mg/mL)
Cytotoxic Agent
FOR INTRAVENOUS USE ONLY
Single Use Vial Only
Sterile - Preservative Free
See package insert for further required dilution
DO NOT MIX OR ADD TO SODIUM CHLORIDE/CHLORIDE-CONTAINING SOLUTIONS
Rx only
20 mL Single-Use Vial
SUN PHARMA

| OXALIPLATIN | | | |
|--|--------------------------|---------------------------|---------------|
| oxaliplatin injection, solution, concentrate | | | |
| Product Information | | | |
| Product Type | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:47335-046 |
| Route of Administration | INTRAVENOUS | | |
| Active Ingredient/Active Moiety | | | |
| Ingredient Name | Basis of Strength | Strength | |
| OXALIPLATIN (UNII: 04ZR38536J) (OXALIPLATIN - UNII:04ZR38536J) | OXALIPLATIN | 50 mg in 10 mL | |
| Inactive Ingredients | | | |
| Ingredient Name | Strength | | |
| WATER (UNII: 059QF0K00R) | | | |

| Packaging | | | | |
|-----------|------------------|--|----------------------|--------------------|
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:47335-046-40 | 1 in 1 CARTON | 04/09/2014 | |
| 1 | | 10 mL in 1 VIAL, SINGLE-USE; Type 0: Not a Combination Product | | |

| Marketing Information | | | |
|-----------------------|--|----------------------|--------------------|
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
| ANDA | ANDA202922 | 04/09/2014 | |

| OXALIPLATIN | | | | |
|--|--|--|----------------------|--------------------|
| oxaliplatin injection, solution, concentrate | | | | |
| Product Information | | | | |
| Product Type | HUMAN PRESCRIPTION DRUG | Item Code (Source) | NDC:47335-047 | |
| Route of Administration | INTRAVENOUS | | | |
| Active Ingredient/Active Moiety | | | | |
| Ingredient Name | Basis of Strength | Strength | | |
| OXALIPLATIN (UNII: 04ZR38536J) (OXALIPLATIN - UNII:04ZR38536J) | OXALIPLATIN | 100 mg in 20 mL | | |
| Inactive Ingredients | | | | |
| Ingredient Name | Strength | | | |
| WATER (UNII: 059QF0K00R) | | | | |
| Packaging | | | | |
| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
| 1 | NDC:47335-047-40 | 1 in 1 CARTON | 04/09/2014 | |
| 1 | | 20 mL in 1 VIAL, SINGLE-USE; Type 0: Not a Combination Product | | |
| Marketing Information | | | | |
| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date | |
| ANDA | ANDA202922 | 04/09/2014 | | |

Labeler - Sun Pharmaceutical Industries, Inc. (146974886)

| Establishment | | | |
|---------------------------------------|---------|-----------|--|
| Name | Address | ID/FEI | Business Operations |
| Sun Pharmaceutical Industries Limited | | 725959238 | ANALYSIS(47335-046, 47335-047) , MANUFACTURE(47335-046, 47335-047) |

Revised: 10/2018

Sun Pharmaceutical Industries, Inc.