NICOTINE- nicotine polacrilex gum, chewing Lucy Goods, Inc.

Nicotine Polacrilex Gum Coated Ice Mint 4 mg ANDA204833 - Equate

Drug Facts

Active ingredient (in each chewing piece)

Nicotine Polacrilex 4 mg (nicotine)

Purpose

Stop smoking aid

Use

reduces withdrawal symptoms, including nicotine craving, associated with quitting smoking

Warnings

If you are pregnant or breast-feeding, only use this medicine on the advice of your health care provider. Smoking can seriously harm your child. Try to stop smoking without using any nicotine replacement medicine. This medicine is believed to be safer than smoking. However, the risks to your child from this medicine are not fully known.

Ask a doctor before use if you have

- a sodium-restricted diet
- heart disease, recent heart attack, or irregular heartbeat. Nicotine can increase your heart rate.
- high blood pressure not controlled with medication. Nicotine can increase blood pressure.
- stomach ulcer or diabetes
- history of seizures

Ask a doctor or pharmacist before use if you are

- using a non-nicotine stop smoking drug
- taking prescription medicine for depression or asthma. Your prescription dose may need to be adjusted.

Stop use and ask a doctor if

- mouth, teeth, or jaw problems occur
- irregular heartbeat or palpitations occur
- you get symptoms of nicotine overdose such as nausea, vomiting, dizziness,

diarrhea, weakness, and rapid heartbeat

• you have symptoms of an allergic reaction (such as difficulty breathing or rash)

Keep out of reach of children and pets.

Pieces of nicotine gum may have enough nicotine to make children and pets sick. Wrap used pieces of gum in paper and throw away in the trash. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

- if you are under 18 years of age, ask a doctor before use
- before using this product, read the enclosed User's Guide for complete directions and other important information
- begin using the gum on your quit day
- if you smoke your first cigarette more than 30 minuted after waking up, use 2 mg nicotine gum
- if you smoke your first cigarette within 30 minutes of waking up, use 4 mg nicotine gum according to the following 12 week schedule

Weeks 1 to 6	Weeks 7 to 9	Weeks 10 to 12
1 piece every 1 to 2 hours	1 piece every 2 to 4 hours	1 piece every 4 to 8 hours

- nicotine gum is a medicine and must be used a certain way to get the best results
- chew the gum slowly until it tingles. Then park it between your cheek and gum. When the tingle is gone, begin chewing again, until the tingle returns.
- repeat this process until most of the tingle is gone (about 30 minutes)
- do not eat or drink for 15 minutes before chewing the nicotine gum, or while chewing a piece
- to improve your chances of quitting, use at least 9 pieces per day for the first 6 weeks
- if you experience strong or frequent cravings, you may use a second piece within the hour. However, do not continuously use one piece after another since this may cause hiccups, heartburn, nausea or other side effects.
- do not use more than 24 pieces a day
- it is important to complete treatment. If you feel you need to use the gum for a longer period to keep from smoking, talk to your health care provider.

other information

- each piece contains: calcium 115 mg, sodium 8 mg
- store at 20-25°C (68-77°F)
- protect from light

Inactive ingredients

acacia, acesulfame potassium, carnauba wax, D&C yellow #10, FD&C blue #2, FD&C red #40, FD&C yellow #6, flavors, gum base, hydroxypropyl cellulose, magnesium oxide, sodium bicarbonate, sodium carbonate, sucralose, talc, titanium dioxide, xylitol.

Questions or comments?

Call

Principal Display Panel

*Compare to the active ingredient in Nicorette® Gum

Nicotine Gum

Nicotine Polacrilex Gum USP, 4 mg (nicotine)

Stop Smoking Aid

FOR THOSE WHO SMOKE THEIR FIRST CIGARETTE WITHIN 30 MINUTES OF WAKING UP

If you smoke your first cigarette <u>MORE THAN</u> 30 MINUTES after waking up, use Nicotine Polacrilex gum, 2 mg

Sugar Free

Coated Fruit

TO INCREASE YOUR SUCCESS IN QUITTING:

1. You must be motivate to quit

2. **Use Enough** - Chew **at least 9 pieces** of Nicotine Polacrilex Gum per day during the first six weeks

- 3. Use long enough Use Nicotine Polacrilex Gum for the full 12 weeks
- 4. Use with a Support Program as directed in the enclosed User's Guide

To remove the gum, tear off single unit

Peel off backing starting at corner with loose edge

Push gum through foil

- not for sale to those under 18 years of age
- proof of age required
- not for sale in vending machines or from any source where proof of age cannot be verified

BLISTER PACKAGED FOR YOUR PROTECTION. DO NOT USE IF INDIVIDUAL SEALS ARE OPEN OR TORN.

*This product is not manufactured or distributed by Haleon group of companies, distributor of Nicorette $\ensuremath{\mathbb{B}}$ Gum

Marketed By: Lucy Goods Inc.

2510 E Sunset Road

Suite 5 #A656 Las Vegas, NV 89120

Package Label



TIURA

aeug-Jeffing

	stop use and ask a dector if Stop use and ask a dector if	und Date ingeneration and the second se) emohumeos, aceacias 23 DAN ben 3407, joskel 2% ei	240, yoow actumus work of the 240 your actument of the second of the 240 years of the second of the second of the 240 years of the second o
gm S, R2U muD ແສ່ຈະດັບດີ ອາຊ່າວດັກ ອອກ ແລະ ອຸດຄົນ ແລະ 201 ອູກ S, R2U muD ແສ່ຈະດັບດີ ອາຊ່າວດັກ ອອກ ແລະ ລູດ ອົກການສາມ	ana ung Yi ana ung Airanada tangka tangka tangka sa vatasa gaba pakerangka sa varangka sa sa sa sa sa sa sa sa sa sa s	Other information = stars at 20 -25°C (68 -71 = stars at 20 -25°C (68 -71) = stars at 20 -25°C (78 -71) = stars at 20	multiples up a cit multiples	Dur g
(anitooin) gm 4, G2U mug xafinoslog anitooil) bid gnöking data poa two skoke them Plast crohert too coa two skoket of the skoket	Add a costor before used from lower the source state of the point with the source states, or implice heartheat Monthes can increase your heart disease, recert heart allocity of Mit medication. Montheat mitigh the group measure and carinoles with medication. Montheat mitight the group measure and analogies with medication. Montheat point and the source of the point of the source of the source of the source of the source of the point of the source of the source of the source of the source of the point of the source of the point of the source of the point of the source of the sour	weeks within the hour experience strang since this may cause you by not use more than 24 25	j or fraquent crawinge, yo sy do not continuously us ta incoups, haarfburn, na ta braafmont, if you haa die braafmont, if you haal	Aon used printe gue Smuller nove based state alge allectr nove based state succher numit new a second bace
Micotine Gum	Warmhings before an exist based for the easily meeting and the advice of your needs without integration of the meeting meeting for the proved to be set to be another proved, can associety herm your child for the behaved to be set to be an another proved and another behaved to be an and fully howen.	ctemps a bace a company of the company of the company a company of the company of the company of the company a company of the company of the company of the company a company of the c	i Xiaq neufit. Singles. Then park it good, begin chreater in the standard of the steps innest of the steps of an and a setting of the steps of the steps of the steps of the steps of the steps of the steps of the steps of the step of t	behween your cheek and ain, until the tingle returns.
	080 the behaviors, general entroin general, including learning associated with ording	1 to 5 tions 1 biece eveny	S ID 4 HOURS	ynave aneig 1 Eruori 8 oc 4 Aeruori 8 oc 4
TUCK	asoqruf (soolq gniword) riose nij trialbargni evitok is prisens eriz	gm 2 eau qui gaixiew to eaturian OG nittlew otkreata territ nevy aixone unit ti m silabelica sterw 31 gaiwoliel erti al pathrocce mug entitodin 21 al 01 zaleeW 2 al 7 zaleeW 8 ad 1 zaleeW		
	spej brug	Drug Facts (continu	(per	

Product Information					
Product Type	HUMAN OTC DRUG	ltem Code	(Source)	NDC:7	2224-931
Route of Administration	ORAL				
Active Ingredient/Activ	e Moiety				
Ingre	edient Name		Basis of Str	ength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6	۲)	NICOTINE		4 mg
Inactive Ingredients					
	Ingredient Name				Strength
ACACIA (UNII: 5C5403N260)					
ACESULFAME POTASSIUM (UN	. ,				
CARNAUBA WAX (UNII: R12CBM					
FD&C RED NO. 40 (UNII: WZ B9: HYDROXYPROPYL CELLULOSE					
D&C YELLOW NO. 10 (UNII: 35					
SODIUM BICARBONATE (UNII: 8					
SODIUM CARBONATE (UNII: 45					
TALC (UNII: 7SEV7J4R1U)	,				
TITANIUM DIOXIDE (UNII: 15FIX	9V2JP)				
XYLITOL (UNII: VCQ006KQ1E)	· ·				
SUCRALOSE (UNII: 96K6UQ3ZD4	4)				
	10G171G)				
MAGNESIUM OXIDE (UNII: 3A3U					
MAGNESIUM OXIDE (UNII: 3A3U FD&C BLUE NO. 2 (UNII: L06K8					

Pr	roduct Chara	cteristics			
Со	lor	brown (Light Tan)	Score	no score	
Sh	аре	SQUARE	Size	14mm	
Fla	avor	FRUIT (Coated Fruit)	mprint Code		
Со	ontains				
Pa	ackaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
	NDC:72224-931- 10	100 in 1 CARTON	04/25/2025		
1		1 in 1 BLISTER PACK; Type 0: Not a Combinatio Product	on		
	arkating I	nformation			
Μ	arketing I				
Μ	Marketing Category	Application Number or Monograp Citation	h Marketing Start Date	Marketing End Date	

Labeler - Lucy Goods, Inc. (080875415)

Revised: 4/2025

Lucy Goods, Inc.