

**ALKA-SELTZER PLUS MAXIMUM STRENGTH COLD AND COUGH LIQUID GELS-  
acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide,  
phenylephrine hydrochloride capsule, liquid filled  
Bayer HealthCare LLC.**

-----

**Alka-Seltzer Plus® Maximum Strength Cold & Cough Liquid Gels**

***Drug Facts***

***Active ingredients (in each capsule)***

Acetaminophen 325 mg

Chlorpheniramine maleate 2 mg

Dextromethorphan hydrobromide 10 mg

Phenylephrine hydrochloride 5 mg

***Purposes***

Pain reliever/fever reducer

Antihistamine

Cough suppressant

Nasal decongestant

***Uses***

- temporarily relieves these symptoms due to a cold or flu:
- minor aches and pains · headache
- nasal congestion · sinus congestion and pressure
- cough · sore throat
- runny nose · sneezing
- temporarily reduces fever

***Warnings***

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening · blisters · rash · hives

· facial swelling · asthma (wheezing) · shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use to sedate children.**

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

**Ask a doctor before use if you have**

- liver disease ● heart disease ● high blood pressure
- thyroid disease ● diabetes ● glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- **do not exceed recommended dosage**
  - may cause marked drowsiness
- avoid alcoholic drinks

- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

**Stop use and ask a doctor if**

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

- nervousness, dizziness, or sleeplessness occurs

**If pregnant or breast-feeding**, ask a health professional before use.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

***Directions***

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

**Other information**

**Other information**

- store at room temperature. Avoid excessive heat above 40°C (104° F).

***Inactive ingredients*** D&C red #33, FD&C blue #1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol sorbitan solution, titanium dioxide

**Questions or Comments**

**Questions or comments?** 1-800-986-0369 (Mon-Fri 9AM - 5PM EST)

**PDP**



## ALKA-SELTZER PLUS MAXIMUM STRENGTH COLD AND COUGH LIQUID GELS

acetaminophen, chlorpheniramine maleate, dextromethorphan hydrobromide, phenylephrine hydrochloride capsule, liquid filled

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:0280-1570
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>CHLORPHENIRAMINE MALEATE</b> (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	2 mg
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

## Inactive Ingredients

Ingredient Name	Strength
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>D&amp;C RED NO. 33</b> (UNII: 9DBA0SBB0L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL 400</b> (UNII: B6978945GQ)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SHELLAC</b> (UNII: 46N107B71O)	
<b>SODIUM HYDROXIDE</b> (UNII: 55X04QC32I)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>SORBITOL</b> (UNII: 506T60A25R)	

## Product Characteristics

<b>Color</b>	purple	<b>Score</b>	no score
<b>Shape</b>	OVAL (Elliptical)	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	ASP;CC
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-1570-20	2 in 1 CARTON	03/23/2021	
1		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
2	NDC:0280-1570-10	1 in 1 CARTON	03/23/2021	
2		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		
3	NDC:0280-1570-40	4 in 1 CARTON	03/23/2021	
3		10 in 1 BLISTER PACK; Type 0: Not a Combination Product		

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	07/01/2019	

**Labeler** - Bayer HealthCare LLC. (112117283)