

**GENTLE LAXATIVE- bisacodyl tablet, delayed release**  
**Bryant Ranch Prepack**

-----  
**Sunmark 44-327**

***Active ingredient (in each tablet)***

Bisacodyl USP, 5 mg

***Purpose***

Stimulant laxative

***Uses***

- for relief of occasional constipation and irregularity
- this product generally produces bowel movement in 6 to 12 hours

***Warnings***

**Do not use**

if you cannot swallow without chewing.

**Ask a doctor before use if you have**

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

**When using this product**

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- you may have stomach discomfort, faintness and cramps

**Stop use and ask a doctor if**

- you have rectal bleeding or fail to have a bowel movement after use of a laxative. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

***Directions***

- take with a glass of water

adults and children 12 years and over	take 1 to 3 tablets in a single daily dose
children 6 to under 12 years	take 1 tablet in a single daily dose
children under 6 years	ask a doctor

***Other information***

- **TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN**
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- avoid excessive humidity
- see end flap for expiration date and lot number

***Inactive ingredients***

acacia, ammonium hydroxide, calcium carbonate, carnauba wax, colloidal anhydrous silica, corn starch, D&C yellow #10 aluminum lake, FD&C yellow #6 aluminum lake, hypromellose, iron oxide black, lactose anhydrous, magnesium stearate, methylparaben, polydextrose, polyethylene glycol, polyvinyl acetate phthalate, povidone, propylene glycol, propylparaben, shellac glaze, simethicone, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate

***Questions or comments?***

**1-800-426-9391**

**HOW SUPPLIED**

- NDC: 71335-1754-0: 25 Tablets in a BLISTER PACK
- NDC: 71335-1754-1: 30 Tablets in a BLISTER PACK
- NDC: 71335-1754-2: 2 Tablets in a BLISTER PACK
- NDC: 71335-1754-3: 3 Tablets in a BLISTER PACK
- NDC: 71335-1754-4: 4 Tablets in a BLISTER PACK
- NDC: 71335-1754-5: 10 Tablets in a BLISTER PACK
- NDC: 71335-1754-6: 90 Tablets in a BLISTER PACK
- NDC: 71335-1754-7: 8 Tablets in a BLISTER PACK
- NDC: 71335-1754-8: 100 Tablets in a BLISTER PACK
- NDC: 71335-1754-9: 20 Tablets in a BLISTER PACK

**Bisacodyl 5mg Tablet**



GTIN 00371335175412  
 Lot 20820  
 Exp 4/30/2026  
 SN 0123456789

Drug Facts	
<b>Active ingredient (in each tablet)</b>	<b>Purpose</b>
Bisacodyl 5 mg	Stimulant Laxative
<b>Uses</b>	
For relief of occasional constipation and irregularity - this product generally produces bowel movement in 6 to 12 hours	
<b>Warnings</b>	
Do not use if you cannot swallow without chewing	
Ask a doctor before use if you have	
*stomach pain, nausea, vomiting	
*noticed a sudden change in bowel habits that lasts more than 2 weeks	
When using this product	
*do not chew or crush tablet(s)	
*do not use within 1 hour after taking an antacid or milk	
*it may cause stomach discomfort, faintness and cramps	
*Stop use and ask a doctor if - you have rectal bleeding or no bowel movement after using this product	
These could be signs of a serious condition.	
*you need to use a laxative for more than 1 week	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
<b>Other Information</b>	
*Tamper Evident: Do not use if imprinted seal under cap is missing or broken	
*Store at room temperature 15°-30°C (59°-86°F)	
*Avoid excessive humidity - packaging not child resistant	
<b>Directions</b>	
*Take with a glass of water	
*Adults and children 12 years and older: take 1 to 3 tablets in a single daily dose	
*Children 12 and under: ask a doctor	
<b>Inactive Ingredients</b>	
Acacia, ammonium hydroxide, calcium carbonate, corn starch, D&C yellow #10 lake, FD&C yellow #6 lake, hypromellose, iron oxide black, lactose, magnesium stearate, methylparaben, PEG, polydextrose, polyvinyl acetate phthalate, propylparaben, propylene glycol, povidone, shellac, simethicone, silica, sodium alginate, sodium benzoate, sodium bicarbonate, stearic acid, sucrose, talc, titanium dioxide, triacetin, triethyl citrate, wax	

NDC 71335-1754-1

**Bisacodyl Tablets, USP**

5 mg

30 Tablets



Repackaged by: **Bryant Ranch Prepack, Inc.** Manufactured by: **LNK International, Inc.**  
 Burbank, CA 91504 USA



**GENTLE LAXATIVE**

bisacodyl tablet, delayed release

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:71335-1754(NDC:49348-032)
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>BISACODYL</b> (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>ACACIA</b> (UNII: 5C5403N26O)	
<b>AMMONIA</b> (UNII: 5138Q19F1X)	
<b>CALCIUM CARBONATE</b> (UNII: H0G9379FGK)	
<b>CARNAUBA WAX</b> (UNII: R12CBM0EIZ)	
<b>SILICON DIOXIDE</b> (UNII: ETJ7Z6XBU4)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>D&amp;C YELLOW NO. 10 ALUMINUM LAKE</b> (UNII: CQ3XH3DET6)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>FERROSFERRIC OXIDE</b> (UNII: XM0M87F357)	
<b>ANHYDROUS LACTOSE</b> (UNII: 3SY5LH9PMK)	
<b>MAGNESIUM STEARATE</b> (UNII: 70097M6I30)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>POLYDEXTROSE</b> (UNII: VH2XOU12IE)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)	
<b>POLYVINYL ACETATE PHTHALATE</b> (UNII: 58QVG85GW3)	
<b>POVIDONE, UNSPECIFIED</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	

<b>SHELLAC</b> (UNII: 46N107B71O)
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)
<b>WATER</b> (UNII: 059QF0KO0R)
<b>SODIUM ALGINATE</b> (UNII: C269C4G2ZQ)
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)
<b>SODIUM BICARBONATE</b> (UNII: 8MDF5V39QO)
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)
<b>SUCROSE</b> (UNII: C151H8M554)
<b>TALC</b> (UNII: 7SEV7J4R1U)
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)
<b>TRIACETIN</b> (UNII: XHX3C3X673)
<b>TRIETHYL CITRATE</b> (UNII: 8Z96QXD6UM)

### Product Characteristics

<b>Color</b>	orange	<b>Score</b>	no score
<b>Shape</b>	ROUND	<b>Size</b>	6mm
<b>Flavor</b>		<b>Imprint Code</b>	5
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71335-1754-0	25 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
2	NDC:71335-1754-1	30 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
3	NDC:71335-1754-2	2 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
4	NDC:71335-1754-3	3 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
5	NDC:71335-1754-4	4 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
6	NDC:71335-1754-5	10 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
7	NDC:71335-1754-6	90 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
8	NDC:71335-1754-7	8 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
9	NDC:71335-1754-8	100 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	
10	NDC:71335-1754-9	20 in 1 BOTTLE; Type 0: Not a Combination Product	02/24/2022	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	505G(a)(3)	03/25/2002	05/04/2025

**Labeler** - Bryant Ranch Prepack (171714327)

**Registrant** - Bryant Ranch Prepack (171714327)

**Establishment**

Name	Address	ID/FEI	Business Operations
Bryant Ranch Prepack		171714327	REPACK(71335-1754) , RELABEL(71335-1754)

Revised: 4/2024

Bryant Ranch Prepack