

WALGREENS GLYCERIN- glycerin suspension enema **Walgreens**

WALG GLYCERIN APPLICATR 7.5ML 6S

Hyperosmotic Laxative

for relief of occasional constipation.

This product usually produces a bowel movement in 15 minutes to 1 hour.

May cause rectal discomfort or a burning sensation.

For rectal use only

Do not use when abdominal pain, nausea, or vomiting are present unless directed by a doctor.

Ask a doctor before using any laxative if you

- have a sudden change in bowel habits lasting more than 2 weeks
- already used a laxative for more than 1 week

Stop using this product and consult a doctor if you have

- rectal bleeding
- no bowel movement within 1 hour of using this product

These symptoms may indicate a serious condition.

If swallowed, get medical help or contact a Poison Control Center right away at 1-800-222-1222

Single Daily Dosage (per 24 hours)

- **Adults and children 6 years and over:** 1 suppository or as directed by a doctor
- **Children under 6 years:** consult a doctor

Left-side position: Lie on left side with right knee bent and arms resting comfortably.

Knee-chest position: Kneel, then lower head and chest forward until left side of face is resting on surface with left arm folded comfortably.

Administering the laxative:

CAUTION: REMOVE ORANGE PROTECTIVE SHIELD BEFORE INSERTING.

Hold unit upright, grasping the bottle with fingers. Grasp protective shield with other hand, pull gently to remove. With steady pressure, insert tip into rectum with a slight side-to-side movement, with tip pointing toward navel. **Do not force the applicator tip into the rectum, as this can cause injury.** Insertion may be easier if person receiving suppository bears down as if having a bowel movement. This helps relax the muscles around the anus. Squeeze completely the accordion bottle to expel liquid. While continuing to squeeze the accordion bottle, remove tip from rectum and discard unit. The unit contains more than the amount of liquid needed for effective use. A small amount of liquid will remain in the unit after use.

edetate disodium, purified water

Store at room temperature.

Keep bottles in carton until use.

Questions? 1-800-WALGREENS (1-800-925-4733)



WALGREENS GLYCERIN

glycerin suspension enema

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:0363-0499 |
| Route of Administration | RECTAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| GLYCERIN (UNII: PDC6A3C00X) (GLYCERIN - UNII:PDC6A3C00X) | GLYCERIN | 5.4 g |

Inactive Ingredients

| Ingredient Name | Strength |
|-------------------------------------|----------|
| EDETATE DISODIUM (UNII: 7FLD91C86K) | |

Product Characteristics

| | | | |
|----------|-------------------|--------------|--|
| Color | white (colorless) | Score | |
| Shape | | Size | |
| Flavor | | Imprint Code | |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:0363-0499-06 | 6 in 1 CARTON | 02/28/2025 | |
| 1 | | 1 in 1 BOTTLE, UNIT-DOSE; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M007 | 02/28/2025 | |

Labeler - Walgreens (008965063)

Registrant - MSH Pharma Inc. (243301992)

Establishment

| Name | Address | ID/FEI | Business Operations |
|-----------------|---------|-----------|------------------------|
| MSH Pharma Inc. | | 243301992 | manufacture(0363-0499) |

Revised: 2/2025

Walgreens