

MEDIQUE ASPIRN- medique aspirin tablet, film coated
HF Acquisition Co LLC, DBA HealthFirst

Aspirin (NSAID*) 325mg *nonsteroidal anti-inflammatory drug

Pain Reliever/Fever Reducer

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or Flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: hives, skin reddening, facial swelling, rash, asthma (wheezing), blisters or shock. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: this contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems, take a blood thinner (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others), have 3 or more alcoholic drinks everyday while using this product, take more for a longer time than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer; right before or after heart surgery, if you are taking a prescription drug for gout, diabetes or arthritis.

Ask a doctor before use if: stomach bleeding warning applies to you, you have a history of stomach problems such as heartburn, you have high blood pressure, heart disease, liver cirrhosis or kidney disease, you are taking a diuretic.

Ask a Doctor or Pharmacist before use if you are: under a doctor's care for any serious condition, taking any other drug.

When using this product: take with food or milk if stomach upset occurs

Stop use and Ask a Doctor if: you experience any of the following signs of stomach bleeding, you feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, you have difficulty swallowing, if ringing in the ears or loss of hearing occurs, redness or swelling is present in painful areas, or any new symptoms appear.

If pregnant or Breastfeeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of Children. In case of overdose, get medical help or contact a poison control center right away.

DIRECTIONS

Do not use more than directed - the smallest effective dosage should be used. Drink a

full glass of water with each dose. Do not take longer than 10 days, unless directed by a doctor.

Adults and Children (12 years and older): take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours or as directed as a doctor.

Children under 12 years: Do not give to children under 12 years of age

Avoid excessive heat and humidity, do not use any open or torn packets.

USE

Temporarily relieves minor aches and pains associated with: headache, muscular aches, minor arthritis pain, backache, common cold, toothache, menstrual cramps, temporarily reduces fever.

Ask a doctor before use if: stomach bleeding warning applies to you, you have a history of stomach problems such as heartburn, you have high blood pressure, heart disease, liver cirrhosis or kidney disease, you are taking a diuretic.

Ask a Doctor or Pharmacist before use if you are: under a doctor's care for any serious condition, taking any other drug.

Stop use and Ask a Doctor if: you experience any of the following signs of stomach bleeding, you feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, you have difficulty swallowing, if ringing in the ears or loss of hearing occurs, redness or swelling is present in painful areas, or any new symptoms appear.

If pregnant or Breastfeeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless directed to do so by a doctor because it may cause problems in the unborn child or complications during deliver.

Keep out of reach of Children. In case of overdose, get medical help or contact a poison control center right away.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer; right before or after heart surgery, if you are taking a prescription drug for gout, diabetes or arthritis.

1 (2 Pack) 325 MG Packet

Adults and Children (12 years and older): take 1 or 2 tablets with water every 4 hours as needed. Do not take more than 12 tablets in 24 hours or as directed as a doctor.

Pain Reliever/Fever Reducer

Reye's Syndrome: Children and teenagers who have or are recovering from chicken pox or Flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy Alert: Aspirin may cause a severe allergic reaction which may include: hives, skin reddening, facial swelling, rash, asthma (wheezing), blisters or shock. If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: this contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you: are age 60 or older; have had stomach ulcers or bleeding problems, take a blood thinner (anticoagulant) or steroid drug; take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen or others), have 3 or more alcoholic drinks everyday while using this product, take more for a longer time than directed.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer; right before or after heart surgery, if you are taking a prescription drug for gout, diabetes or arthritis.

Ask a doctor before use if: stomach bleeding warning applies to you, you have a history of stomach problems such as heartburn, you have high blood pressure, heart disease, liver cirrhosis or kidney disease, you are taking a diuretic.

Ask a Doctor or Pharmacist before use if you are: under a doctor's care for any serious condition, taking any other drug.

When using this product: take with food or milk if stomach upset occurs

Stop use and Ask a Doctor if: you experience any of the following signs of stomach bleeding, you feel faint, vomit blood, have bloody or black stools, have stomach pain that does not get better, gets worse or lasts more than 10 days, fever gets worse or lasts more than 3 days, you have difficulty swallowing, if ringing in the ears or loss of hearing occurs, redness or swelling is present in painful areas, or any new symptoms appear.

If pregnant or Breastfeeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of Children. In case of overdose, get medical help or contact a poison control center right away.

Do not use more than directed - the smallest effective dosage should be used. Drink a full glass of water with each dose. Do not take longer than 10 days, unless directed by a doctor.

CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)

CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)

TITANIUM DIOXIDE (UNII: 15FIX9V2JP)

MINERAL OIL (UNII: T5L8T28FGP)

HYPROMELLOSES (UNII: 3NXW29V3WO)

STARCH, CORN (UNII: O8232NY3SJ)

(01) 00351662127728
(17) 200630
(21) 77328389192
(10) 5462



See manufacturer's package insert
Distributed by HF Acquisition Co., LLC
Mukilteo, WA 98275

2 Tablets **Médique**
Aspirin

Active ingredient (in each tablet) **Purpose**
Aspirin (ASA) 325 mg Pain reliever/fever reducer
Nonsteroidal anti-inflammatory drug

Use: Temporarily relieves minor aches and pains due to
● headache ● muscular aches ● minor arthritis pain ● backache
● common cold ● toothache ● menstrual cramps
Temporarily reduces fever.

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include ● hives ● skin redness ● facial swelling ● chest ● wheezing ● asthma ● shock.

If an allergic reaction occurs, stop use and seek medical help right away.
Stomach bleeding warning: This product contains an NSAID, which may cause serious stomach bleeding. The chance is higher if you

● are age 65 or older ● have had stomach ulcers or bleeding problems ● take a blood thinner (anticoagulant) or blood-thinning drug ● take other drugs containing aspirin or nonsteroidal NSAIDs (aspirin, ibuprofen, naproxen, or others) ● have 3 or more alcoholic drinks every day while using this product ● use this drug for a longer time. Use caution.

Store at 20°-25° (68°-77°F) - avoid excessive heat and humidity.
-Keep out of reach of children - Do not use any expired or open packets
MSL for MEDIQUE PRODUCTS - Fort Myers, FL 33907

MEDIQUE ASPIRIN

medique aspirin tablet, film coated

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:51662-1277(NDC:47682-097)

Route of Administration ORAL

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|----------|
| ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E) | ASPIRIN | 325 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| CROSCARMELLOSE SODIUM (UNII: M28OL1HH48) | |
| HYPROMELLOSES (UNII: 3NXW29V3WO) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U) | |
| TITANIUM DIOXIDE (UNII: 15FIX9V2JP) | |
| MINERAL OIL (UNII: T5L8T28FGP) | |

Product Characteristics

| | | | |
|----------|-------|--------------|----------|
| Color | white | Score | no score |
| Shape | ROUND | Size | 10mm |
| Flavor | | Imprint Code | TCL;011 |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:51662-1277-2 | 2 in 1 PACKET; Type 0: Not a Combination Product | 12/30/2008 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | 505G(a)(3) | 12/30/2008 | |

Labeler - HF Acquisition Co LLC, DBA HealthFirst (045657305)

Registrant - HF Acquisition Co LLC, DBA HealthFirst (045657305)

Establishment

| Name | Address | ID/FEI | Business Operations |
|--|---------|-----------|---------------------|
| HF Acquisition Co LLC, DBA HealthFirst | | 045657305 | relabel(51662-1277) |