

BLOOD CLOTTING- benzethonium chloride and lidocaine aerosol, spray
HART Health

Blood Clotting Spray

Active Ingredients

Benzethonium Chloride 0.2% w/w
Lidocaine 4% w/w

Purpose

Topical Antiseptic
Topical Anesthetic

Uses

- for help in control of superficial bleeding
- for the temporary relief of pain
- to help protect against skin infection in minor cuts, scrapes, and burns

Warnings

For external use only

Flammable

- keep away from flame
- contents under pressure
- do not puncture or incinerate container
- do not expose to temperatures above 120°F

Do not use

- in or near the eyes or other mucous membranes
- over large portions of the body
- on deep or puncture wounds, raw surfaces, blistered areas, animal bites, or serious burns

Stop use and ask doctor if

- redness, swelling, or irritation occurs
- infection occurs
- condition persists or worsens
- symptoms persist for more than 7 days

Keep out of reach of children.

If swallowed, get medical help or contact a Poison Control Center right away. 1-800-222-1222

▯Directions

Adults and children 23 years of age and over:

- shake can well before using
- hold 6-8 inches from moist injured area
- spray until area is covered and superficial bleeding is stopped
- repeat as necessary
- scab-like cover will gradually disappear as healing takes place or may be removed with soap and water

Children under 12 years of age: ask a doctor

Other Information

- avoid inhaling
- use only as directed
- intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal

Inactive Ingredients

isobutane, isopropyl myristate, karaya, n-butane, propane, silica, sorbitan monooleate, talc, tragacanth

Principal Display

6.75" x 4"
PMS 2602C

Purpose
Benzethonium chloride 0.2% w/w..... Topical antiseptic
Lidocaine 4% w/w..... Topical anesthetic

Uses ■ for help control of superficial bleeding ■ for the temporary relief of pain
■ to help protect against skin infection in minor cuts, scrapes, and burns

Warnings For external use only

Flammable ■ keep away from fire or flame ■ contents under pressure
■ do not puncture or incinerate container ■ do not expose to temperatures above 120°F

Do not use ■ in or near the eyes or other mucous membranes
■ over large portions of the body
■ on deep or puncture wounds, raw surfaces, blistered areas, animal bites, or serious burns

Stop use and ask a doctor if ■ redness, swelling or irritation occurs ■ infection occurs
■ condition persists or worsens ■ symptoms persist for more than 7 days

Keep out of reach of children.
If swallowed, get medical help or contact a Poison Control Center right away. 1-800-222-1222

Directions Adults and children 12 years of age and over:
■ shake can well before using ■ hold can 6 to 8 inches away from moist injured area
■ spray until area is covered and superficial bleeding is stopped ■ repeat as necessary
■ scab-like cover will gradually disappear as healing takes place or may be removed with soap and water
Children under 12 years of age: ask a doctor

Other information ■ avoid inhaling ■ use only as directed
■ intentional misuse by deliberately concentrating and inhaling the contents may be harmful or fatal

Inactive ingredients
isobutane, isopropyl myristate, keratin, n-butane, propane, silica, sorbitan monooleate, talc, triglycerin

2760-21
N 50332-02171 6



NDC 50332-0217-1 Item 2760

Blood Clotting Spray
Coagulant

Helps control superficial bleeding
Helps prevent infection

Contains no fluorocarbons environmentally safe
3 oz. (85 gm)

HART
HART Health
South, WI 53158 www.harthealth.com

BLOOD CLOTTING

benzethonium chloride and lidocaine aerosol, spray

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50332-0217
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	3400 mg in 85 g
BENZETHONIUM CHLORIDE (UNII: PH41D05744) (BENZETHONIUM - UNII:1VU15B70BP)	BENZETHONIUM CHLORIDE	170 mg in 85 g

Inactive Ingredients

Ingredient Name	Strength
ISOBUTANE (UNII: BXR49TP611)	

ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS)	
KARAYA GUM (UNII: 73W9IQY50Q)	
BUTANE (UNII: 6LV4FOR43R)	
PROPANE (UNII: T75W9911L6)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
3-(TRIETHOXYSILYL)PROPYLAMINE (UNII: L8S6UBW552)	
SORBITAN MONOOLEATE (UNII: 06XEA2VD56)	
TALC (UNII: 7SEV7J4R1U)	
TRAGACANTH (UNII: 294435702O)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:50332-0217-1	85 g in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product	04/01/1974	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M	04/01/1974	

Labeler - HART Health (069560969)

Revised: 1/2026

HART Health