

IBUPROFEN 100- ibuprofen tablet, coated
IBUPROFEN 250- ibuprofen tablet, coated
JHK Inc dba American Safety & First Aid

IBUPROFEN

Drug Facts

Active ingredient (In Each Tablets)

Ibuprofen, USP 200 mg (NSAID)

*Nonsteroidal Anti-Inflammatory Drug

Purpose

pain reliever/fever reducer

Use

temporarily relieves fever and minor aches and pains associated with:

- common cold
- backache
- headache
- toothache
- arthritis
- muscular aches
- menstrual cramps

Warnings

Allergy Alert

Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin.

Symptoms may include:

- hives
- asthma (wheezing)
- facial swelling
- shock
- skin reddening
- rash
- blisters
- If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning

This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic
- drinks every day while using this product
- take more or for a longer time than directed

Do not use

- if you ever had an allergic reaction to any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

Ask a doctor or pharmacist before use if you are

- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) drug
- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding:

- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. IT IS ESPECIALLY IMPORTANT NOT TO USE IBUPROFEN DURING THE LAST 3 MONTHS OF PREGNANCY UNLESS DEFINITELY DIRECTED TO DO SO BY A DOCTOR BECAUSE IT MAY CAUSE PROBLEMS IN THE UNBORN CHILD OR COMPLICATIONS DURING DELIVERY.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed
- the smallest effective dose should be used
- do not take longer than 10 days, unless directed by a doctor (see Warnings)

Adults: 1 tablet every 4 to 6 hours while symptoms occur; if pain or fever does not respond to 1 tablet, 2 tablets may be used, but do not exceed 6 tablets in 24 hours unless directed by a doctor.

Children under 12 years: Ask a doctor.

Other Information

- do not use if packet is opened or torn
- Store at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F)

Inactive Ingredients

crosscarmellose sodium¹, FD& C Blue #2¹, FD& C red #40¹, FD& C Yellow #6¹, hypromellose¹, iron oxide red¹, lactose monohydrate¹, magnesium stearate¹, maltodextrin¹, medium chain triglycerides¹, microcrystalline cellulose¹, polydextrose¹, polyethylene glycol¹, polyvinyl alcohol¹, povidone (K-30)¹, silicon dioxide, sodium starch glycolate¹, starch, stearic acid¹, talc¹, titanium dioxide, triacetin¹,

1 contains one or more of these ingredients

Questions?

1-866-651-3660 Monday-Friday

8:00am-5:00pm EST

PRINCIPAL DISPLAY PANEL - 100 Tablet Packet Box

This product is not manufactured or distributed by
Pfizer Consumer Healthcare
Owner of the registered trademark Advil®.

Product #1102

Ibuprofen

Pain Reliever/Fever Reducer

Active ingredient:

Ibuprofen

- Temporary relief from pain, fever and inflammation
- Soothes muscles aches and minor arthritis pain
- Non-drowsy, aspirin-free formula

These quality comfort tablets are formulated and packaged for use in the

workplace and meet ANSI Z308.1-2015 standards. They are sealed in tamper-evident packets for safety and convenience.

100 Tablets • 2 Tablets Per Packet

This product is not manufactured or distributed by
Pfizer Consumer Healthcare
Owner of the registered trademark Advil®.

Product #1102



Active ingredient:
Ibuprofen

Pain Reliever/Fever Reducer

- Temporary relief from pain, fever and inflammation
- Soothes muscles aches and minor arthritis pain
- Non-drowsy, aspirin-free formula

These quality comfort tablets are formulated and packaged for use in the workplace and meet ANSI Z308.1-2015 standards. They are sealed in tamper-evident packets for safety and convenience.

100 Tablets • 2 Tablets Per Packet

Drug Facts (continued)

If pregnant or breast-feeding, ask a health professional before use. IT IS ESPECIALLY IMPORTANT NOT TO USE IBUPROFEN DURING THE LAST 3 MONTHS OF PREGNANCY UNLESS DEFINITELY DIRECTED TO DO SO BY A DOCTOR BECAUSE IT MAY CAUSE PROBLEMS IN THE UNBORN CHILD OR COMPLICATIONS DURING DELIVERY.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions • do not take more than directed • the smallest effective dose should be used • do not take longer than 10 days, unless directed by a doctor (see Warnings)
Adults: 1 tablet every 4 to 6 hours while symptoms occur; if pain or fever does not respond to 1 tablet, 2 tablets may be used, but do not exceed 6 tablets in 24 hours unless directed by a doctor. *Children under 12 years: Ask a doctor.*

Other Information

• do not use if packet is opened or torn • Store at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F)

Inactive Ingredients croscarmellose sodium*, FD&C Blue #2*, FD&C red #40*, FD&C Yellow #6*, hypromellose*, iron oxide red*, lactose monohydrate*, magnesium stearate*, maltodextrin*, medium chain triglycerides*, microcrystalline cellulose*, polydextrose*, polyethylene glycol*, polyvinyl alcohol*, povidone (K-30)*, silicon dioxide, sodium starch glycolate*, starch, stearic acid*, talc*, titanium dioxide, triacetin*, *contains one or more of these ingredients

Questions? 1-866-651-3660 Monday-Friday
8:00am-5:00pm EST

Manufactured for
American Safety & First Aid
Osceola, IN 46561


Pain Reliever/Fever Reducer

Ibuprofen



Product #1102

100 Tablets • 2 Tablets Per Packet



PRINCIPAL DISPLAY PANEL - 250 Tablet Packet Box

Product #1104

Ibuprofen

Pain Reliever/Fever Reducer

Active ingredient:

Ibuprofen

- Temporary relief from pain, fever and inflammation
- Soothes muscles aches and minor arthritis pain
- Non-drowsy, aspirin-free formula

These quality comfort tablets are formulated and packaged for use in the workplace and meet ANSI Z308.1-2015 standards. They are sealed in tamper-evident packets for safety and convenience.

250 Tablets • 2 Tablets Per Packet

This product is not manufactured or distributed by
Pfizer Consumer Healthcare
Owner of the registered trademark Advil®.

Product #1104



Active ingredient:
Ibuprofen

Pain Reliever/Fever Reducer

- Temporary relief from pain, fever and inflammation
- Soothes muscles aches and minor arthritis pain
- Non-drowsy, aspirin-free formula

These quality comfort tablets are formulated and packaged for use in the workplace and meet ANSI Z308.1-2015 standards. They are sealed in tamper-evident packets for safety and convenience.



Pain Reliever/Fever Reducer

- Temporary relief from pain, fever and inflammation
- Soothes muscle aches and minor arthritis pain
- Non-drowsy, aspirin-free formula

250 Tablets • 2 Tablets Per Packet

Manufactured for
American Safety & First Aid
Osceola, IN 46561

Pain Reliever/Fever Reducer

Ibuprofen

Product #1104

250 Tablets • 2 Tablets Per Packet

Drug Facts

Active Ingredient (In Each Tablet).....**Purpose**
Ibuprofen, USP 200 mg (NSAID).....pain reliever/fever reducer
*Nonsteroidal Anti-Inflammatory Drug

Use Temporarily relieves fever and minor aches and pains associated with: • common cold • backache
• headache • toothache • arthritis • muscular aches • menstrual cramps

Warnings

Allergy Alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: •hives •asthma (wheezing) •facial swelling •shock •skin reddening •rash •blisters
If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause severe stomach bleeding. The chance is higher if you: •are age 60 or older •have had stomach ulcers or bleeding problems •take a blood thinning (anticoagulant) or steroid drug •take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others) •have 3 or more alcoholic drinks every day while using this product •take more or for a longer time than directed

Do not use

• If you ever had an allergic reaction to any other pain reliever/fever reducer • right before or after heart surgery

Ask a doctor before use if •stomach bleeding warning applies to you •you have a history of stomach problems, such as heartburn •you have high blood pressure, heart disease, liver cirrhosis, or kidney disease •you are taking a diuretic

Ask a doctor or pharmacist before use if you are •taking any other drug containing an NSAID (prescription or nonprescription) •taking a blood thinning (anticoagulant) drug •under a doctor's care for any serious condition •taking any other drug

When using this product •take with food or milk if stomach upset occurs
• long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if you experience any of the following signs of stomach bleeding: •feel faint •vomit blood •have bloody or black stools •have stomach pain that does not get better •pain gets worse or lasts more than 10 days •fever gets worse or lasts more than 3 days •redness or swelling is present in the painful area •any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. IT IS ESPECIALLY IMPORTANT NOT TO USE IBUPROFEN DURING THE LAST 3 MONTHS OF PREGNANCY UNLESS DEFINITELY DIRECTED TO DO SO BY A DOCTOR BECAUSE IT MAY CAUSE PROBLEMS IN THE UNBORN CHILD OR COMPLICATIONS DURING DELIVERY.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions •do not take more than directed •the smallest effective dose should be used
•do not take longer than 10 days, unless directed by a doctor (see Warnings)

Adults: 1 tablet every 4 to 6 hours while symptoms occur; if pain or fever does not respond to 1 tablet, 2 tablets may be used, but do not exceed 6 tablets in 24 hours unless directed by a doctor.
Children under 12 years: Ask a doctor.

Other Information

•do not use if packet is opened or torn
•Store at 20-25°C (68-77°F) avoid excessive heat above 40°C (104°F)

Inactive Ingredients croscarmellose sodium*, FD&C Blue #2*, FD&C Red #40*, FD&C Yellow #6*, hypromellose, iron oxide red, lactose monohydrate, magnesium stearate, maltodextrin, medium chain triglycerides, microcrystalline cellulose, polydextrose, polyethylene glycol, polyvinyl alcohol, povidone (K-30), silicon dioxide, sodium starch glycolate, starch, stearic acid, talc, titanium dioxide, triacetin
*contains one or more of these ingredients

Questions? 1-866-651-3660 Monday-Friday 8:00am-5:00pm EST



Pain Reliever/Fever Reducer

PULL TO OPEN

Compare Active Ingredients To: Advil®

Do not use if individual packet is open or torn.

V1_10/9/19



IBUPROFEN 100

ibuprofen tablet, coated

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:73598-1102

Route of Administration

ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE K30 (UNII: U725QWY32X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	BROWN	Score	2 pieces
Shape	ROUND	Size	10mm
Flavor		Imprint Code	G2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73598-1102-1	50 in 1 BOX	02/02/2000	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA072096	02/02/2000	

IBUPROFEN 250

ibuprofen tablet, coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73598-1104
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XYI10QM) (IBUPROFEN - UNII:WK2XYI10QM)	IBUPROFEN	200 mg

Inactive Ingredients

Ingredient Name	Strength
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C BLUE NO. 2 (UNII: L06K8R7DQK)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
LACTOSE MONOHYDRATE (UNII: EWQ57Q8I5X)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MEDIUM-CHAIN TRIGLYCERIDES (UNII: C9H2L21V7U)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
POLYDEXTROSE (UNII: VH2XOU12IE)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
POVIDONE K30 (UNII: U725QWY32X)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM STARCH GLYCOLATE TYPE A (UNII: H8AV0SQX4D)	
STARCH, CORN (UNII: O8232NY3SJ)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

Product Characteristics

Color	BROWN	Score	2 pieces
Shape	ROUND	Size	10mm

Flavor		Imprint Code	G2
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73598-1104-1	125 in 1 BOX	02/02/2000	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA072096	02/02/2000	

Labeler - JHK Inc dba American Safety & First Aid (867236309)

Revised: 4/2024

JHK Inc dba American Safety & First Aid