

**ACETAMINOPHEN- acetaminophen tablet**  
**WALMART INC**

-----  
**EXTRA STRENGTH**

**Pain Reliever**

**Acetaminophen 500mg**

**Pain Reliever/Fever Reducer**

**FOR ADULTS**

**Contains No Aspirin**

**Active ingredient (in each caplet)**

Acetaminophen USP, 500 mg

**Purpose**

Pain reliever/fever reducer

**Uses**

- temporarily relieves minor aches and pains due to:
- the common cold
  
- headache
- backache
  
- minor pain of arthritis
- toothache
  
- muscular aches
- premenstrual and menstrual cramps
- temporarily reduces fever

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Allergy alert**

acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are allergic to acetaminophen or any of the inactive ingredients in this product

### **Ask a doctor before use if you have**

liver disease

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

### **Stop use and ask a doctor if**

- pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - new symptoms occur
  - redness or swelling is present
- These could be signs of a serious condition

### **If pregnant or breast-feeding,**

ask a health professional before use

### **Keep out of reach of children**

Keep out of reach of children

### **Overdose warning:**

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

#### **■ do not take more than directed (see overdose warning)**

adults and children 12 years and over

- take 2 caplets every 6 hours while symptoms last
- do not take more than 6 caplets in 24 hours, unless directed by a doctor
- do not use for more than 10 days unless directed by a doctor

children under 12 years

- ask a doctor

## Other information

- store between 20-25°C (68-77°F). See USP Controlled Room Temperature.
- see bottom of the label for lot number and expiration date

## Inactive ingredients

hydroxypropyl methyl cellulose, polyethylene glycol, povidone k-30, pregelatinized starch, sodium starch glycolate, stearic acid

## Questions or comments?

1-888-287-1915

**equat<sup>™</sup>** NDC 79903-296-10  
**EXTRA STRENGTH**  
**Pain Reliever Acetaminophen 500 MG**  
 Compare to Tylenol® Extra Strength active ingredient\*  
**Pain Reliever/Fever Reducer For Adults**  
 Contains No Aspirin  
 Actual Size  
**500 mg EACH**  
**100 CAPLETS\***  
 (\*Capsule-Shaped Tablets)  
 IMPORTANT: READ THE DIRECTIONS AND WARNINGS BEFORE USE  
 TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING  
**Drug Facts**  
**Active ingredient (in each caplet) Purpose**  
 Acetaminophen USP, 500 mg.....Pain reliever/fever reducer  
**Uses**  
 ■ temporarily relieves minor aches and pains due to:  
 ■ the common cold ■ headache  
 ■ backache ■ minor pain of arthritis  
 ■ toothache ■ muscular aches  
 ■ premenstrual and menstrual cramps  
 ■ temporarily reduces fever  
**Warnings**  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take  
 ■ more than 4,000 mg of acetaminophen in 24 hours  
 ■ with other drugs containing acetaminophen  
 ■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** acetaminophen may cause severe skin reactions. Symptoms may include:  
 ■ skin reddening ■ blisters ■ rash  
 ■ if a skin reaction occurs, stop use and seek medical help right away  
**Do not use**  
 ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
 ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product  
 Satisfaction guaranteed - For questions or comments please call 1-888-287-1915  
 94346 37708  
 634803 (CONTINUED ON BACK OF LABEL)  
 200000005675  
 700000004656  
 PLASTIC BOTTLE  
 LOT  
 EXP  
 PEEL HERE

## Inside (adhesive side)

**Drug Facts (continued)**  
 ingredients in this product  
 Ask a doctor before use if you have liver disease  
 Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin  
**Stop use and ask a doctor if**  
 ■ pain gets worse or lasts more than 10 days  
 ■ fever gets worse or lasts more than 3 days  
 ■ new symptoms occur  
 ■ redness or swelling is present  
 These could be signs of a serious condition  
**If pregnant or breast-feeding,** ask a health professional before use  
**Keep out of reach of children**  
**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away, (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.  
**Directions** ■ do not take more than directed (see overdose warning)  
 adults and children 12 years and over  
 ■ take 2 caplets every 6 hours while symptoms last  
 ■ do not take more than 6 caplets in 24 hours, unless directed by a doctor  
 ■ do not use for more than 10 days unless directed by a doctor  
 children under 12 years  
 ■ ask a doctor  
**Other information**  
 ■ store between 20-25°C (68-77°F). See USP Controlled Room Temperature.  
 ■ see bottom of the label for lot number and expiration date  
**Inactive ingredients**  
 hydroxypropyl methyl cellulose, polyethylene glycol, povidone k-30, pregelatinized starch, sodium starch glycolate, stearic acid  
**Questions or comments?**  
 1-888-287-1915

## ACETAMINOPHEN

acetaminophen tablet

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:79903-296
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	500 mg

### Inactive Ingredients

Ingredient Name	Strength
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDW1A)	
<b>POVIDONE K30</b> (UNII: U725QWY32X)	
<b>SODIUM STARCH GLYCOLATE TYPE A POTATO</b> (UNII: 5856J3G2A2)	
<b>HYPROMELLOSE, UNSPECIFIED</b> (UNII: 3NXW29V3WO)	
<b>STARCH, CORN</b> (UNII: O8232NY3SJ)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z 65AP)	

### Product Characteristics

<b>Color</b>	white	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	17mm
<b>Flavor</b>		<b>Imprint Code</b>	G;551
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-296-10	100 in 1 BOTTLE; Type 0: Not a Combination Product	12/03/2024	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/03/2024	

**Labeler** - WALMART INC (051957769)